

Some upper bounds on ordinal-valued Ramsey numbers for colourings of pairs

Leszek Aleksander Kołodziejczyk¹ and Keita Yokoyama²

¹Institute of Mathematics

University of Warsaw

E-mail: lak@mimuw.edu.pl

²School of Information Science

Japan Advanced Institute of Science and Technology

E-mail: y-keita@jaist.ac.jp

November 7, 2018

Abstract

We study Ramsey's theorem for pairs and two colours in the context of the theory of α -large sets introduced by Ketonen and Solovay. We prove that any 2-colouring of pairs from an ω^{300n} -large set admits an ω^n -large homogeneous set. We explain how a formalized version of this bound gives a more direct proof, and a strengthening, of the recent result of Patey and Yokoyama [Adv. Math. 330 (2018), 1034–1070] stating that Ramsey's theorem for pairs and two colours is $\forall\Sigma_2^0$ -conservative over the axiomatic theory RCA_0 (recursive comprehension).

Introduction

The work described in this paper is mostly finite combinatorics. Much of the motivation, on the other hand, comes from logic.

We contribute to the quantitative study of Ramsey's theorem for pairs in a setting where the pairs always come from a finite subset of \mathbb{N} , but the size of the subset is given by a countable ordinal rather than just the finite ordinal specifying its cardinality. More concretely, we use the framework of α -large sets originally due to Ketonen and Solovay [10], in which, for instance:

- a set $X \subseteq \mathbb{N}$ is n -large, for $n \in \mathbb{N}$, exactly if X has at least n elements,
- X is ω -large if $X \setminus \{\min X\}$ is $\min X$ -large, that is, if X has strictly more than $\min X$ elements,
- X is ω^2 -large if $X \setminus \{\min X\}$ can be split into $\min X$ many sets $X_1, \dots, X_{\min X}$ such that $\max X_i < \min X_{i+1}$ and each X_i is ω -large,

The work of the first author is partially supported by grant number 2017/27/B/ST1/01951 of the National Science Centre, Poland.

The work of the second author is partially supported by JSPS KAKENHI (grant numbers 16K17640 and 15H03634) and JSPS Core-to-Core Program (A. Advanced Research Networks), and JAIST Research Grant 2018(Houga).

and so on (for precise definitions, see below). Our main aim is to obtain a good upper bound on the size of a set X guaranteeing that each 2-colouring of $[X]^2$ will have an ω^n -large homogeneous set, for $n \in \mathbb{N}$.

This sort of work can be viewed simply as a specific kind of finite combinatorics: essentially, the study of bounds on Ramsey numbers that happen to take ordinal values rather than finite ones. Among the papers developing Ramsey theory in the context of α -largeness—e.g. [1, 2, 3, 20, 12, 19]—many do in fact focus on the purely combinatorial side of things. However, the original motivation for studying α -largeness was the desire to understand the combinatorial underpinnings of (un)provability in strong axiom systems. For example, the seminal work of [10] showed that the size of a set needed to guarantee the existence of ω -large homogeneous sets for colourings of n -tuples grows extremely fast with n . This provided a combinatorial explanation for the unprovability of a statement known as the Paris-Harrington theorem in Peano Arithmetic.

Our work is also inspired by a question from logic. It follows from a general-purpose result on colourings of n -tuples [2, Theorem 5] that

$$\omega^{\omega^{n \cdot 2}} \rightarrow (\omega^n)_2^2. \quad (1)$$

That is, every 2-colouring of pairs from an $\omega^{\omega^{n \cdot 2}}$ -large set has an ω^n -large homogeneous set. It has been known that determining whether this upper bound is more or less tight would have important consequences for a longstanding open problem about the logical strength of *infinite* Ramsey’s theorem for pairs (see e.g. [17, Question 4.4] or [14, Question 2] for the question and e.g. [5, 6, 7, 4] for some important related work). Recently, Patey and the second author [15] solved that open problem by showing that (1) is *not* tight. However, the argument in [15] was non-constructive and required a detour via infinite combinatorics and forcing; as a consequence, it did not give any specific bound.

Our main theorem here is

$$\omega^{300n} \rightarrow (\omega^n)_2^2. \quad (2)$$

This *is* more or less tight, at least in the sense that it is impossible to get the left-hand side down from $\omega^{O(n)}$ to $\omega^{(1+o(1))n}$ [12]. Moreover, our arguments use only relatively basic finite-combinatorial tools, which means that they can be formalized in axiomatic theories of modest strength. In effect, we obtain a new, significantly more direct proof of the main result of [15]: any simple enough statement provable using infinite Ramsey’s theorem for pairs and two colours can also be proved in the axiomatic theory RCA_0 , which corresponds to a form of “computable mathematics” and (unlike infinite Ramsey’s theorem) is too weak to imply the existence of any non-computable sets. In fact, we also obtain some improvements of that result, which provide additional information concerning the proof-theoretic properties of Ramsey’s theorem.

The paper consists of three sections. In Section 1, we provide the necessary definitions and background. In Section 2, we prove the main theorem. Those two sections involve no logic beyond elementary facts about small infinite ordinals. The connections to logic are explained in Section 3.

1 α -largeness and Ramsey α -largeness

We fix a primitive recursive notation for ordinals below ω^ω by writing them in *Cantor normal form*: $\alpha = \sum_{i < k} \omega^{n_i}$ where $n_i \in \mathbb{N}$ and $n_0 \geq \dots \geq n_{k-1}$.

Let $\alpha = \sum_{i < k} \omega^{n_i}$ and $\beta = \sum_{i < k'} \omega^{m_i}$. We write $\beta \triangleright \alpha$ if $m_{k'-1} \geq n_0$. If $\beta \triangleright \alpha$, we can define the sum of β and α as $\beta + \alpha = \sum_{i < k+k'} \omega^{t_i}$ where $t_i = m_i$ for $i < k'$ and $t_{j+k'} = n_j$ for $j < k$.

In what follows, we only consider sums of this form. We let $\beta > \alpha$ if there is $i \leq k, k'$ such that $n_j = m_j$ for any $j < i$ and $(n_i < m_i$ or $i = k < k')$. By definition, $\beta \supseteq \alpha$ implies $\beta \geq \alpha$.

We write 1 for ω^0 , and $\omega^n \cdot k$ for $\sum_{i < k} \omega^n$. With this notation, one can write $\alpha < \omega^\omega$ as $\alpha = \omega^n \cdot k_n + \dots + \omega^0 \cdot k_0$, and put $\text{MC}(\alpha) = \max\{k_n, \dots, k_0\}$ (MC stands for the *maximal coefficient* of α).

For a given $\alpha < \omega^\omega$ and $m \in \mathbb{N}$, define $0[m] = 0$, $\alpha[m] = \beta$ if $\alpha = \beta + 1$, and $\alpha[m] = \beta + \omega^{n-1} \cdot m$ if $\alpha = \beta + \omega^n$ for some $n \geq 1$. By definition, $m \leq n$ implies $\alpha[m] \leq \alpha[n]$.

The following definition combines a fundamental concept from [10] with a variant from [15].

Definition 1.1 (largeness). Let $\alpha < \omega^\omega$, and let $n, k, m \in \mathbb{N}$.

1. A set $X = \{x_0 < \dots < x_{\ell-1}\} \subseteq_{\text{fin}} \mathbb{N}$ is said to be α -large if $\alpha[x_0] \dots [x_{\ell-1}] = 0$. In other words, any finite set is 0-large, and X is said to be α -large if
 - $X \setminus \{\min X\}$ is β -large if $\alpha = \beta + 1$,
 - $X \setminus \{\min X\}$ is $(\beta + \omega^{n-1} \cdot \min X)$ -large if $\alpha = \beta + \omega^n$.
2. A set $X \subseteq_{\text{fin}} \mathbb{N}$ is said to be RT_k^n - α -large if for any $P : [X]^n \rightarrow k$, there exists $Y \subseteq X$ such that Y is P -homogeneous and α -large.

The above definition of ω^n -largeness causes minor issues if $\min X$ is a very small number – for instance, the set $\{0\}$ ends up being ω^n -large for every n . To avoid this and simplify the notation, we will always consider finite sets $X \subseteq_{\text{fin}} \mathbb{N}$ satisfying $\min X \geq 3$. We will first check several basic properties.

Lemma 1.2. Let $\alpha, \beta < \omega^\omega$ and $m \in \mathbb{N}$. If $\alpha \leq \beta$ and $\text{MC}(\alpha) < m$, then $\alpha[m] \leq \beta[m]$.

Proof. The case $\alpha = \beta$ is trivial, so we assume $\alpha < \beta$. Write $\beta = \beta' + \omega^n$. If $\alpha \leq \beta'$, then $\alpha[m] \leq \beta' \leq \beta[m]$. Otherwise, $n \geq 1$ and there exists $\gamma \triangleleft \beta'$ such that $\alpha = \beta' + \gamma$ and $\gamma < \omega^n$. Since $\text{MC}(\alpha) < m$, we also have $\text{MC}(\gamma) < m$, thus $\gamma < \omega^{n-1} \cdot m = \omega^n[m]$. Therefore, we obtain $\alpha[m] \leq \alpha < \beta' + \omega^n[m] = \beta[m]$. \square

Lemma 1.3. Let $\alpha < \omega^\omega$ and $X, Y \subseteq_{\text{fin}} \mathbb{N}$ where $X = \{x_0 < \dots < x_{\ell-1}\}$, $Y = \{y_0 < \dots < y_{\ell'-1}\}$ for $\ell \leq \ell'$. Assume that $y_i \leq x_i$ for each $i < \ell$ and that X is α -large. Then Y is α -large.

In particular, if X is α -large and $X \subseteq Y$, then Y is α -large.

Proof. We will show the following by induction on i :

for any $i < \ell$, there exists $j_i < \ell$ such that $j_i \geq i$ and $\alpha[y_0] \dots [y_i] = \alpha[x_0] \dots [x_{j_i}]$.

The base case, which corresponds to $i = -1$, is the trivial statement $\alpha = \alpha$.

Assume $\beta := \alpha[y_0] \dots [y_i] = \alpha[x_0] \dots [x_{j_i}]$ and $i + 1 < \ell$. If $\beta = 0$, put $j_{i+1} = \max\{j_i, i + 1\}$. If $\beta = \beta' + 1$, then $\beta[y_{i+1}] = \beta[x_{j_i+1}]$, so put $j_{i+1} = j_i + 1$. Note that x_{j_i+1} must exist, because $\alpha[x_0] \dots [x_{j_i}] = \beta \neq 0 = \alpha[x_0] \dots [x_{\ell-1}]$.

If $\beta = \beta' + \omega^n$ for some $n \geq 1$, then

$$\beta[x_{j_i+1}] = \beta' + \omega^{n-1} \cdot (x_{j_i+1}) = \beta[y_{i+1}] + \omega^{n-1} \cdot (x_{j_i+1} - y_{i+1}).$$

Since $\beta[x_{j_i+1}] \dots [x_{\ell-1}] = 0$, we have $\omega^{n-1} \cdot (x_{j_i+1} - y_{i+1})[x_{j_i+2}] \dots [x_{\ell-1}] = 0$. (Otherwise, $\beta[x_{j_i+1}] \dots [x_{\ell-1}] = \beta[y_{i+1}] + \omega^{n-1} \cdot (x_{j_i+1} - y_{i+1})[x_{j_i+2}] \dots [x_{\ell-1}] > 0$.) Let j_{i+1} be the smallest j such that $\omega^{n-1} \cdot (x_{j_i+1} - y_{i+1})[x_{j_i+2}] \dots [x_j] = 0$. We then have $\beta[x_{j_i+1}] \dots [x_{j_i+1}] = \beta[y_{i+1}]$.

Now, since $j_{\ell-1}$ must equal $\ell - 1$, we have $\alpha[y_0] \dots [y_{\ell-1}] = \alpha[x_0] \dots [x_{\ell-1}] = 0$. \square

For a given α -large set $X = \{x_0 < \dots < x_{\ell-1}\} \subseteq_{\text{fin}} \mathbb{N}$, take the minimum $i < \ell$ such that $\alpha[x_0] \cdots [x_i] = 0$ and define $X \upharpoonright \alpha$ to be the set $\{x_0, \dots, x_i\}$. (Thus, $X \upharpoonright \alpha$ is the smallest α -large initial segment of X .)

Lemma 1.4. *Let $\alpha = \alpha_{k-1} + \dots + \alpha_0 < \omega^\omega$ where $\alpha_{k-1} \supseteq \dots \supseteq \alpha_0$. Then, a set $X \subseteq_{\text{fin}} \mathbb{N}$ is α -large if and only if there is a partition $X = X_0 \sqcup \dots \sqcup X_{k-1}$ such that $\max X_i < \min X_{i+1}$ and X_i is α_i -large.*

Proof. Let $X = \{x_0 < \dots < x_{\ell-1}\}$ be α -large. By Lemma 1.3, we can assume without loss of generality that $X = X \upharpoonright \alpha$. For each $i < k$, let X_i be $(X \upharpoonright (\alpha_i + \dots + \alpha_0)) \setminus (X \upharpoonright (\alpha_{i-1} + \dots + \alpha_0))$. One checks by induction on i that X_i equals $(X \setminus (X_0 \cup \dots \cup X_{i-1})) \upharpoonright \alpha_i$. It follows that $\max X_i < \min X_{i+1}$ and X_i is α_i -large.

Conversely, if $X = X_0 \sqcup \dots \sqcup X_{k-1}$ such that $\max X_i < \min X_{i+1}$ and X_i is α_i -large, put $Y_i = X_i \upharpoonright \alpha_i$. Then, $Y = Y_0 \sqcup \dots \sqcup Y_{k-1}$ is α -large by the definition, and thus X is α -large by Lemma 1.3. \square

In [10], Ketonen and Solovay use α -largeness to analyze the Ramsey-theoretic statement known as the Paris-Harrington principle and to clarify the relationship between the principle and hierarchies of fast growing functions. In the process, they prove the following result concerning RT_k^2 - ω -largeness.

Theorem 1.5 (Ketonen-Solovay [10], Lemma 6.4). *Let $n \geq 2$. If $X \subseteq_{\text{fin}} \mathbb{N}$ is ω^{n+4} -large and $\min X \geq 3$, then it is RT_n^2 - ω -large.*

We will give a new proof of this theorem in Subsection 2.2.

Theorem 1.5 and its generalization to RT_k^m proved in [10] only deal with the question how much α -largeness is guaranteed to imply RT_k^m - ω -largeness, that is, the existence of an ω -large homogeneous set for any given colouring. Our target is a generalization of the case $m = k = 2$ to bounds implying RT_2^2 - ω^n -largeness for larger $n \in \mathbb{N}$. As already mentioned, even though this sort of work is purely combinatorial, much of the motivation comes from the study of the proof-theoretic strength of infinite Ramsey's theorem for pairs. We discuss this in more detail in Section 3.

Our main result is as follows.

Theorem 1.6. *If $X \subseteq_{\text{fin}} \mathbb{N}$ is ω^{300n} -large and $\min X \geq 3$, then X is RT_2^2 - ω^n -large.*

2 Calculation

In this section, we prove Theorem 1.6.

To simplify our calculations, we only consider “sparse enough” finite sets. A set X with $\min X \geq 3$ is said to be *exp-sparse* if for any $x, y \in X$, $x < y$ implies $4^x < y$. More generally, X is said to be α -sparse if for any $x, y \in X$, $x < y$ implies that the interval $(x, y]$ is α -large. Trivially, any subset of an α -sparse set is α -sparse. By an easy calculation, one checks that any ω^3 -sparse set is exp-sparse: $y > 2x$ whenever $(x, y]$ is ω -large, $y > x2^x$ whenever $(x, y]$ is ω^2 -large, and $y > 2^{2^{\dots^x}}$ (where there are x applications of the exponential function) whenever $(x, y]$ is ω^3 -large.

Lemma 2.1. *Let $n, m \in \mathbb{N}$. If $X \subseteq_{\text{fin}} \mathbb{N}$ is $(\omega^{n+m} + 1)$ -large and $\min X \geq 3$, then there exists $Y \subseteq X$ such that Y is ω^n -large and ω^m -sparse. In particular, if $X \subseteq_{\text{fin}} \mathbb{N}$ is $(\omega^{n+3} + 1)$ -large and $\min X \geq 3$, then there exists $Y \subseteq X$ such that Y is ω^n -large and exp-sparse.*

Proof. We will show the following slightly stronger condition by induction on n :

if $X \subseteq_{\text{fin}} \mathbb{N}$ is $(\omega^{n+m} + 1)$ -large and $\min X \geq 3$, then there exists $Y \subseteq X \setminus \{\max X\}$ such that Y is ω^n -large and $Y \cup \{\max X\}$ is ω^m -sparse.

For the case $n = 0$, let X be $(\omega^m + 1)$ -large and take $Y = \{\min X\}$. Then Y is ω^0 -large, i.e. 1-large, and it follows from Lemma 1.3 and the $(\omega^m + 1)$ -largeness of X that $\{\min X, \max X\}$ is ω^m -sparse.

We turn to the case $n \geq 1$. If X is $(\omega^{n+m} + 1)$ -large, then $X \setminus \{\min X\}$ is ω^{n+m} -large, thus there exist X_0, \dots, X_{k-1} such that

- $X = \{\min X, \min(X \setminus \{\min X\})\} \sqcup X_0 \sqcup \dots \sqcup X_{k-1}$,
- $k = \min(X \setminus \{\min X\}) \geq 1 + \min X$,
- $\max X_i < \min X_{i+1}$,
- each X_i is ω^{n+m-1} -large.

Put $x_i = \max X_i$. By the induction hypothesis applied to $\{x_i\} \cup X_{i+1}$ for $0 \leq i \leq k-2$, there exist Y_0, \dots, Y_{k-2} such that $Y_i \subseteq \{x_i\} \cup X_{i+1} \setminus \{x_{i+1}\}$, Y_i is ω^{n-1} -large and $Y_i \cup \{x_{i+1}\}$ is ω^m -sparse. Now we can check that $Y = \{\min X\} \cup Y_0 \cup \dots \cup Y_{k-2}$ is ω^n -large and $Y \cup \{\max X\}$ is ω^m -sparse. \square

The following lemma means that if a large set X is 2-coloured, we can always choose a ‘‘majority’’ colour without losing too much of its largeness. This fact underlies most of the constructions in the core part of our proof, as presented in Subsection 2.1. The lemma follows from the more general [1, Theorem 1], but our proof is very simple and—crucially for our purposes—involves no use of transfinite induction.

Lemma 2.2. *For each $n \in \mathbb{N}$, the following holds.*

1. *If $X = Y_0 \cup Y_1 \subseteq_{\text{fin}} \mathbb{N}$ is $\omega^n \cdot 2$ -large and exp-sparse, then Y_0 is ω^n -large or Y_1 is ω^n -large.*
2. *If $X = Y_0 \cup Y_1 \subseteq_{\text{fin}} \mathbb{N}$ is $\omega^n \cdot (4k)$ -large and exp-sparse, then Y_0 is $\omega^n \cdot k$ -large or Y_1 is $\omega^n \cdot k$ -large.*

Proof. First, we show that 1. implies 2. for each $n \in \mathbb{N}$. If X is $\omega^n \cdot (4k)$ -large, then there exists a partition $X = X_0 \sqcup X_1 \sqcup \dots \sqcup X_{2k-1}$ such that $\max X_i < \min X_{i+1}$ and X_i is $\omega^n \cdot 2$ -large. Then, by 1., at least one of $Y_0 \cap X_i$ and $Y_1 \cap X_i$ is ω^n -large for each $i < 2k$. Depending on which case happens for at least half the i 's, at least one of $Y_0 \cap X$ and $Y_1 \cap X$ must be $\omega^n \cdot k$ -large.

We now show 1., and thus also 2., by induction on n . The case $n = 0$ is trivial, so assume $n \geq 1$. Let $X = Y_0 \cup Y_1 \subseteq_{\text{fin}} \mathbb{N}$ be $\omega^n \cdot 2$ -large and exp-sparse. Take a partition $X = X_0 \sqcup X_1$ so that $\max X_0 < \min X_1$ and X_0, X_1 are both ω^n -large. If $X_0 \subseteq Y_0$ or $X_0 \subseteq Y_1$, we are done. Otherwise, there are $c_0, c_1 \in X_0$ such that $c_0 \in Y_0$ and $c_1 \in Y_1$. Put $c = \max\{c_0, c_1\}$. Then, by exp-sparseness, $4^c < \min X_1$, hence $X_1 \setminus \{\min X_1\}$ is $\omega^{n-1} \cdot (4c)$ -large. By 2. of the induction hypothesis, at least one of $Y_0 \cap X_1$ and $Y_1 \cap X_1$ is $\omega^{n-1} \cdot c$ -large. Thus, at least one of $\{c_0\} \cup (Y_0 \cap X_1) \subseteq Y_0$ and $\{c_1\} \cup (Y_1 \cap X_1) \subseteq Y_1$ is ω^n -large. \square

2.1 The grouping principle

In this subsection, we consider the notion of grouping, introduced in [15, Section 7] as a useful tool in the analysis of Ramsey's theorem for pairs. We will obtain an upper bound on the largeness of a set needed to guarantee the existence of sufficiently large groupings.

Definition 2.3 (grouping). Let $\alpha, \beta < \omega^\omega$. Let $X \subseteq \mathbb{N}$ and let $P : [X]^2 \rightarrow 2$ be a colouring. A finite family (sequence) of finite sets $\langle F_i \subseteq X : i < \ell \rangle$ is said to be an (α, β) -grouping for P if

1. $\forall i < j < \ell \max F_i < \min F_j$,
2. for any $i < \ell$, F_i is α -large,
3. $\{\max F_i : i < \ell\}$ is β -large, and,
4. $\forall i < j < \ell \forall x, x' \in F_i \forall y, y' \in F_j [P(x, y) = P(x', y')]$.

Moreover, $\langle F_i \subseteq X : i < \ell \rangle$ is said to be a *strong* (α, β) -grouping for P if the fourth condition is replaced with

- 4'. $\exists c < 2 \forall i < j < \ell \forall x \in F_i \forall y \in F_j [P(x, y) = c]$.

The intuition is that each F_i is a “group” and that the colour of a pair consisting of representatives of two distinct groups depends only on the groups, not on the representatives. We say that a set $X \subseteq \mathbb{N}$ admits an (α, β) -grouping if for any colouring $P : [X]^2 \rightarrow 2$, there exists an (α, β) -grouping for P . Our target theorem in this subsection is the following.

Theorem 2.4. *Let $n, k \in \mathbb{N}$. If $X \subseteq_{\text{fin}} \mathbb{N}$ is ω^{n+6k} -large and exp-sparse, then X admits an (ω^n, ω^k) -grouping.*

To obtain a grouping, we need to stabilize the colour between elements of any two fixed groups. We first show how to stabilize the colour between one set and each individual element of another set. This will have to be done both “from below” and “from above”.

Lemma 2.5. *Let $X \subseteq_{\text{fin}} \mathbb{N}$ be ω^{n+1} -large and exp-sparse, and let $c \in \mathbb{N}$ such that $4^c \leq \min X$. Then, we have the following.*

1. *For any $W \subseteq_{\text{fin}} \mathbb{N}$ such that $|W| \leq c$ and $\max W < \min X$ and for any colouring $P : [W \cup X]^2 \rightarrow 2$, there exists $Y \subseteq X$ such that Y is ω^n -large and $P(w, y) = P(w, y')$ for any $w \in W$ and $y, y' \in Y$.*
2. *For any $W \subseteq_{\text{fin}} \mathbb{N}$ such that $|W| \leq c$ and $\max X < \min W$ and for any colouring $P : [X \cup W]^2 \rightarrow 2$, there exists $Y \subseteq X$ such that Y is ω^n -large and $P(y, w) = P(y', w)$ for any $w \in W$ and $y, y' \in Y$.*

Proof. We only show 1., as the proof of 2. is virtually identical. Since X is ω^{n+1} -large and $4^c \leq \min X$, we know that $X \setminus \{\min X\}$ is $\omega^n \cdot 4^c$ -large. Put $Y_{-1} = X \setminus \{\min X\}$. Without loss of generality, we may assume that $|W| = c$, so let $\{w_i : i < c\}$ be an enumeration of W . Construct a sequence $Y_0 \supseteq Y_1 \supseteq \dots \supseteq Y_c$ so that Y_{i+1} is $\omega^n \cdot 4^{c-i-1}$ -large and $\forall y, y' \in Y_{i+1} (P(w_i, y) = P(w_i, y'))$. Indeed, Lemma 2.2 guarantees that at least one of $\{y \in Y_i : P(w_i, y) = 0\}$ or $\{y \in Y_i : P(w_i, y) = 1\}$ can be chosen as Y_{i+1} . Take Y_c as the desired set Y . \square

Next, we obtain a constant-length grouping.

Lemma 2.6. *Let $X \subseteq_{\text{fin}} \mathbb{N}$ be ω^{n+3} -large and exp-sparse, and let $d \in \mathbb{N}$ such that $d \leq \min X$. Then, X admits an (ω^n, d) -grouping.*

Proof. Fix a colouring $P : [X]^2 \rightarrow 2$. We will construct an (ω^n, d) -grouping for P .

First, we stabilize the colour from below in the sense of Lemma 2.5. Since $d \leq \min X$, we know that $X \setminus \{\min X\}$ is $\omega^{n+2} \cdot d$ -large. Take a partition $X \setminus \{\min X\} = X_0 \sqcup \dots \sqcup X_{d-1}$ so that $\max X_i < \min X_{i+1}$ and X_i is ω^{n+2} -large. Put $Y_0 = X_0$, and for $i \geq 1$ take $Y_i \subseteq X_i$ so that Y_i is ω^{n+1} -large and $P(x, y) = P(x, y')$ for any $x \in X \cap [0, \max X_{i-1}]$ and any $y, y' \in Y_i$. This can be done using Lemma 2.5.1. with $W = X \cap [d, \max X_{i-1}]$ and $c = \max X_{i-1}$, because $4^{\max X_{i-1}} < \min X_i$ by the exp-sparseness of X . Then, $\langle Y_i : i < d \rangle$ is a family of ω^{n+1} -large sets such that for any $0 \leq i < j < d$ and for any $x \in Y_i, y, y' \in Y_j$, we have $P(x, y) = P(x, y')$.

Now, we stabilize the colour from above. Note that $4^d \leq \min Y_i$ for each $i < d$, because $d \leq \min X < \min Y_i$ and all Y_i are subsets of X which is exp-sparse. Put $Z_{d-1} = Y_{d-1}$, and for $i < d-1$ take $Z_i \subseteq Y_i$ so that Z_i is ω^n -large and $P(z, x) = P(z', x)$ for any $x \in \{\min Y_j : i < j < d\}$ and any $z, z' \in Z_i$. This can be done using Lemma 2.5.2. with $W = \{\min Y_j : i < j < d\}$ and $c = d - i - 1$. Then, $\langle Z_i : i < c \rangle$ is a family of ω^n -large sets, and for any $0 \leq i < j < d$ and any $x, x' \in Z_i, y, y' \in Z_j$, we have $P(x, y) = P(x, \min Y_j) = P(x', \min Y_j) = P(x', y')$. Thus, $\langle Z_i : i < c \rangle$ is an (ω^n, c) -grouping for P . \square

By applying Lemma 2.6 twice, we obtain an ω -length grouping.

Lemma 2.7. *Let $X \subseteq_{\text{fin}} \mathbb{N}$ be ω^{n+6} -large and exp-sparse. Then, X admits an (ω^n, ω) -grouping.*

Proof. Fix a colouring $P : [X]^2 \rightarrow 2$. By Lemma 2.6, since $2 \leq \min X$, there is an $(\omega^{n+3}, 2)$ -grouping $\langle Y_0, Y_1 \rangle$ for P . Again by Lemma 2.6, since $\max Y_0 < \min Y_1$, there is an $(\omega^n, \max Y_0)$ -grouping $\langle Z_i : i < \max Y_0 \rangle$ for P with $Z_i \subseteq Y_1$ for each i . One can easily check that $\langle Y_0, Z_0, \dots, Z_{\max Y_0 - 1} \rangle$ is an (ω^n, ω) -grouping for P . \square

Finally we prove Theorem 2.4 by using the previous lemma repeatedly.

Proof of Theorem 2.4. We prove the statement by induction on k . The case $k = 0$ is trivial, and the case $k = 1$ is Lemma 2.7. Assume that $k \geq 2$ and let $X \subseteq_{\text{fin}} \mathbb{N}$ be ω^{n+6k} -large and exp-sparse. Fix a colouring $P : [X]^2 \rightarrow 2$. By Lemma 2.7, there is an $(\omega^{n+6(k-1)}, \omega)$ -grouping $\langle Y_i : i \leq \ell \rangle$ for P . Since $\{\max Y_i : i \leq \ell\}$ is ω -large, we know that $\ell \geq \max Y_0$. By the induction hypothesis, for each $1 \leq i \leq \ell$ there is an (ω^n, ω^{k-1}) -grouping $\langle Z_j^i : j \leq m_i \rangle$ for P such that $Z_j^i \subseteq Y_i$ for each j . Since $\{\max Z_j^i : j \leq m_i\}$ is ω^{k-1} -large for any $1 \leq i \leq \ell$, the set $\{\max Y_0\} \cup \{\max Z_j^i : j \leq m_i, 1 \leq i \leq \ell\}$ is ω^k -large. One can check that $\langle Y_0, Z_0^1, \dots, Z_{m_1}^1, \dots, Z_0^\ell, \dots, Z_{m_\ell}^\ell \rangle$ is an (ω^n, ω^k) -grouping for P . \square

2.2 Proof of Theorem 1.5

In this subsection, we give a simple proof of Theorem 1.5. The proof is still based on the original idea in [10], but the calculation is simplified. We include the argument to make the paper more self-contained and to facilitate the discussion of axiomatic requirements in Section 3.

For a given $P : [X]^2 \rightarrow n$ and $x \in X$, define the *hereditarily minimal prehomogeneous (h.m.p.h.) sequence* $\sigma_x \in [X]^{<\mathbb{N}}$ as follows:

$$\begin{aligned} \sigma_x(0) &= \min X, \\ \sigma_x(i+1) &= \min\{y \in X : y > \sigma_x(i) \wedge \forall j \leq i P(\sigma_x(j), x) = P(\sigma_x(j), y)\}, \\ &\text{stop this construction when } \sigma_x(i) = x. \end{aligned}$$

One can easily check the following from the definition.

- For any $i < j < k < |\sigma_x|$, $P(\sigma_x(i), \sigma_x(j)) = P(\sigma_x(i), \sigma_x(k))$.
- $\sigma_x(i) = y < x$ if and only if $\sigma_y = \sigma_x \upharpoonright_{i+1} \neq \sigma_x$. In particular, any nonempty initial segment of σ_x has the form σ_y for some $y < x$.

For a given colour $c < n$, let $\text{ho}(\sigma_x, c) = \{\sigma_x(i) : i < |\sigma_x| - 1 \wedge P(\sigma_x(i), x) = c\}$. The set $\text{ho}(\sigma_x, c) \cup \{x\}$ is P -homogeneous with colour c . We let $\text{col}(\sigma_x) = \{c < n : \text{ho}(\sigma_x, c) \neq \emptyset\}$. Clearly, $\sigma_x \subseteq \sigma_y$ implies $\text{col}(\sigma_x) \subseteq \text{col}(\sigma_y)$. For $x \in X \setminus \{\min X\}$, we write σ_x^- to denote the longest initial segment $\sigma_y \subsetneq \sigma_x$ such that $\text{col}(\sigma_y) \subsetneq \text{col}(\sigma_x)$. Note that this definition would not make sense for $x = \min X$, because $\text{col}(\sigma_{\min X}) = \emptyset$.

Lemma 2.8. *Let, $n \geq 2$, $X \subseteq_{\text{fin}} \mathbb{N}$ and let $P : [X]^2 \rightarrow n$ be a colouring. Then we have the following.*

1. For any $m \in \mathbb{N}$, $|\{x \in X : |\sigma_x| \leq m\}| \leq n^m$.
2. For any $x \in X$ and $c \in \text{col}(\sigma_x)$, $\min \text{ho}(\sigma_x, c) \leq \sigma_x^-(|\sigma_x^-| - 1)$.

Proof. By the definition of h.m.p.h. sequences, if $\sigma_y = \sigma_x \widehat{\langle y \rangle}$ and $\sigma_z = \sigma_x \widehat{\langle z \rangle}$, then $P(x, y) \neq P(x, z)$. Thus, for any $x \in X$, there are at most n -many y 's in X such that $y > x$, $\sigma_y \supseteq \sigma_x$ and $|\sigma_y| = |\sigma_x| + 1$. Hence the size of $\{x \in X : |\sigma_x| \leq m\}$ is at most $1 + n + \dots + n^{m-1} \leq n^m$, which gives 1.

For a given $x \in X$, put $y = \max\{\min \text{ho}(\sigma_x, c) : c \in \text{col}(\sigma_x)\}$. Then $\text{col}(\sigma_y) \subsetneq \text{col}(\sigma_x)$. Thus, $\sigma_y \subseteq \sigma_x^-$, and we have 2. \square

Proof of Theorem 1.5. Let $X_0 \subseteq_{\text{fin}} \mathbb{N}$ be ω^{n+4} -large and $\min X_0 \geq 3$. Then one can find a subset $X \subseteq X_0$ which is $\omega^n + 1$ -large, ω^3 -sparse and such that $\min X > n$. Indeed, $X'_0 = X_0 \setminus \{\min X_0\}$ is at least $\omega^{n+3} \cdot 3$ -large. Put $X'_1 = X'_0 \upharpoonright_{\omega^{n+3}}$, $X'_2 = (X'_0 \setminus X'_1) \upharpoonright_{\omega^{n+3}}$ and $X'_3 = (X'_0 \setminus X'_1 \cup X'_2) \upharpoonright_{\omega^{n+3}}$. Note that $|X'_1| > n$. By Lemma 2.1, the set $\{\min X'_2, \max X'_2\}$ is ω^3 -sparse. Moreover, $\{\max X'_2\} \cup X'_3$ is $\omega^{n+3} + 1$ -large, so it contains an ω^n -large ω^3 -sparse subset X'' . We can take $X = \{\min X'_2\} \cup X''$ as the desired set.

Now we show that X chosen as above is RT_n^2 - ω -large by way of contradiction. Assume that $P : [X]^2 \rightarrow n$ is a colouring with no ω -large homogeneous set. Write $X = \{x_0 < \dots < x_{\ell-1}\}$. Let $\sigma_i := \sigma_{x_i}$ be the h.m.p.h. sequence defined by P and x_i . For each $1 \leq d \leq n$, we say that $i < \ell$ is d -critical if $|\text{col}(\sigma_i)| = d$ and for any $j < i$, $\sigma_i^- \neq \sigma_j^-$. For $1 \leq i < \ell$ and $1 \leq d \leq n$, define an ordinal $\gamma_i^d < \omega^n$ as follows. If no $j \leq i$ is d -critical, put $\gamma_i^d = 0$. Otherwise, take the largest d -critical number $j_0 \leq i$ and let $m_{i,1}^d = |\{k \leq i : |\text{col}(\sigma_k)| = d\}|$, $m_{i,2}^d = |\{k \leq i : k \text{ is } (d+1)\text{-critical}\}|$ (where $m_{i,2}^d = 0$ for $d = n$); then put $\gamma_i^d = \omega^{n-d} \cdot (x_{j_0} - m_{i,1}^d - m_{i,2}^d)$.

Claim. *If there is a d -critical number $j \leq i$, then $\gamma_i^d > 0$.*

Proof of Claim. Let $j_0 \leq i$ be the largest d -critical number $\leq i$; since $d \geq 1$, we know that $j_0 > 0$. Note that for any $k \leq i$ such that $|\text{col}(\sigma_k)| = d$, we have $\sigma_k^- = \sigma_j^-$ for some $j \leq j_0$ (if not, there would be a d -critical number bigger than j_0) and therefore also $\sigma_k^- = \sigma_j$ for some $j < j_0$; this implies $\sigma_k^-(|\sigma_k^-| - 1) \leq x_{j_0-1}$. Fix $k \leq i$ such that $|\text{col}(\sigma_k)| = d$. Then, for any $c \in \text{col}(\sigma_k)$, $\min \text{ho}(\sigma_k, c) \leq \sigma_k^-(|\sigma_k^-| - 1) \leq x_{j_0-1}$, where the first inequality follows from Lemma 2.8.2. Since $\text{ho}(\sigma_k, c) \cup \{x_k\}$ is P -homogeneous and thus not ω -large, we have $|\text{ho}(\sigma_k, c) \cup \{x_k\}| \leq x_{j_0-1}$, and hence $|\sigma_k| \leq nx_{j_0-1}$. Therefore, by Lemma 2.8.1, we have $m_{i,1}^d \leq n^{nx_{j_0-1}}$.

If $k, k' \leq i$ are both $d+1$ -critical, then $\sigma_k^- \neq \sigma_{k'}^-$ and $|\text{col}(\sigma_k^-)| = |\text{col}(\sigma_{k'}^-)| = d$. Thus, $m_{i,2}^d \leq m_{i,1}^d \leq n^{nx_{j_0-1}}$. Finally, since X is ω^3 -sparse and $x_{j_0-1} > n$, one can easily check that $x_{j_0} > 2n^{nx_{j_0-1}} \geq m_{i,1}^d + m_{i,2}^d$. This completes the proof of the claim.

Now, define $\gamma_0 = \omega^n$ and $\gamma_i = \gamma_i^1 + \dots + \gamma_i^n$ for $i = 1, \dots, \ell - 1$. Note that 1 is 1-critical, because $|\text{col}(\sigma_1)| = 1$ and $\sigma_1^- = \emptyset$ while σ_0^- does not exist. Thus, by the Claim, $\gamma_i > 0$ for any $i < \ell$.

For $i < \ell - 1$, consider the difference between γ_i and γ_{i+1} . Let $d = |\text{col}(\sigma_{i+1})|$. There are two cases:

- if $i + 1$ is d -critical, then γ_{i+1} is obtained from γ_i by removing one $\omega^{n-(d-1)}$ and adding at most x_{i+1} -many ω^{n-d} 's (note that if $i > 0$, then $d > 1$ and $\gamma_i^{d-1} > 0$ because there must be a $(d-1)$ -critical $j \leq i$),
- if $i + 1$ is not d -critical, then γ_{i+1} is obtained from γ_i simply by removing one ω^{n-d} .

In either case, $\gamma_{i+1} \leq \gamma_i[x_{i+1}]$. Note also that $\text{MC}(\gamma_i) < x_{i+1}$. This lets us check by induction that $\gamma_i \leq \gamma_0[x_1] \dots [x_i]$ for any $1 \leq i < \ell$. Indeed, $\gamma_i \leq \gamma_0[x_1] \dots [x_i]$ and $\text{MC}(\gamma_i) < x_{i+1}$ implies that $\gamma_{i+1} \leq \gamma_i[x_{i+1}] \leq \gamma_0[x_1] \dots [x_i][x_{i+1}]$ by Lemma 1.2. Since $\gamma_0 = (\omega^n + 1)[x_0]$, we have $0 < \gamma_i \leq (\omega^n + 1)[x_0] \dots [x_i]$ for any $i < \ell$. However, $(\omega^n + 1)[x_0] \dots [x_{\ell-1}] = 0$ since X is $\omega^n + 1$ -large. This implies $0 < \gamma_{\ell-1} \leq 0$, which is a contradiction. \square

2.3 Decomposition of Ramsey's theorem for pairs

A colouring $P : [X]^2 \rightarrow 2$ is said to be *transitive* if both $P^{-1}(0)$ and $P^{-1}(1)$ are transitive relations on X . Here $[X]^2$ is formally understood as the set of ordered pairs from X in which the second element is strictly greater than the first: in other words, for a transitive P , if $x < y < z$ and $P(x, y) = P(y, z)$, then $P(x, z)$ must have the same value as well.

Using this notion, RT_2^2 can be decomposed as $\text{RT}_2^2 = \text{EM} + \text{ADS}$ where

- EM: for any colouring $P : [\mathbb{N}]^2 \rightarrow 2$, there exists an infinite set $H \subseteq \mathbb{N}$ such that P is transitive on $[H]^2$,
- ADS: for any transitive colouring $P : [\mathbb{N}]^2 \rightarrow 2$, there exists an infinite set $H \subseteq \mathbb{N}$ such that H is P -homogeneous.

EM and ADS were originally introduced as combinatorial principles about ordered graphs and linear orders, respectively; see [9, 4, 13]. We consider a similar decomposition for RT_2^2 - α -largeness.

Definition 2.9. Let $\alpha < \omega^\omega$.

1. A set $X \subseteq_{\text{fin}} \mathbb{N}$ is said to be EM- α -large if for any colouring $P : [X]^2 \rightarrow 2$, there exists $Y \subseteq X$ such that P is transitive on $[Y]^2$ and Y is α -large.
2. A set $X \subseteq_{\text{fin}} \mathbb{N}$ is said to be ADS- α -large if for any transitive colouring $P : [X]^2 \rightarrow 2$, there exists $Y \subseteq X$ such that Y is P -homogeneous and Y is α -large.

We prove Theorem 1.6 by combining appropriate upper bounds for EM- α -largeness and ADS- α -largeness.

Theorem 2.10. *If $X \subseteq_{\text{fin}} \mathbb{N}$ is ω^{36n} -large and exp-sparse, then it is EM- ω^n -large.*

Note that [15, Lemma 7.2] essentially says that for every n there is an m such that an ω^m -large set is EM- ω^n -large. Theorem 2.10 strengthens this by providing a concrete upper bound on m , which is possible thanks to Theorem 2.4.

Proof. We follow the proof of [15, Lemma 7.2], replacing the use of [15, Lemma 7.1] by Theorem 2.4. It is enough to show that if X is $\omega^{36(n-1)+6}$ -large and exp-sparse then it is EM- ω^n -large. We prove this by induction on n .

The case $n = 1$ is just a weakening of Theorem 1.5. Assume that $n \geq 2$ and let $X \subseteq_{\text{fin}} \mathbb{N}$ be $\omega^{36(n-1)+6}$ -large. Fix $P : [X]^2 \rightarrow 2$. By Theorem 2.4, there exists an $(\omega^{36(n-2)+6}, \omega^6)$ -grouping $\langle Y_i : i \leq \ell \rangle$ for P . Theorem 1.5 applied to the ω^6 -large set $\{\max Y_i : i \leq \ell\}$ gives an $(\omega^{36(n-2)+6}, \omega)$ -subgrouping $\langle Y_{i_j} : j \leq \ell' \rangle$ which is strong, i.e. there is a fixed colour c such that for any x, y from different groups, $P(x, y) = c$. By the induction hypothesis, for each $j \leq \ell'$ there is some $Z_j \subseteq Y_{i_j}$ such that Z_j is ω^{n-1} -large and P is transitive on $[Z_j]^2$. Since $\max Z_0 \leq \max Y_{i_0} \leq \ell'$, the set $H = \{\max Z_0\} \cup \bigcup_{1 \leq j \leq \ell'} Z_j$ is ω^n -large. Moreover, by construction, P is transitive on $[H]^2$. \square

Theorem 2.11. *If $X \subseteq_{\text{fin}} \mathbb{N}$ is ω^{4n+4} -large and $\min X \geq 3$, then it is ADS- ω^n -large.*

Theorem 2.11 is a reformulation of [15, Lemma 4.4]. The proof below is still based on the idea of the original proof.

Proof. Let $X \subseteq_{\text{fin}} \mathbb{N}$ be an ω^{4n+4} -large set with $\min X \geq 3$. Assume towards a contradiction that X is not ADS- ω^n -large. Thus, there is a transitive colouring $P : [X]^2 \rightarrow 2$ without an ω^n -large homogeneous set. Given $x, y \in X$ with $x < y$, we say that an interval $[x, y]$ is (i, α) -long if $P(x, y) = i$ and there exists an α -large set $H \subseteq [x, y] \cap X$ such that $x, y \in H$ and H is P -homogeneous with colour i . Define a new colouring $Q : [X]^2 \rightarrow 4n$ as follows:

$$Q(x, y) = \begin{cases} 4k & \text{if } [x, y] \text{ is } (0, \omega^k)\text{-long but not } (0, \omega^k + 1)\text{-long,} \\ 4k + 1 & \text{if } [x, y] \text{ is } (0, \omega^k + 1)\text{-long but not } (0, \omega^{k+1})\text{-long,} \\ 4k + 2 & \text{if } [x, y] \text{ is } (1, \omega^k)\text{-long but not } (1, \omega^k + 1)\text{-long,} \\ 4k + 3 & \text{if } [x, y] \text{ is } (1, \omega^k + 1)\text{-long but not } (1, \omega^{k+1})\text{-long,} \end{cases}$$

where $0 \leq k < n$. Since there is no ω^n -large P -homogeneous set, Q is well-defined. By Theorem 1.5, there exists an ω -large Q -homogeneous set $\bar{H} \subseteq X$. Write $\bar{H} = \{x_0, \dots, x_m\}$ where $x_0 < \dots < x_m$. By ω -largeness, $m \geq x_0$.

We now claim that $Q(x_0, x_1) \neq Q(x_0, x_m)$, which will contradict the Q -homogeneity of \bar{H} . The proof of the claim splits into four cases depending on $Q(x_0, x_1)$. Consider for instance the case where $[x_0, x_1]$, and thus each of $[x_i, x_{i+1}]$, is $(0, \omega^k + 1)$ -long but not $(0, \omega^{k+1})$ -long. For each $i \leq m - 1$, let H_i be the $\omega^k + 1$ -large Q -homogeneous subset of $[x_i, x_{i+1}]$ whose existence follows from the assumption that $[x_i, x_{i+1}]$ is $(0, \omega^k + 1)$ -long. Let H be $\bigcup_{i < m} H_i$. Note that $x_0 \in H_0$ and that $x_i = \max H_{i-1} = \min H_i$ for $1 \leq i \leq m - 1$; in particular, $H_{i-1} \cap H_i \neq \emptyset$. Thus, by the transitivity of P , the set H is Q -homogeneous with colour 0. Moreover, $m \geq x_0$ and $H = \{x_0\} \cup \bigcup_{i < m} (H_i \setminus \{x_i\})$ imply that H is ω^{k+1} -large. Hence, $[x_0, x_m]$ is $(0, \omega^{k+1})$ -long, which implies $Q(x_0, x_1) \neq Q(x_0, x_m)$. The other cases are similar or easier. \square

Proof of Theorem 1.6. We show that if $X \subseteq_{\text{fin}} \mathbb{N}$ is $(\omega^{(4n+4) \cdot 36+3} + 1)$ -large, then it is RT_2^2 - ω^n -large. Fix a colouring $P : [X]^2 \rightarrow 2$. First, using Lemma 2.1, take $X_0 \subseteq X$ which is $\omega^{(4n+4) \cdot 36}$ -large and exp-sparse. Next, using Theorem 2.10, take $X_1 \subseteq X_0$ such that X_1 is ω^{4n+4} -large and P is transitive on $[X]^2$. Finally, Theorem 2.11 gives $Y \subseteq X_1$ which is ω^n -large and P -homogeneous. \square

Remark. One may obtain slightly better bounds for some of the theorems/lemmas above. For example, in Lemma 2.6, if $d = 2$ then we only need X to be ω^{n+2} -large, because we only need to shrink X_1 in the first stage of the proof and $Y_0 = X_0$ in the second stage. This could actually

be used to obtain a slightly better upper bound (ω^{n+5k} -largeness) in Theorem 2.4 but such small improvements are not particularly important from our perspective.

On the other hand, the bound in Theorem 2.4 cannot be reduced to $\omega^{n+o(n)}$ -largeness. Indeed, Kotlarski et al. [12, Theorem 5.4] showed that if a set X is RT_2^2 - ω^n -large, then it is ω^{2n} -large.

3 Finite consequences of Ramsey’s theorem for pairs

In this section, we explain the relevance of Theorem 1.6 to logic, or more specifically to proof theory. Ramsey-theoretic principles are well-known to display interesting behaviour with respect to provability in axiomatic theories. For instance, the already mentioned Paris-Harrington principle, which states:

“for every $n, \ell \in \mathbb{N}$ there exists a finite set $X \subseteq_{\text{fin}} \mathbb{N}$ such that $\min X \geq \ell$ and X is RT_2^n - ω -large”

is unprovable in Peano Arithmetic (see e.g. [8]). In contrast, it was recently proved in [15] that (infinite) Ramsey’s theorem for pairs and two colours is in a certain sense proof-theoretically “tame”. Theorem 1.6 makes it possible to give a more direct proof of that result and, in fact, to strengthen it.

To understand the proofs in this section, the reader will need some familiarity with axiomatic theories of first- and second-order arithmetic and their models—see [18, 8] for details. The following very brief review will hopefully suffice for understanding the statements of the results. The *language of second-order arithmetic* has two types of variables: *first-order* variables x, y, z, \dots or k, ℓ, n, \dots to stand for natural numbers (which can also be used to code other finite objects, such as finite subsets of \mathbb{N}) and *second-order* variables to stand for subsets of \mathbb{N} (which can also be used to code relations on \mathbb{N}). A formula in this language is Σ_n^0 if it has no second order quantifiers and consists of at most n first-order quantifiers (beginning with \exists) followed by a formula in which all quantifiers have to be *bounded*, i.e. of the form $\exists x < y$ or $\forall x < y$. The dual class of formulas beginning with \forall is called Π_n^0 , while $\forall \Sigma_n^0$ stands for the class of formulas consisting of universal (possibly second-order) quantifiers followed by a Σ_n^0 formula. RCA_0 is an axiomatic theory in this language which has: (a) some basic axioms specifying that \mathbb{N} is a discrete ordered semiring, (b) the Δ_1^0 -*comprehension* axiom, which states that for every decidable property R of natural numbers (as given by an appropriate syntax) the set $\{n \in \mathbb{N} : R(n)\}$ exists, and (c) the Σ_1^0 -induction axiom, which allows the use of mathematical induction for any property expressed by a Σ_1^0 formula (which in fact means: for any recursively enumerable property). RCA_0 may be viewed as embodying the methods of “computable mathematics”. For each n , any $\forall \Sigma_n^0$ statement provable in RCA_0 is provable in the weaker theory IS_1^0 , which only has axioms of type (a), (c). *EFA* (*Elementary Function Arithmetic*) is an even weaker theory in which mathematical induction can only be used for properties defined without using any unbounded quantifiers; to counteract this weakness, *EFA* has to include an additional axiom guaranteeing the basic properties of the exponential function on \mathbb{N} , including the fact that 2^n exists for every $n \in \mathbb{N}$.

The main result of [15] concerns the theory $\text{WKL}_0 + \text{RT}_2^2$, which is obtained by adding a weak version of König’s Lemma and a natural statement of Ramsey’s theorem for pairs and two colours to RCA_0 .

Theorem. [15, Theorem 7.4] $\text{WKL}_0 + \text{RT}_2^2$ is $\forall \Sigma_2^0$ -conservative over RCA_0 . That is, each $\forall \Sigma_2^0$ statement provable in $\text{WKL}_0 + \text{RT}_2^2$ is already provable in RCA_0 .

The combinatorial core of the proof of this theorem in [15] is contained in the following result about α -largeness. Here and below, ordinals smaller than ω^ω are represented in RCA_0 by letting the number coding $\langle n_0, \dots, n_{k-1} \rangle$ stand for $\sum_{i < k} \omega^{n_i}$.

Proposition. [15, Proposition 7.7] *For every natural number n there exists a natural number m such that RCA_0 proves: for every $X \subseteq_{\text{fin}} \mathbb{N}$ with $\min X \geq 3$, if X is ω^m -large, then X is RT_2^2 - ω^n -large.*

However, the proof of [15, Theorem 7.4] does not work with [15, Proposition 7.7] directly, but instead makes use of an intermediate notion of “density”. Moreover, even though [15, Proposition 7.7] is a statement of finite combinatorics, its proof involves a major detour through an infinitary principle (cf. [15, Section 6]). Our proof of Theorem 1.6 is considerably more direct and it is readily seen to give the following stronger version of [15, Proposition 7.7]:

Corollary 3.1. RCA_0 (and, in fact, the weaker theory EFA) *proves the following: for every $n \in \mathbb{N}$ and every $X \subseteq_{\text{fin}} \mathbb{N}$ with $\min X \geq 3$, if X is ω^{300n} -large, then X is RT_2^2 - ω^n -large.*

Proof. An inspection of the arguments in Sections 1 and 2 (including the proof of Theorem 1.5 as presented in Subsection 2.2) reveals that they only make use of elementary manipulations of finite combinatorial objects such as finite sets, finite trees and Cantor Normal Forms, and of the usual principle of mathematical induction applied to properties that can be expressed using bounded quantifiers, possibly with exponentially large bounds. These tools are available within EFA . (A different proof of Theorem 1.5 in EFA was recently given by Pelupecy [16].)

Crucially, none of the arguments involve transfinite induction up to ω^ω (which is not available in RCA_0) or mathematical induction for Σ_1^0 or Π_1^0 properties whose definitions require unbounded quantifiers (this would be available in RCA_0 but not in EFA). Regarding the second point, note that all apparent uses of Π_1^0 -induction—as in, for instance, the proof of Theorem 2.4, where we seem to be using induction for a statement quantifying over all $X \subseteq_{\text{fin}} \mathbb{N}$ —can be replaced by bounded induction: for any given X , the universal quantifier in the induction property can be restricted to range over subsets of X . \square

The extra strength provided by Corollary 3.1 can be used to obtain a strengthening of [15, Theorem 7.4], by means of a relatively simple proof that avoids the concept of density. To express the strengthening, let $\text{WO}(\alpha)$, for $\alpha < \omega^\omega$, denote the statement that there is no infinite descending sequence of ordinals starting from α . The following lemma lists some basic properties of ordinals below ω^ω provable within RCA_0 . The properties are well-known and their easy proofs seem to be part of the folklore.

Lemma 3.2. *The following are provable within RCA_0 .*

1. *For any $\alpha < \omega^\omega$, $\text{WO}(\alpha)$ if and only every set of ordinals smaller than α has a minimum element.*
2. *For any $\alpha < \omega^\omega$, $\text{WO}(\alpha)$ if and only if any infinite set contains an α -large subset.*
3. *For any $m \in \mathbb{N}$, $\text{WO}(\omega^m)$ implies $\text{WO}(\omega^{2^m})$.*

In contrast, RCA_0 is unable to prove “ $\text{WO}(\alpha)$ holds for every $\alpha < \omega^\omega$ ”.

Theorem 3.3. $\text{WKL}_0 + \text{RT}_2^2$ *is conservative over RCA_0 with respect to sentences of the form:*

$$\forall \alpha < \omega^\omega (\text{WO}(\alpha) \rightarrow \varphi(\alpha))$$

where φ is $\forall\Sigma_2^0$.

Note that the class of sentences considered in Theorem 3.3 is strictly larger than the one in [15, Theorem 7.4] because $\text{WO}(\alpha)$ is not a Σ_3^0 statement (it is in fact $\forall\Sigma_2^0$).

Proof. (In this argument, we follow the notational conventions of [15], using the symbol ω to denote the smallest infinite ordinal as formalized in RCA_0 and reserving ω for the set of actual (standard) natural numbers.)

Let $\varphi(\alpha) \equiv \forall X \forall x \exists y \forall z \varphi_0(X[z], x, y, z, \alpha)$, where φ_0 is Σ_0^0 , be a $\forall\Sigma_2^0$ -formula such that RCA_0 does not prove $\forall \alpha < \omega^\omega (\text{WO}(\alpha) \rightarrow \varphi(\alpha))$. Take a countable nonstandard model $(M, S) \models \text{RCA}_0 + \exists \alpha < \omega^\omega (\text{WO}(\alpha) \wedge \neg \varphi(\alpha))$. There exist $A \in S$ and $a, \alpha \in M$ such that

$$(M, S) \models \alpha < \omega^\omega \wedge \text{WO}(\alpha) \wedge \forall y \exists z \neg \varphi_0(A[z], a, y, z, \alpha).$$

Take some $c \in M \setminus \{0\}$ such that $\alpha < \omega^c$ and $\text{WO}(\omega^c)$ holds in (M, S) . (If $\alpha = \omega^{c_0} + \beta$, then Lemma 3.2 part 3. lets us take $c := c_0 + 1$.) Also take some $b \in M$ which is greater than each of a, c , and the code for α . Use primitive recursion in (M, S) to define a sequence $\langle x_i : i \in M \rangle$ such that $x_0 = b$ and $x_{i+1} = \min\{x > x_i : \forall y < x_i \exists z < x \neg \varphi_0(A[z], a, y, z, \alpha)\}$. By Δ_1^0 -comprehension in (M, S) , the set $Y = \{x_i : i \in M\}$ belongs to S . Moreover, Y is infinite in (M, S) .

By Lemma 3.2 parts 2. and 3., every infinite set contains an ω^{nc} -large finite subset for each $n \in \omega$. It follows that Y has an ω^{nc} -large M -finite subset for each $n \in \omega$. By overspill, there exists an M -finite set $X \subseteq Y$ which is $\omega^{300^d c}$ -large for some $d \in M \setminus \omega$.

Let $\{E_i\}_{i \in \omega}$ be an enumeration of all M -finite sets which are not ω^c -large, and $\{P_i\}_{i \in \omega}$ be an enumeration of all M -finite functions from $[0, \max X]^2$ to 2. We will construct an ω -length sequence of M -finite sets $X = X_0 \supseteq X_1 \supseteq \dots$ such that for each $i \in \omega$, the set X_i is $\omega^{300^{d-i} c}$ -large, the colouring P_i is constant on $[X_{2i+1}]^2$, and $[\min X_{2i+2}, \max X_{2i+2}] \cap E_i = \emptyset$.

To achieve this, we do the following for each $i \in \omega$. At stage $2i + 1$ of the construction, we take $X_{2i+1} \subseteq X_{2i}$ such that P_i is constant on $[X_{2i+1}]^2$. Assuming X_{2i} was $\omega^{300^{d-2i} c}$ -large, Corollary 3.1 lets us choose X_{2i+1} so that it is $\omega^{300^{d-2i-1} c}$ -large. Then, at stage $2i + 2$, consider the colouring $Q: [X_{2i+1}]^2 \rightarrow 2$ such that $Q(x, y) = 0$ if and only if $E_i \cap [x, y] = \emptyset$. Again by Corollary 3.1, we take $X_{2i+2} \subseteq X_{2i+1}$ such that Q is constant on $[X_{2i+2}]^2$ and X_{2i+2} is $\omega^{300^{d-2i-2} c}$ -large. X_{2i+2} is in particular $(\omega^c + 1)$ -large, so if the colour of Q on $[X_{2i+2}]^2$ was 1, then by Lemma 1.3 the set E_i would be ω^c -large. Therefore, the colour of Q on $[X_{2i+2}]^2$ must be 0, which implies $[\min X_{2i+2}, \max X_{2i+2}] \cap E_i = \emptyset$.

Now, let $I = \sup\{\min X_i : i \in \omega\} \subseteq_e M$. The even-numbered stages of our construction ensure that I is a cut in M and that $X_j \cap I$ is unbounded in I for each $j \in \omega$ (consider the case where E_i is a singleton set). They also ensure that any set $E \in \text{Cod}(M/I)$ which is unbounded in I has an ω^c -large subset. To see this, assume E has no ω^c -large subset and take an M -finite set \hat{E} such that $E = \hat{E} \cap I$. By overspill, there exists $e \in M \setminus I$ such that $\hat{E} \cap [0, e]$ has no ω^c -large subset, but then $\hat{E} \cap [0, e] = E_i$ for some $i \in \omega$ and so by construction $E = \hat{E} \cap I = E_i \cap I$ must be bounded in I .

It follows in particular that I is a semi-regular cut—that is, for every $e \in I$, any $E \in \text{Cod}(M/I)$ which is unbounded in I has an M -finite subset with at least e elements. By standard arguments, this implies $(I, \text{Cod}(M/I)) \models \text{WKL}_0$. Thus, by Lemma 3.2 part 2., we also get $(I, \text{Cod}(M/I)) \models \text{WO}(\omega^c)$.

On the other hand, the odd-numbered stages ensure that $(I, \text{Cod}(M/I)) \models \text{RT}_2^2$. To see this, let $P: [I]^2 \rightarrow 2$ be a function in $\text{Cod}(M/I)$. Then $P = P_i \cap I$ for some $i \in \omega$. Hence P is constant on $[X_{2i+1} \cap I]^2$, and $X_{2i+1} \cap I \in \text{Cod}(M/I)$ is an infinite set in I .

Finally, since $X \cap I$ is unbounded in I , so is $Y \cap I$. Thus, we have

$$(I, \text{Cod}(M/I)) \models \forall y \exists z \neg \varphi_0((A \cap I)[z], a, y, z, \alpha),$$

and hence $(I, \text{Cod}(M/I)) \models \neg \varphi(\alpha)$. We have $(I, \text{Cod}(M/I)) \models \text{WO}(\alpha)$ because $\alpha < \omega^c$. Therefore, $\text{WKL}_0 + \text{RT}_2^2$ does not prove $\forall \alpha < \omega^\omega (\text{WO}(\alpha) \rightarrow \varphi(\alpha))$. \square

The following consequence of Theorem 3.3 states, intuitively speaking, that RT_2^2 does not imply any new closure properties of ordinals below ω^ω compared to RCA_0 .

Corollary 3.4. *For any primitive recursive function $p: \omega^\omega \rightarrow \omega^\omega$ (defined on codes of ordinals), if $\text{RT}_2^2 + \text{WKL}_0$ proves*

$$\forall \alpha < \omega^\omega (\text{WO}(\alpha) \rightarrow \text{WO}(p(\alpha))),$$

then RCA_0 proves the same statement.

As a special case, $\text{RT}_2^2 + \text{WKL}_0$ does not prove $\forall x (\text{WO}(\omega^x) \rightarrow \text{WO}(\omega^{2^x}))$, as this is not provable within RCA_0 . (Note, though, that already the model constructed in the proof of Theorem 3.3 satisfies $\text{WO}(\omega^c) \wedge \neg \text{WO}(\omega^{2^c})$.)

Another strengthening of [15, Theorem 7.4] – in fact, the original motivation for Corollary 3.1 – concerns proof lengths. Corollary 3.1 can be used to obtain the theorem below, which states that $\text{WKL}_0 + \text{RT}_2^2$ has no significant proof speedup for proofs of $\forall \Sigma_2^0$ sentences over RCA_0 . This answers Question 9.5 of [15] in the negative.

Theorem 3.5. *There is a polynomial-time computable mapping which, given a proof p of a $\forall \Sigma_2^0$ sentence φ in $\text{WKL}_0 + \text{RT}_2^2$ as input, returns a proof p' of φ in RCA_0 as output. In particular, the size of p' is at most polynomially larger than the size of p .*

Proving Theorem 3.5 requires a more extensive development of the logical framework. The proof will be provided in the forthcoming paper [11].

References

- [1] Teresa Bigorajska and Henryk Kotlarski. A partition theorem for α -large sets. *Fund. Math.*, 160(1):27–37, 1999.
- [2] Teresa Bigorajska and Henryk Kotlarski. Some combinatorics involving ξ -large sets. *Fund. Math.*, 175(2):119–125, 2002.
- [3] Teresa Bigorajska and Henryk Kotlarski. Partitioning α -large sets: some lower bounds. *Trans. Amer. Math. Soc.*, 358(11):4981–5001, 2006.
- [4] Andrey Bovykin and Andreas Weiermann. The strength of infinitary Ramseyan principles can be accessed by their densities. *Ann. Pure Appl. Logic*, 168(9):1700–1709, 2017.
- [5] Peter A. Cholak, Carl G. Jockusch, and Theodore A. Slaman. On the strength of Ramsey’s theorem for pairs. *The Journal of Symbolic Logic*, 66(1):1–15, 2001.
- [6] C. T. Chong, Theodore A. Slaman, and Yue Yang. The metamathematics of Stable Ramsey’s Theorem for Pairs. *J. Amer. Math. Soc.*, 27(3):863–892, 2014.

- [7] C. T. Chong, Theodore A. Slaman, and Yue Yang. The inductive strength of Ramsey's Theorem for Pairs. *Adv. Math.*, 308:121–141, 2017.
- [8] Petr Hájek and Pavel Pudlák. *Metamathematics of First-Order Arithmetic*. Springer-Verlag, Berlin, 1993. XIV+460 pages.
- [9] Denis R. Hirschfeldt and Richard A. Shore. Combinatorial principles weaker than Ramsey's theorem for pairs. *J. Symb. Log.*, 72:171–206, 2007.
- [10] Jussi Ketonen and Robert Solovay. Rapidly growing Ramsey functions. *Ann. of Math. (2)*, 113(2):267–314, 1981.
- [11] Leszek Aleksander Kołodziejczyk, Tin Lok Wong, and Keita Yokoyama. Ramsey's theorem for pairs, collection, and proof size. In preparation.
- [12] Henryk Kotlarski, Bożena Piekart, and Andreas Weiermann. More on lower bounds for partitioning α -large sets. *Ann. Pure Appl. Logic*, 147:113–126, 2007.
- [13] Manuel Lerman, Reed Solomon, and Henry Towsner. Separating principles below Ramsey's theorem for pairs. *J. Math. Log.*, 13(02):1350007, 2013.
- [14] A. Montalbán. Open questions in reverse mathematics. *Bull. Symb. Log.*, 17(3):431–454, 2011.
- [15] Ludovic Patey and Keita Yokoyama. The proof-theoretic strength of Ramsey's theorem for pairs and two colors. *Adv. Math.*, 330:1034–1070, 2018.
- [16] Florian Pelupessy. On α -largeness and the Paris-Harrington principle in RCA and RCA_0^* , 2016. Available at <https://arxiv.org/abs/1611.08988>.
- [17] David Seetapun and Theodore A. Slaman. On the strength of Ramsey's theorem. *Notre Dame Journal of Formal Logic*, 36(4):570–582, 1995.
- [18] Stephen G. Simpson. *Subsystems of Second Order Arithmetic*. Perspectives in Mathematical Logic. Springer-Verlag, 1999. XIV + 445 pages; Second Edition, Perspectives in Logic, Association for Symbolic Logic, Cambridge University Press, 2009, XVI+ 444 pages.
- [19] Michiel De Smet and Andreas Weiermann. Partitioning α -large sets for $\alpha < \varepsilon_\omega$. Available at <http://arxiv.org/abs/1001.2437>.
- [20] Andreas Weiermann. A classification of rapidly growing Ramsey functions. *Proc. Amer. Math. Soc.*, 132(2):553–561, 2004.