

# Completion of real Johnson-Wilson theory gives fixed points of Morava $E$ -theory

Maia Averett

Mills College

CAT '09  
Warsaw, Poland  
July 2009

# Two cohomology theories

Fix prime  $p = 2$ .

- **Johnson-Wilson theory**  $E(n)$ : Landweber exact theory with

$$E(n)_* = \mathbb{Z}_{(2)}[v_1, \dots, v_{n-1}, v_n^\pm], \quad |v_i| = 2(2^i - 1)$$

- **Morava  $E$ -theory**  $E_n$ : Landweber exact theory with

$$(E_n)_* = W(\mathbb{F}_{2^n})[[u_1, \dots, u_{n-1}]] [u^\pm], \quad |u_i| = 0, |u| = 2$$

- Related by completion and homotopy fixed points:

$$\widehat{E(n)} = L_{K(n)} E(n), \quad E_n(\text{Gal}) = E_n^{hG}$$

$$\widehat{E(n)} \simeq E_n(\text{Gal})$$

$$\widehat{E(n)}_* = (E(n)_*)_{I_n}^\wedge = \widehat{\mathbb{Z}}_2[[v_1, \dots, v_{n-1}]] [v_n^\pm]$$

# A natural question

We have  $\widehat{E}(n) \simeq E_n(\text{Gal})$  and...

$\mathbb{Z}/2$  acts on  $\widehat{E}(n)$

Complex conjugation action

$\mathbb{Z}/2$  acts on  $E_n(\text{Gal})$

Action of the subgroup of  
Morava stabilizer group  
generated by the formal inverse

# A natural question

We have  $\widehat{E}(n) \simeq E_n(\text{Gal})$  and...

$\mathbb{Z}/2$  acts on  $\widehat{E}(n)$

Complex conjugation action

$\mathbb{Z}/2$  acts on  $E_n(\text{Gal})$

Action of the subgroup of  
Morava stabilizer group  
generated by the formal inverse

**Natural question: Are these actions related?**

# A natural question

We have  $\widehat{E}(n) \simeq E_n(\text{Gal})$  and...

$\mathbb{Z}/2$  acts on  $\widehat{E}(n)$

Complex conjugation action

$\mathbb{Z}/2$  acts on  $E_n(\text{Gal})$

Action of the subgroup of  
Morava stabilizer group  
generated by the formal inverse

**Natural question: Are these actions related?**

**YES**

# A natural question

We have  $\widehat{E}(n) \simeq E_n(\text{Gal})$  and...

$\mathbb{Z}/2$  acts on  $\widehat{E}(n)$

Complex conjugation action

$\mathbb{Z}/2$  acts on  $E_n(\text{Gal})$

Action of the subgroup of  
Morava stabilizer group  
generated by the formal inverse

**Natural question: Are these actions related?**

**YES**

First a little more background...

# Real theories

Complex conjugation action on  $E(n)$  arises in context of Real theories ( $\mathbb{Z}/2$ -equivariant  $RO(\mathbb{Z}/2)$ -graded)

- Atiyah, 1966: Real  $K$ -theory  $\mathbb{K}\mathbb{R}$

$$\mathbb{K}\mathbb{R}(X) = G \left\{ \begin{array}{l} \text{cplx v.b. } \pi : E \rightarrow X \\ \left. \begin{array}{l} E, X \text{ } \mathbb{Z}/2\text{-spaces} \\ \text{antilin. on fibers, } \pi \text{ equiv} \end{array} \right\} \end{array} \right.$$

# Real theories

Complex conjugation action on  $E(n)$  arises in context of Real theories ( $\mathbb{Z}/2$ -equivariant  $RO(\mathbb{Z}/2)$ -graded)

- Atiyah, 1966: Real  $K$ -theory  $\mathbb{K}R$

$$\mathbb{K}R(X) = G \left\{ \begin{array}{l} \text{cplx v.b. } \pi : E \rightarrow X \\ \left. \begin{array}{l} E, X \text{ } \mathbb{Z}/2\text{-spaces} \\ \text{antilin. on fibers, } \pi \text{ equiv} \end{array} \right\} \end{array} \right\}$$

- Landweber, 1967: Real cobordism  $\mathbb{M}R$

Uses  $\mathbb{Z}/2$ -action of complex conjugation on  $BU(k)$ .

# Real theories

Complex conjugation action on  $E(n)$  arises in context of Real theories ( $\mathbb{Z}/2$ -equivariant  $RO(\mathbb{Z}/2)$ -graded)

- Atiyah, 1966: Real  $K$ -theory  $\mathbb{K}R$

$$\mathbb{K}R(X) = G \left\{ \begin{array}{l} \text{cplx v.b. } \pi : E \rightarrow X \\ \left. \begin{array}{l} E, X \text{ } \mathbb{Z}/2\text{-spaces} \\ \text{antilin. on fibers, } \pi \text{ equiv} \end{array} \right\} \end{array} \right\}$$

- Landweber, 1967: Real cobordism  $\mathbb{M}R$

Uses  $\mathbb{Z}/2$ -action of complex conjugation on  $BU(k)$ .

- Araki, 1978: Defined  $\mathbb{B}PR$  using a Quillen idempotent argument

# Real theories

Complex conjugation action on  $E(n)$  arises in context of Real theories ( $\mathbb{Z}/2$ -equivariant  $RO(\mathbb{Z}/2)$ -graded)

- Atiyah, 1966: Real  $K$ -theory  $\mathbb{K}\mathbb{R}$

$$\mathbb{K}\mathbb{R}(X) = G \left\{ \begin{array}{l} \text{cplx v.b. } \pi : E \rightarrow X \\ \left. \begin{array}{l} E, X \text{ } \mathbb{Z}/2\text{-spaces} \\ \text{antilin. on fibers, } \pi \text{ equiv} \end{array} \right\} \end{array} \right\}$$

- Landweber, 1967: Real cobordism  $\mathbb{M}\mathbb{R}$

Uses  $\mathbb{Z}/2$ -action of complex conjugation on  $BU(k)$ .

- Araki, 1978: Defined  $\mathbb{B}\mathbb{P}\mathbb{R}$  using a Quillen idempotent argument

- Hu & Kriz, 2001: Defined  $\mathbb{K}\mathbb{R}(n)$  and  $\mathbb{E}\mathbb{R}(n)$  as  $\mathbb{M}\mathbb{R}$ -modules

# Kitchloo and Wilson's real Johnson-Wilson theory

Real theory  $\mathbb{E} \rightsquigarrow$  naïve  $\mathbb{Z}/2$ -equivariant theory  $\mathbb{E}_{\{e\}}$

- $\mathbb{K}R_{\{e\}} = KU$
- $\mathbb{M}R_{\{e\}} = MU$
- $\mathbb{E}R(n)_{\{e\}} = E(n)$

# Kitchloo and Wilson's real Johnson-Wilson theory

Real theory  $\mathbb{E} \rightsquigarrow$  naïve  $\mathbb{Z}/2$ -equivariant theory  $\mathbb{E}_{\{e\}}$

- $\mathbb{K}R_{\{e\}} = KU$
- $\mathbb{M}R_{\{e\}} = MU$
- $\mathbb{E}R(n)_{\{e\}} = E(n)$

Taking homotopy fixed points gives new theories:

- $KU^{h\mathbb{Z}/2} = KO$
- $E(n)^{h\mathbb{Z}/2} = ER(n)$ , Kitchloo and Wilson's "real Johnson-Wilson"

# Kitchloo and Wilson's real Johnson-Wilson theory

The  $ER(n)$  are higher real  $K$ -theories.

$$E(1) = KU_{(2)} \quad ER(1) = KO_{(2)}$$

Kitchloo-Wilson: There is a fibration

$$\Sigma^{\lambda(n)} ER(n) \xrightarrow{x(n)} ER(n) \rightarrow E(n)$$

that reduces when  $n = 1$  to the classical fibration

$$\Sigma KO_{(2)} \xrightarrow{\eta} KO_{(2)} \rightarrow KU_{(2)}$$

Makes computations feasible (Bockstein spectral sequence).

$$\lambda(n) = 2^{2n+1} - 2^{n+2} + 1$$

# Bockstein Spectral Sequence

The Bockstein spectral sequence arising from the fibration

$$\Sigma^{\lambda(n)} ER(n) \xrightarrow{x(n)} ER(n) \rightarrow E(n)$$

can be used to compute the coefficients of  $ER(n)$ .

# Bockstein Spectral Sequence

The Bockstein spectral sequence arising from the fibration

$$\Sigma^{\lambda(n)} ER(n) \xrightarrow{x(n)} ER(n) \rightarrow E(n)$$

can be used to compute the coefficients of  $ER(n)$ .

$$ER(n)_* = \mathbb{Z}_{(2)}[\hat{v}_k(l) \mid 0 \leq k < n, l \in \mathbb{Z}][x, v_n^{\pm 2^{n+1}}]/J$$

$J$  is the ideal generated by the relations

$$\hat{v}_0(0) = 2 \quad x^{2^{k+1}-1} \hat{v}_k(l) = 0$$

$$|x| = \lambda(n) = 2^{2n+1} - 2^{n+2} + 1$$

$$ER(n)_* \rightarrow E(n)_* \text{ sends } \hat{v}_k(l) \mapsto v_k v_n^{-(2^k-1)(2^n-1)+l2^{k+1}(2^n-1)}$$

# Completed Johnson-Wilson

- Bousfield localization gives a completion  $E(n) \rightarrow \widehat{E(n)} = L_{K(n)}E(n)$  such that

$$E(n)^* \rightarrow \widehat{E(n)}^*$$

is  $I_n$ -adic completion for  $I_n = (v_0, \dots, v_{n-1})$ .

$$\mathbb{Z}_{(2)}[v_1, \dots, v_{n-1}, v_n^\pm] \rightarrow \widehat{\mathbb{Z}}_2[[v_1, \dots, v_{n-1}]] [v_n^\pm]$$

# Completed Johnson-Wilson

- Bousfield localization gives a completion  $E(n) \rightarrow \widehat{E(n)} = L_{K(n)}E(n)$  such that

$$E(n)^* \rightarrow \widehat{E(n)}^*$$

is  $I_n$ -adic completion for  $I_n = (v_0, \dots, v_{n-1})$ .

$$\mathbb{Z}_{(2)}[v_1, \dots, v_{n-1}, v_n^\pm] \rightarrow \widehat{\mathbb{Z}}_2[[v_1, \dots, v_{n-1}]] [v_n^\pm]$$

- Complex conjugation action on  $E(n)$  gives action on  $\widehat{E(n)}$

# Two sides of a picture

Summary so far:

- $\mathbb{Z}/2$ -action on  $E(n)$  of complex conjugation gives action on  $\widehat{E}(n)$
- $ER(n) := E(n)^{h\mathbb{Z}/2}$

# Two sides of a picture

Summary so far:

- $\mathbb{Z}/2$ -action on  $E(n)$  of complex conjugation gives action on  $\widehat{E}(n)$
- $ER(n) := E(n)^{h\mathbb{Z}/2}$

Other side is Morava  $E$ -theory and stabilizer group action...

# The Morava stabilizer group

Morava  $E$ -theory:

- Landweber exact cohomology theory  $E_n$
- $(E_n)^* = W(\mathbb{F}_{2^n})[[u_1, \dots, u_{n-1}]] [u^\pm]$

# The Morava stabilizer group

Morava  $E$ -theory:

- Landweber exact cohomology theory  $E_n$
- $(E_n)^* = W(\mathbb{F}_{2^n})[[u_1, \dots, u_{n-1}]]\langle u^\pm \rangle$
- $F_{E_n}$  is the universal deformation of the Honda formal group law  $H_n$

# The Morava stabilizer group

Morava  $E$ -theory:

- Landweber exact cohomology theory  $E_n$
- $(E_n)^* = W(\mathbb{F}_{2^n})[[u_1, \dots, u_{n-1}]][u^\pm]$
- $F_{E_n}$  is the universal deformation of the Honda formal group law  $H_n$

Morava stabilizer group:  $S_n := \text{Aut}(H_n)$

# The Morava stabilizer group

Morava  $E$ -theory:

- Landweber exact cohomology theory  $E_n$
- $(E_n)^* = W(\mathbb{F}_{2^n})[[u_1, \dots, u_{n-1}]]\langle u^\pm \rangle$
- $F_{E_n}$  is the universal deformation of the Honda formal group law  $H_n$

Morava stabilizer group:  $S_n := \text{Aut}(H_n)$

*Note:*  $S_n$  has a subgroup of order two generated by the formal inverse  $i(x)$ .

# The Morava stabilizer group

Morava  $E$ -theory:

- Landweber exact cohomology theory  $E_n$
- $(E_n)^* = W(\mathbb{F}_{2^n})[[u_1, \dots, u_{n-1}]] [u^\pm]$
- $F_{E_n}$  is the universal deformation of the Honda formal group law  $H_n$

Morava stabilizer group:  $S_n := \text{Aut}(H_n)$

*Note:*  $S_n$  has a subgroup of order two generated by the formal inverse  $i(x)$ .

Extended Morava stabilizer group:  $\mathbb{G}_n := \text{Gal}(\mathbb{F}_{2^n}/\mathbb{F}_2) \rtimes S_n$

# Hopkins-Miller-Goerss

- Lubin-Tate theory  $\Rightarrow \mathbb{G}_n$  acts on  $(E_n)_*$

# Hopkins-Miller-Goerss

- Lubin-Tate theory  $\Rightarrow \mathbb{G}_n$  acts on  $(E_n)_*$
- Hopkins-Miller-Goerss  $\Rightarrow \mathbb{G}_n$  acts on  $E_n$  by  $E_\infty$ -maps

# Hopkins-Miller-Goerss

- Lubin-Tate theory  $\Rightarrow \mathbb{G}_n$  acts on  $(E_n)_*$
- Hopkins-Miller-Goerss  $\Rightarrow \mathbb{G}_n$  acts on  $E_n$  by  $E_\infty$ -maps
- Get interesting  $E_\infty$ -ring spectra by  $E_n^{hK}$  for  $K \subseteq \mathbb{G}_n$   
e.g.  $E_n^{h\mathbb{G}_n} = L_{K(n)}\mathcal{S}$

## Hopkins-Miller-Goerss

- Lubin-Tate theory  $\Rightarrow \mathbb{G}_n$  acts on  $(E_n)_*$
- Hopkins-Miller-Goerss  $\Rightarrow \mathbb{G}_n$  acts on  $E_n$  by  $E_\infty$ -maps
- Get interesting  $E_\infty$ -ring spectra by  $E_n^{hK}$  for  $K \subseteq \mathbb{G}_n$   
e.g.  $E_n^{h\mathbb{G}_n} = L_{K(n)}\mathcal{S}$

Define  $E_n(\text{Gal}) := E_n^{hK}$  for  $K = \text{Gal}(\mathbb{F}_{2^n}/\mathbb{F}_2) \rtimes \mathbb{F}_{2^n}^\times$ .

## Hopkins-Miller-Goerss

- Lubin-Tate theory  $\Rightarrow \mathbb{G}_n$  acts on  $(E_n)_*$
- Hopkins-Miller-Goerss  $\Rightarrow \mathbb{G}_n$  acts on  $E_n$  by  $E_\infty$ -maps
- Get interesting  $E_\infty$ -ring spectra by  $E_n^{hK}$  for  $K \subseteq \mathbb{G}_n$   
e.g.  $E_n^{h\mathbb{G}_n} = L_{K(n)}\mathcal{S}$

Define  $E_n(\text{Gal}) := E_n^{hK}$  for  $K = \text{Gal}(\mathbb{F}_{2^n}/\mathbb{F}_2) \rtimes \mathbb{F}_{2^n}^\times$ .

- $E_n(\text{Gal})_* = \widehat{\mathbb{Z}}_2[[v_1, \dots, v_{n-1}]] [v_n^\pm]$
- Order 2 subgroup generated by  $i(x)$  acts on  $E_n(\text{Gal})$

# A natural question

We have  $\widehat{E}(n) \simeq E_n(\text{Gal})$  and...

$\mathbb{Z}/2$  acts on  $\widehat{E}(n)$

Complex conjugation action

$\mathbb{Z}/2$  acts on  $E_n(\text{Gal})$

Action of the subgroup of  
Morava stabilizer group  
generated by the formal inverse

# A natural question

We have  $\widehat{E}(n) \simeq E_n(\text{Gal})$  and...

$\mathbb{Z}/2$  acts on  $\widehat{E}(n)$

Complex conjugation action

$\mathbb{Z}/2$  acts on  $E_n(\text{Gal})$

Action of the subgroup of  
Morava stabilizer group  
generated by the formal inverse

**Natural question: Are these actions related?**

# A natural question

We have  $\widehat{E(n)} \simeq E_n(\text{Gal})$  and...

$\mathbb{Z}/2$  acts on  $\widehat{E(n)}$

Complex conjugation action

$\mathbb{Z}/2$  acts on  $E_n(\text{Gal})$

Action of the subgroup of  
Morava stabilizer group  
generated by the formal inverse

**Natural question: Are these actions related?**

**YES**

# Answer: Yes

## Theorem (A.)

*There is an equivalence*

$$\widehat{E(n)}^{h\mathbb{Z}/2} \simeq E_n(\text{Gal})^{h\mathbb{Z}/2}$$

*and the natural map*

$$ER(n) = E(n)^{h\mathbb{Z}/2} \rightarrow \widehat{E(n)}^{h\mathbb{Z}/2}$$

*induces an algebraic completion on coefficients.*

# Consequences

## Corollary

*After completion,  $ER(n)$  is an  $E_\infty$ -ring spectrum.*

# Consequences

## Corollary

After completion,  $ER(n)$  is an  $E_\infty$ -ring spectrum.

## Corollary

$$(E_n(\text{Gal})^{h\mathbb{Z}/2})_* = \widehat{\mathbb{Z}}_2[[\hat{v}_k(l) \mid 0 \leq k < n, l \in \mathbb{Z}]] [x, v_n^{\pm 2^{n+1}}] / J$$

$J$  is the ideal generated by the relations

$$\hat{v}_0(0) = 2 \quad x^{2^{k+1}-1} \hat{v}_k(l) = 0$$

$$\text{and for } k \leq m, \quad \hat{v}_m(l) \hat{v}_k(2^{m-k}s) = \hat{v}_m(l+s) \hat{v}_k(0)$$

$$|x| = \lambda(n) = 2^{2n+1} - 2^{n+2} + 1 \quad |v_n^{2^{n+1}}| = 2^{n+2}(2^n - 1)^2$$

$$|\hat{v}_k(l)| = 2(2^k - 1) + l^{2^{k+2}}(2^n - 1)^2 - 2(2^k - 1)(2^n - 1)^2$$

## A bit about the proof

We'd like *equivariant* map  $\varphi : \widehat{E}(n) \rightarrow E_n(\text{Gal})$  that is also an equivalence.

But we only have a *homotopy equivariant* one.

# A bit about the proof

We'd like *equivariant* map  $\varphi : \widehat{E}(n) \rightarrow E_n(\text{Gal})$  that is also an equivalence.

But we only have a *homotopy equivariant* one.

- Try replacing  $\widehat{E}(n)$  by  $\widehat{E}(n) \wedge F(\widehat{E}(n), E_n(\text{Gal}))_\varphi$
- $\varphi$  homotopy equivariant  $\Rightarrow$  conjugation action on  $F(\widehat{E}(n), E_n(\text{Gal}))_\varphi$
- $ev : \widehat{E}(n) \wedge F(\widehat{E}(n), E_n(\text{Gal}))_\varphi \rightarrow E_n(\text{Gal})$  is honestly equivariant

# A bit about the proof

We'd like *equivariant* map  $\varphi : \widehat{E}(n) \rightarrow E_n(\text{Gal})$  that is also an equivalence.

But we only have a *homotopy equivariant* one.

- Try replacing  $\widehat{E}(n)$  by  $\widehat{E}(n) \wedge F(\widehat{E}(n), E_n(\text{Gal}))_\varphi$
- $\varphi$  homotopy equivariant  $\Rightarrow$  conjugation action on  $F(\widehat{E}(n), E_n(\text{Gal}))_\varphi$
- $ev : \widehat{E}(n) \wedge F(\widehat{E}(n), E_n(\text{Gal}))_\varphi \rightarrow E_n(\text{Gal})$  is honestly equivariant
- If  $F(\widehat{E}(n), E_n(\text{Gal}))_\varphi \simeq pt$ , then  $\widehat{E}(n) \wedge F(\widehat{E}(n), E_n(\text{Gal}))_\varphi \simeq \widehat{E}(n)$ .
- Need appropriate category so that  $F(\widehat{E}(n), E_n(\text{Gal}))_\varphi \simeq pt$ . Try  $S$ -algebra maps.
- Problem: not known if  $\mathbb{Z}/2$ -action on  $E(n)$  is a  $S$ -algebra map.

# A bit about the proof

- Instead use  $F_{S\text{-alg}}(v_n^{-1}\widehat{MU}, E_n(\text{Gal}))$ . New problem: not contractible.
- Dirty trick: create  $S$ -algebra  $T$  so that  $F_{T\text{-alg}}(v_n^{-1}\widehat{MU}, E_n(\text{Gal}))$  is homotopy discrete.

# A bit about the proof

- Instead use  $F_{S\text{-alg}}(v_n^{-1}\widehat{MU}, E_n(\text{Gal}))$ . New problem: not contractible.
- Dirty trick: create  $S$ -algebra  $T$  so that  $F_{T\text{-alg}}(v_n^{-1}\widehat{MU}, E_n(\text{Gal}))$  is homotopy discrete.
  - $T =$  free commutative  $S$ -algebra on a bunch of spheres
  - $\pi_*(v_n^{-1}\widehat{MU}) = \pi_*(\widehat{E}(n) \wedge T)$
  - Compute BKSS for  $F_{T\text{-alg}}(-, E_n(\text{Gal}))$
  - A map  $\widehat{E}(n) \wedge T \rightarrow v_n^{-1}\widehat{MU}$  gives a map of spectral sequences that is an iso on  $E_2$
  - Since that for  $\widehat{E}(n) \wedge T$  collapses, so does that for  $v_n^{-1}\widehat{MU}$

# A bit about the proof

- Now

$$v_n^{-1}\widehat{MU} \wedge F_{T\text{-alg}}(v_n^{-1}\widehat{MU}, E_n(\text{Gal}))_\nu \rightarrow E_n(\text{Gal})$$

is equivariant.

- After taking homotopy fixed points, obtain a factorization

$$\begin{array}{ccc}
 v_n^{-1}\widehat{MU}^{h\widetilde{\mathbb{Z}/2}} & \rightarrow & E_n(\text{Gal})^{h\widetilde{\mathbb{Z}/2}} \\
 \downarrow & \nearrow \simeq & \\
 \widehat{E(n)}^{h\widetilde{\mathbb{Z}/2}} & & 
 \end{array}$$

Thank you!