

INEQUALITIES FOR SECOND-ORDER RIESZ TRANSFORMS ASSOCIATED WITH BESSEL EXPANSION

ADAM OSĘKOWSKI

ABSTRACT. The paper contains the proofs of L^p , logarithmic and weak-type estimates for the second-order Riesz transforms arising in the context of multidimensional Bessel expansions. Using a novel probabilistic approach, which rests on martingale methods and the representation of Riesz transforms via associated Bessel-heat processes, we show that these estimates hold with constants independent of the dimension.

1. INTRODUCTION

One of the basic examples of Calderón-Zygmund singular integral operators in \mathbb{R}^d are the so-called Riesz transforms, given by

$$R_j f(x) = \frac{\Gamma\left(\frac{d+1}{2}\right)}{\pi^{\frac{d+1}{2}}} \text{p. v.} \int_{\mathbb{R}^d} \frac{x_j - y_j}{|x - y|^{d+1}} f(y) dy, \quad j = 1, 2, \dots, d.$$

These operators and their second-order analogues (linear combinations of the expressions of the form $R_j R_k$, $j, k = 1, 2, \dots, d$) play an important role in harmonic analysis and the theory of PDEs. In particular, it is well-known that tight information about the norms of these objects, considered as operators on various function spaces, can be used in the study of regularity of certain elliptic partial differential equations. The literature on the subject is extremely vast and exploits a variety of different techniques coming from analysis and probability theory.

In the recent years, much effort has been done to extend the estimates for Riesz transforms to other contexts, in which \mathbb{R}^d , equipped with Lebesgue's measure, classical laplacian and classical Fourier transform, is replaced with some other homogeneous space possessing an appropriate differential operator. This problem has been studied in various setups, and the typical approach rests on the careful examination of the pointwise behavior of certain associated kernels. The purpose of this paper is to introduce a novel, probabilistic approach to the study of second-order Riesz transforms arising in the multidimensional setting of Bessel expansions [3, 4, 5]. To formulate our results, we need to introduce the basic setup which will be used throughout the paper. Let $d \geq 1$ be a fixed dimension and let $\lambda \in \mathbb{R}_+^d$ be a fixed multiindex. Consider the space $X = \mathbb{R}_+^d$ equipped with its Borel subsets and the product measure

$$d\mu_\lambda(x) = \prod_{j=1}^d x_j^{2\lambda_j} dx_j.$$

2010 *Mathematics Subject Classification*. Primary: 42B15, 42B20. Secondary: 42B10, 60G44.

Key words and phrases. Riesz transform, Hankel transform, Bessel expansion, martingale, differential subordination.

Then the Bessel differential operator, defined initially on $C_c^\infty(\mathbb{R}_+^d)$ by

$$\Delta^\lambda = -\Delta - \sum_{j=1}^d \frac{2\lambda_j}{x_j} \partial_{x_j},$$

has a symmetric and nonnegative extension to $L^2(\mathbb{R}_+^d, d\mu_\lambda)$. It is easy to check that Δ^λ admits the decomposition $\Delta^\lambda = \sum_{j=1}^d \delta_j^* \delta_j$, where $\delta_j = \partial_{x_j}$ and $\delta_j^* = -\partial_{x_j} - \frac{2\lambda_j}{x_j}$ is the formal adjoint of δ_j , $j = 1, 2, \dots, d$. The following information on the spectral properties of Δ^λ will be needed later. For any $z \in \mathbb{R}_+^d$, consider the function

$$\varphi_z^\lambda(x) = \prod_{j=1}^d (z_j x_j)^{-\lambda_j + 1/2} J_{\lambda_j - 1/2}(z_j x_j), \quad x \in \mathbb{R}_+^d,$$

where J_ν stands for the oscillating Bessel function of the first kind and order ν :

$$J_\nu(z) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k! \Gamma(\nu + k + 1)} (z/2)^{2k + \nu}$$

(see [14] for more on the subject). Then φ_z^λ is an eigenfunction of the Bessel operator, corresponding to the eigenvalue $|z|^2$: $\Delta^\lambda \varphi_z^\lambda = |z|^2 \varphi_z^\lambda$. Furthermore, the family $(\varphi_z^\lambda)_z$ can be used to introduce another important object, the (modified) Hankel transform \mathcal{H}^λ . This operator, defined initially on $C_c^\infty(\mathbb{R}_+^d)$, acts by the formula

$$\mathcal{H}^\lambda f(x) = \int_{\mathbb{R}_+^d} \varphi_x^\lambda(y) f(y) \mu_\lambda(dy),$$

and plays the role of the Fourier transform from the Euclidean setting. It can be extracted from the reasoning of Betancor and Stempak [5] that \mathcal{H}^λ extends to an isometry on $L^2(d\mu_\lambda)$ and satisfies $\mathcal{H}^\lambda = (\mathcal{H}^\lambda)^{-1}$. In addition, for any $f \in C_c^\infty(\mathbb{R}_+^d)$ and any $j = 1, 2, \dots, d$, we have the identity

$$(1) \quad \mathcal{H}^\lambda(\delta_j^* \delta_j f)(z) = |z_j|^2 \mathcal{H}^\lambda f(z), \quad z \in \mathbb{R}_+^d.$$

The Bessel heat semigroup $W_t^\lambda = \exp(-t\Delta^\lambda)$, corresponding to the generator $-\Delta^\lambda$, is given by

$$W_t^\lambda f = \mathcal{H}^\lambda(e^{-t|\cdot|^2} \mathcal{H}^\lambda f)$$

and admits the following representation. If $f \in L^2(d\mu_\lambda)$ and $x \in \mathbb{R}_+^d$, then

$$W_t^\lambda f(x) = \int_{\mathbb{R}_+^d} W_t^\lambda(x, y) f(y) d\mu_\lambda(y),$$

where the kernel $W_t^\lambda(\cdot, \cdot)$ is given by the formula

$$(2) \quad W_t^\lambda(x, y) = \frac{1}{(2t)^d} \exp\left(-\frac{1}{4t}(|x|^2 + |y|^2)\right) \prod_{j=1}^d (x_j y_j)^{-\lambda_j + 1/2} I_{\lambda_j - 1/2}\left(\frac{x_j y_j}{2t}\right).$$

Here $x, y \in \mathbb{R}_+^d$, $t > 0$ and

$$I_\nu(z) = \sum_{k=0}^{\infty} \frac{(z/2)^{2k + \nu}}{k! \Gamma(\nu + k + 1)}$$

stands for the non-oscillating modified Bessel function of the first kind and order ν (see [14] for details).

We are ready to formulate the main results of this paper. In what follows, for any vector $a = (a_1, a_2, \dots, a_d) \in \mathbb{C}^d$, the operator $T^{a,\lambda}$ is a Hankel multiplier with the symbol $|\xi|^{-2} \sum_{j=1}^d a_j |\xi_j|^2$; that is, we have the identity

$$\mathcal{H}^\lambda(T^{a,\lambda}f)(\xi) = |\xi|^{-2} \sum_{j=1}^d a_j |\xi_j|^2 \mathcal{H}^\lambda f(\xi), \quad \xi \in \mathbb{R}_+^d.$$

It is easy to see (using (1), for example) that $T^{a,\lambda}$ can be expressed as a linear combination of second-order Bessel-Riesz transforms:

$$T^{a,\lambda} = \sum_{j=1}^d a_j (R_j^\lambda)^* R_j^\lambda.$$

Here R_j^λ is the first-order Riesz transform given by $R_j^\lambda f(x) = \delta_j \mathcal{H}^\lambda(|\cdot|^{-1} \mathcal{H}^\lambda f)(x)$ and $(R_j^\lambda)^*$ is its formal adjoint: $(R_j^\lambda)^* f(x) = \mathcal{H}^\lambda(|\cdot|^{-1} \mathcal{H}^\lambda(\delta_j^* f))(x)$.

The primary goal of this paper is to show, using probabilistic methods, that the operators $T^{a,\lambda}$ are bounded as operators on various classical spaces. Let us start with L^p -estimates.

Theorem 1.1. *Pick a sequence $a = (a_1, a_2, \dots, a_d)$ such that $|a_j| \leq 1$, $j = 1, 2, \dots, d$. Then for any $1 < p < \infty$ we have*

$$(3) \quad \|T^{a,\lambda}\|_{L^p(\mathbb{R}_+^d, d\mu_\lambda) \rightarrow L^p(\mathbb{R}_+^d, d\mu_\lambda)} \leq p^* - 1,$$

where $p^* = \max\{p, p/(p-1)\}$.

In the boundary cases $p = 1$ and $p = \infty$, we will establish the corresponding LlogL and exponential inequalities. It will be convenient to use the functions $\Phi(t) = e^t - 1 - t$ and $\Psi(t) = (t+1) \log(t+1) - t$, defined for $t \geq 0$.

Theorem 1.2. *Pick a sequence $a = (a_1, a_2, \dots, a_d)$ such that $|a_j| \leq 1$, $j = 1, 2, \dots, d$. Then for any $K > 1$ and any Borel subset A of \mathbb{R}_+^d we have*

$$(4) \quad \int_A |T^{a,\lambda} f(x)| d\mu_\lambda(x) \leq K \int_{\mathbb{R}_+^d} \Psi(|f(x)|) d\mu_\lambda(x) + \frac{\mu_\lambda(A)}{2(K-1)}.$$

Furthermore, if $\|f\|_{L^\infty(\mathbb{R}_+^d, d\mu_\lambda)} \leq 1$, then

$$(5) \quad \int_{\mathbb{R}_+^d} \Phi(|T^{a,\lambda} f(x)|) d\mu_\lambda(x) \leq \frac{1}{2K(K-1)} \|f\|_{L^1(\mathbb{R}_+^d, d\mu_\lambda)}.$$

We will also establish the corresponding weak-type bounds; for any $1 < p < \infty$, we will work with the norming

$$\|f\|_{L^{p,\infty}(\mathbb{R}_+^d, d\mu_\lambda)} = \sup \left\{ \frac{1}{\mu_\lambda(A)^{1-1/p}} \int_A |f(x)| d\mu_\lambda(x) \right\},$$

where the supremum is taken over all Borel subsets A of \mathbb{R}_+^d , satisfying $\mu_\lambda(A) > 0$. Introduce the constants

$$K_p = \begin{cases} \left[\frac{1}{2} \Gamma \left(\frac{2p-1}{p-1} \right) \right]^{(p-1)/p} & \text{if } 1 < p < 2, \\ (p^{p-1}/2)^{1/p} & \text{if } p \geq 2. \end{cases}$$

Theorem 1.3. *Assume that $1 < p < \infty$ and let a_1, a_2, \dots, a_d be elements of unit ball in \mathbb{C} . Then*

$$(6) \quad \left\| T^{a,\lambda} \right\|_{L^p(\mathbb{R}_+^d, d\mu_\lambda) \rightarrow L^{p,\infty}(\mathbb{R}_+^d, d\mu_\lambda)} \leq K_p.$$

A few comments on the method of proof are in order. A classical argument used to establish results of the above type studies various delicate properties of the kernel W^λ . Our approach will be completely different and will exploit probabilistic methods: the above estimates will be deduced from some deep results from martingale theory. As a by-product, we obtain constants which do not depend on the dimension d or the value of parameter λ .

The paper is organized as follows. In the next section we present the probabilistic facts which will be needed to establish the aforementioned inequalities. Section 3 links the probabilistic and analytic aspects of the paper, and is devoted to a martingale representation of the operators $T^{a,\lambda}$. In the final part we put all the facts together and establish Theorems 1.1, 1.2 and 1.3.

2. PROBABILISTIC BACKGROUND

As we have announced in the preceding section, our argumentation depends heavily on probabilistic techniques. Let us introduce the necessary setup and notation. Suppose that $(\Omega, \mathcal{F}, \mathbb{P})$ is a complete probability space, equipped with $(\mathcal{F}_t)_{t \geq 0}$, a nondecreasing family of sub- σ -fields of \mathcal{F} , such that \mathcal{F}_0 contains all the events of probability 0. Assume further that X, Y are two adapted martingales taking values in a certain separable Hilbert space $(\mathcal{H}, |\cdot|)$; with no loss of generality, we may put $\mathcal{H} = \ell^2$. As usual, we impose standard conditions on the trajectories of these processes: we assume that the paths are right-continuous and have limits from the left. Given $p \in [1, \infty]$, we will write $\|X\|_p$ for the p -th moment of X , given by $\|X\|_p = \sup_{t \geq 0} \|X_t\|_p$. The symbol $[X, Y]$ will stand for the quadratic covariance process of X and Y . See e.g. Dellacherie and Meyer [8] for details in the case when the processes are real-valued, and extend the definition to the vector setting by $[X, Y] = \sum_{k=0}^{\infty} [X^k, Y^k]$, where X^k, Y^k are the k -th coordinates of X, Y , respectively. We will say that Y is differentially subordinate to X , if the process $([X, X]_t - [Y, Y]_t)_{t \geq 0}$ is nonnegative and nondecreasing as a function of t . This notion appeared for the first time in the discrete-time setting in the works of Burkholder [6, 7]; the above continuous-time extension was introduced by Bañuelos and Wang [2] and Wang [13]. The differential subordination implies many interesting inequalities between the martingales X and Y ; for an overview of the results, methods and many more, see the monograph by Osękowski [9].

The inequalities (3)-(6) will be deduced from their probabilistic counterparts. We start with the following L^p -estimate, proved by Burkholder [6] in the discrete-time setting and extended to the general context by Wang [13].

Theorem 2.1. *Suppose that Y is differentially subordinate to X . Then for any $1 < p < \infty$ we have*

$$\|Y\|_p \leq (p^* - 1)\|X\|_p.$$

To show (5), we will need the following statement, established in [10]. Then (4) will be obtained with the use of duality-type argument.

Theorem 2.2. *Assume that X, Y are \mathcal{H} -valued martingales such that $\|X\|_\infty \leq 1$ and Y is differentially subordinate to X . Then for any $K > 1$ we have*

$$(7) \quad \sup_{t \geq 0} \mathbb{E} \Phi(|Y_t|/K) \leq \frac{1}{2K(K-1)} \|X\|_1.$$

Finally, to prove (6), we will exploit the following fact from [11], which can be regarded as a dual to the weak-type inequalities between X and Y .

Theorem 2.3. *Assume that X, Y are \mathcal{H} -valued martingales such that Y is differentially subordinate to X . Then for any $1 < q < \infty$ and $p = q/(q-1)$,*

$$(8) \quad \|Y\|_q^q \leq K_p^q \|X\|_1 \|X\|_\infty^{q-1}.$$

In the remainder of this section, we will provide some basic facts concerning Bessel processes. The interested reader is referred to [12, Chapter XI] for the systematic presentation of the subject.

Let $\beta = (\beta_t)_{t \geq 0}$ be a standard, one-dimensional Brownian motion. For every $\delta \geq 0$ and $x \geq 0$, the equation

$$Z_t = x + 2 \int_0^t \sqrt{Z_s} d\beta_s + \delta t$$

has a unique strong solution, which is called the square of δ -dimensional Bessel process started at x (and denoted by $BESQ^\delta(x)$). For any $x \geq 0$, the square root of $BESQ^\delta(x^2)$ is called the Bessel process of dimension δ started at x , and is denoted by $BES^\delta(x)$. The family $(BES^\delta(x))_{x \geq 0}$ is a Markov family with the density

$$(9) \quad p_t^\delta(x, y) = t^{-1} (y/x)^{\delta/2-1} y \exp\left(-\frac{x^2+y^2}{2t}\right) I_{\delta/2-1}\left(\frac{xy}{t}\right), \quad x, y > 0.$$

Obviously, the function $W_t^\lambda(x, y)$ given by (2) is closely related to the product $\prod_{j=1}^d p_{2t}^{2\lambda_j+1}(x_j, y_j)$; both functions are essentially equal (the slight difference in the formulas comes from the fact that W_t^λ is the density of the semigroup with respect to the measure $d\mu_\lambda$, while (9) refers to Lebesgue measure).

3. PROBABILISTIC REPRESENTATION OF SECOND-ORDER RIESZ TRANSFORMS

Now we will explain how second-order Riesz transforms can be expressed in terms of stochastic integrals involving Bessel processes. Suppose that f is a smooth, compactly supported and complex-valued function on \mathbb{R}_+^d and let $U_f : \mathbb{R}_+^d \times [0, \infty) \rightarrow \mathbb{C}$ denote the Bessel-heat extension of f : $U_f(\cdot, 0) = f$ and, for any $x \in \mathbb{R}_+^d$ and $t > 0$,

$$U_f(x, t) = W_t^\lambda f(x) = \int_{\mathbb{R}_+^d} W_t^\lambda(x, y) f(y) d\mu_\lambda(y),$$

where the kernel $W_t^\lambda(\cdot, \cdot)$ is given by (2). This extension is a C^∞ function on $\mathbb{R}_+^d \times \mathbb{R}^+$ and satisfies the PDE

$$(10) \quad \Delta_x^\lambda U_f + \frac{\partial U_f}{\partial t} = 0,$$

where Δ_x^λ is the Bessel differential operator applied to the x -coordinate. We will also need the following ‘‘square’’ regularity property of U_f . Note that W_t^λ extends

to a C^∞ function on $\mathbb{R}^d \times \mathbb{R}^d$, with the use of the formula

$$W_t^\lambda(x, y) = \frac{1}{(2t)^d} \exp\left(-\frac{1}{4t}(|x|^2 + |y|^2)\right) \prod_{j=1}^d \sum_{k=0}^{\infty} \frac{(x_j y_j)^{2k}}{(4t)^{2k+\lambda_j-1/2} k! \Gamma(k + \lambda + 1/2)}$$

and hence $W_t^\lambda(\cdot, y)$ is a smooth function of $x_1^2, x_2^2, \dots, x_d^2$. Clearly, this property is inherited by the function $U_f(\cdot, t)$.

Now, for a fixed $x \in \mathbb{R}_+^d$, let $X^x = (X^{1,x_1}, X^{2,x_2}, \dots, X^{d,x_d})$ be the collection of independent processes such that for each $j = 1, 2, \dots, d$, the coordinate X^{j,x_j} is a Bessel process of dimension $2\lambda_j + 1$, started at x_j . For a given a positive number T , we introduce the associated parabolic process $F = F(x; T; f)$ by

$$F_t = U_f(X_{2t}^x, T - t), \quad t \in [0, T].$$

The next step is to apply Itô's formula to F . We have the SPDEs $d(X^{j,x_j})_t^2 = 2X_t^{j,x_j} dB_t^j + (2\lambda_j + 1)dt$, where $B = (B^1, B^2, \dots, B^d)$ is a Brownian motion in \mathbb{R}^d . Let us rewrite F in the form $F_t = U_f(\sqrt{(X_{2t}^x)^2}, T - t)$, $t \in [0, T]$. Then Itô's formula, combined with (10), yields

$$F_t(x; T; f) = U_f(x, T) + \int_{0+}^t \nabla_x U_f(X_{2s}^x, T - s) \cdot dB_{2s}, \quad t \in [0, T].$$

This in particular implies that the process F is a continuous-path martingale. Note that F is bounded, and hence square-integrable. It follows from classical facts from stochastic analysis that the quadratic variation of F equals

$$[F, F]_t = |U_f(x, T)|^2 + 2 \int_{0+}^t |\nabla_x U_f(X_{2s}^x, T - s)|^2 ds, \quad t \in [0, T].$$

The next step of the construction is to apply a certain transformation to F . Suppose that \mathcal{A} is a diagonal matrix of dimension $d \times d$, such that each entry belongs to the unit ball of \mathbb{C} . Consider the associated transform of F , given by

$$G_t = G_t(x; T; f; \mathcal{A}) = \int_{0+}^t \mathcal{A} \nabla_x U_f(X_{2s}^x, T - s) \cdot dB_{2s}, \quad t \in [0, T].$$

Then G is a square integrable martingale, with the quadratic covariance process equal to

$$[G, G]_t = 2 \int_{0+}^t |\mathcal{A} \nabla_x U_f(X_{2s}^x, T - s)|^2 ds, \quad t \in [0, T].$$

Since the operator norm of \mathcal{A} does not exceed one, we see that G is differentially subordinate to F : the process

$$\begin{aligned} & [F, F]_t - [G, G]_t \\ &= |U_f(x, T)|^2 + 2 \int_{0+}^t |\nabla_x U_f(X_{2s}^x, T - s)|^2 - |\mathcal{A} \nabla_x U_f(X_{2s}^x, T - s)|^2 ds \end{aligned}$$

is nonnegative and nondecreasing. As we will show now, the appropriate projection of the process G leads to an associated Hankel multiplier. To this end, observe that for any $h \in L^2(\mathbb{R}_+^d, d\mu_\lambda)$, we have, by Schwarz inequality and Fubini's theorem

(and the fact that μ_λ is the invariant measure for the Bessel process X),

$$\begin{aligned}
& \mathbb{E} \int_{\mathbb{R}_+^d} |G_T(x; T; f; \mathcal{A})h(X_{2T}^x)| d\mu_\lambda(x) \\
& \leq \left(\int_{\mathbb{R}_+^d} \mathbb{E}|G_T(x; T; f; \mathcal{A})|^2 d\mu_\lambda(x) \right)^{1/2} \left(\int_{\mathbb{R}_+^d} \mathbb{E}|h(X_{2T}^x)|^2 d\mu_\lambda(x) \right)^{1/2} \\
& = \left(\int_{\mathbb{R}_+^d} \mathbb{E}[G(x; T; f; \mathcal{A}), G(x; T; f; \mathcal{A})]_T d\mu_\lambda(x) \right)^{1/2} \left(\mathbb{E} \int_{\mathbb{R}_+^d} |h(X_{2T}^x)|^2 d\mu_\lambda(x) \right)^{1/2} \\
& \leq \left(\int_{\mathbb{R}_+^d} \mathbb{E}[F(x; T; f), F(x; T; f)]_T d\mu_\lambda(x) \right)^{1/2} \|h\|_{L^2(\mathbb{R}_+^d, d\mu_\lambda)} \\
& = \left(\int_{\mathbb{R}_+^d} \mathbb{E}|F_T(x; T; f)|^2 d\mu_\lambda(x) \right)^{1/2} \|h\|_{L^2(\mathbb{R}_+^d, d\mu_\lambda)} \\
& = \left(\int_{\mathbb{R}_+^d} \mathbb{E}|f(X_{2T}^x)|^2 d\mu_\lambda(x) \right)^{1/2} \|h\|_{L^2(\mathbb{R}_+^d, d\mu_\lambda)} \\
& = \|f\|_{L^2(\mathbb{R}_+^d, d\mu_\lambda)} \|h\|_{L^2(\mathbb{R}_+^d, d\mu_\lambda)}.
\end{aligned}$$

Consequently, there is a unique function $g = S^{T, \mathcal{A}}f \in L^2(\mathbb{R}_+^d, d\mu_\lambda)$ defined through the bilinear form

$$\int_{\mathbb{R}_+^d} g(x)h(x)d\mu_\lambda(x) = \mathbb{E} \int_{\mathbb{R}_+^d} G_T(x; T; f; \mathcal{A})h(X_{2T}^x)d\mu_\lambda(x), \quad h \in L^2(\mathbb{R}_+^d, d\mu_\lambda).$$

(Informally, we can treat $g(x)$ as the projection, or rather the ‘‘conditional expectation’’ $\mathbb{E}_{\mathcal{P}}[G_T(x; T; f; \mathcal{A})|X_{2T}^x = x]$ with respect to the product, non-probability measure $\mathcal{P} = \mathbb{P} \otimes \mu_\lambda$). We will prove now that $S^{T, \mathcal{A}}$ is a Hankel multiplier and identify the associated symbol. Using some basic properties of stochastic integrals, we note that for any $x \in \mathbb{R}_+^d$,

$$\begin{aligned}
& \mathbb{E}G_T(x; T; f; \mathcal{A})h(X_{2T}^x) \\
& = \mathbb{E} \int_{0+}^T \mathcal{A}\nabla_x U_f(X_{2s}^x, T-s)dB_{2s} \int_{0+}^T \nabla_x U_h(X_{2s}^x, T-s)dB_{2s} \\
& = 2\mathbb{E} \int_{0+}^T \langle \mathcal{A}\nabla_x U_f(X_{2s}^x, T-s), \nabla_x U_h(X_{2s}^x, T-s) \rangle ds \\
& = 2 \int_{0+}^T \int_{\mathbb{R}_+^d} \langle \mathcal{A}\nabla_x U_f(y, T-s), \nabla_x U_h(y, T-s) \rangle p_{2s}(x, y) dy ds \\
& = 2 \int_{0+}^T \int_{\mathbb{R}_+^d} \langle \mathcal{A}\nabla_x U_f(y, T-s), \nabla_x U_h(y, T-s) \rangle W_s^\lambda(x, y) d\mu_\lambda(y) ds,
\end{aligned}$$

where $p_s(x, y) = \prod_{j=1}^d p_s^{2\lambda_j+1}(x_j, y_j)$ is the transition density of the process X^x . Therefore, by Fubini’s theorem and the fact that $\int_{\mathbb{R}_+^d} W_s^\lambda(x, y) d\mu_\lambda(x) = 1$ for each

y , we may write

$$\begin{aligned}
& \int_{\mathbb{R}_+^d} S^{T,\mathcal{A}} f(x) h(x) d\mu_\lambda(x) \\
&= 2 \int_{\mathbb{R}_+^d} \int_{0+}^T \int_{\mathbb{R}_+^d} \langle \mathcal{A} \nabla_x U_f(y, T-s), \nabla_x U_h(y, T-s) \rangle W_s^\lambda(x, y) d\mu_\lambda(x) ds d\mu_\lambda(y) \\
&= 2 \int_{\mathbb{R}_+^d} \int_{0+}^T \langle \mathcal{A} \nabla_x U_f(y, T-s), \nabla_x U_h(y, T-s) \rangle ds d\mu_\lambda(dy) \\
&= 2 \int_{\mathbb{R}_+^d} \int_{0+}^T \sum_{j=1}^d a_{jj} U_f(y, T-s) \delta_j^* \delta_j U_h(y, T-s) ds d\mu_\lambda(dy).
\end{aligned}$$

Here in the last line we have used integration by parts. Now fix a vector $\xi \in \mathbb{R}_+^d$ and, as a function h , put

$$h(x) = \varphi_\xi^\lambda(x) = \prod_{j=1}^d (\xi_j x_j)^{-\lambda_j + 1/2} J_{\lambda_j - 1/2}(\xi_j x_j),$$

the eigenfunction of Δ^λ corresponding to the eigenvalue $|\xi|^2$. Then

$$U_h(x, t) = \varphi_\xi^\lambda(x) \exp(-t|\xi|^2)$$

and hence $\delta_j^* \delta_j U_h(x, t) = |\xi_j|^2 \varphi_\xi^\lambda(x) \exp(-t|\xi|^2)$. Plugging this above, we get

$$\begin{aligned}
& \mathcal{H}^\lambda(S^{T,\mathcal{A}} f)(\xi) \\
&= \int_{\mathbb{R}_+^d} S^{T,\mathcal{A}} f(x) h(x) dx \\
&= 2 \sum_{j=1}^d a_{jj} |\xi_j|^2 \int_{\mathbb{R}_+^d} \int_{0+}^T U_f(y, T-s) \varphi_\xi^\lambda(y) \exp(-(T-s)|\xi|^2) ds d\mu_\lambda(dy) \\
&= 2 \sum_{j=1}^d a_{jj} |\xi_j|^2 \int_{0+}^T \exp(-(T-s)|\xi|^2) \mathcal{H}^\lambda(U_f(\cdot, T-s)) ds
\end{aligned}$$

However, we have $\mathcal{H}^\lambda(U_f(\cdot, t))(\xi) = e^{-t|\xi|^2} \mathcal{H}^\lambda f(\xi)$, which implies

$$\begin{aligned}
\mathcal{H}^\lambda(S^{T,\mathcal{A}} f)(\xi) &= 2 \sum_{j=1}^d a_{jj} |\xi_j|^2 \mathcal{H}^\lambda f(\xi) \cdot \int_{0+}^T e^{-2(T-s)|\xi|^2} ds \\
&= \mathcal{H}^\lambda(f)(\xi) \frac{\langle \mathcal{A}\xi, \xi \rangle}{|\xi|^2} [1 - e^{-2T|\xi|^2}].
\end{aligned}$$

Thus, $S^{T,\mathcal{A}}$ is a Hankel multiplier with the symbol $\langle \mathcal{A}\xi, \xi \rangle |\xi|^{-2} [1 - e^{-2T|\xi|^2}]$.

4. PROOFS OF THEOREMS 1.1, 1.2 AND 1.3

Equipped with the representation of the preceding section, we are ready for the proofs of the results announced in the introduction.

Proof of (3). By a straightforward approximation argument, it is enough to show that

$$\int_{\mathbb{R}_+^d} |T^{a,\lambda} f|^p d\mu_\lambda \leq (p^* - 1)^p \int_{\mathbb{R}_+^d} |f|^p d\mu_\lambda$$

for any $f \in C_c^\infty(\mathbb{R}_+^d)$. Let A be the diagonal matrix with the terms a_1, a_2, \dots, a_d on the main diagonal, and let F, G be the associated martingales introduced in the preceding section. We will combine the differential subordination of these processes with Theorem 2.1. To this end, recall that $q = p/(p - 1)$ is the conjugate to p and note that for any $h \in L^2(\mathbb{R}^d, d\mu_\lambda)$,

$$\begin{aligned} & \mathbb{E} \int_{\mathbb{R}_+^d} |G_T(x; T; f; \mathcal{A}) h(X_{2T}^x)| d\mu_\lambda(x) \\ & \leq \left(\int_{\mathbb{R}_+^d} \mathbb{E} |G_T(x; T; f; \mathcal{A})|^p d\mu_\lambda(x) \right)^{1/p} \left(\int_{\mathbb{R}_+^d} \mathbb{E} |h(X_{2T}^x)|^q d\mu_\lambda(x) \right)^{1/q} \\ & \leq (p^* - 1) \left(\int_{\mathbb{R}_+^d} \mathbb{E} |F_T(x; T; f)|^p d\mu_\lambda(x) \right)^{1/p} \left(\int_{\mathbb{R}_+^d} \mathbb{E} |h(X_{2T}^x)|^q d\mu_\lambda(x) \right)^{1/q} \\ & = (p^* - 1) \|f\|_{L^p(\mathbb{R}_+^d, d\mu_\lambda)} \|h\|_{L^q(\mathbb{R}_+^d, d\mu_\lambda)}, \end{aligned}$$

by Fubini's theorem. This, by the very definition of $S^{T,\mathcal{A}}$, implies

$$(11) \quad \|S^{T,\mathcal{A}} f\|_{L^p(\mathbb{R}_+^d, d\mu_\lambda)} \leq (p^* - 1) \|f\|_{L^p(\mathbb{R}_+^d, d\mu_\lambda)}.$$

Now, let $T \rightarrow \infty$. Since \mathcal{H}^λ is an isometry, we see that $S^{T,\mathcal{A}} f$ converges in $L^2(\mathbb{R}_+^d, d\mu_\lambda)$ to the function $S^{\mathcal{A}} f$, where $S^{\mathcal{A}}$ is a Hankel multiplier with the symbol $\langle \mathcal{A}\xi, \xi \rangle / |\xi|^2$. Hence we can pick a sequence T_n converging to infinity such that $S^{T_n, \mathcal{A}} f$ converges to $S^{\mathcal{A}} f$ μ_λ -almost everywhere on \mathbb{R}_+^d . So, Fatou's lemma combined with (11) gives

$$\|S^{\mathcal{A}} f\|_{L^p(\mathbb{R}_+^d, d\mu_\lambda)} \leq (p^* - 1) \|f\|_{L^p(\mathbb{R}_+^d, d\mu_\lambda)}$$

and it remains to note that $S^{\mathcal{A}}$ coincides with $T^{a,\lambda}$. \square

Proof of (5). We may and do assume that $\|f\|_{L^1(\mathbb{R}_+^d, d\mu_\lambda)} < \infty$, since otherwise there is nothing to prove. We also know that f is bounded by 1, and hence, by Hölder inequality, $f \in L^2(\mathbb{R}_+^d, d\mu_\lambda)$. Recall the functions $\Phi(t) = e^t - 1 - t$ and $\Psi(t) = (t + 1) \log(t + 1) - t$; one easily checks that Ψ' and Φ' are inverse to each other, so

$$(12) \quad ab \leq \Psi(a) + \Phi(b)$$

for any nonnegative a and b . Now, f is bounded by 1, so the martingale $F(x; T; f)$ also enjoys this property. So, by (12),

$$\begin{aligned}
& \left| \int_{\mathbb{R}_+^d} S^{T, \mathcal{A}} f(x) h(x) d\mu_\lambda(x) \right| \\
& \leq \mathbb{E} \int_{\mathbb{R}_+^d} |G_T(x; T; f; \mathcal{A}) h(X_{2T}^x)| d\mu_\lambda(x) \\
& \leq K \int_{\mathbb{R}_+^d} \Psi(|h(X_{2T}^x)|) d\mu_\lambda(x) + K \int_{\mathbb{R}_+^d} \mathbb{E} \Phi(|G_T(x; T; f; \mathcal{A})|/K) d\mu_\lambda(x) \\
& \leq K \int_{\mathbb{R}_+^d} \Psi(|h(X_{2T}^x)|) d\mu_\lambda(x) + \frac{1}{2(K-1)} \int_{\mathbb{R}_+^d} \mathbb{E} |F_T(x; T; f)| d\mu_\lambda(x) \\
& = K \int_{\mathbb{R}_+^d} \Psi(|h(X_{2T}^x)|) d\mu_\lambda(x) + \frac{1}{2(K-1)} \|f\|_{L^1(\mathbb{R}_+^d, d\mu_\lambda)}.
\end{aligned}$$

Here in the second passage we have exploited the inequality (7), and the last identity follows from Fubini's theorem and the fact that μ_λ is the invariant measure for X . Now fix a positive number M and consider the function

$$h(x) = \frac{S^{T, \mathcal{A}} f(x)}{|S^{T, \mathcal{A}} f(x)|} (\exp(|S^{T, \mathcal{A}} f(x)|/K) - 1) \chi_{\{|S^{T, \mathcal{A}} f(x)| \leq M\}},$$

with the convention $h(x) = 0$ if $S^{T, \mathcal{A}} f(x) = 0$. This function belongs to $L^2(\mathbb{R}_+^d, d\mu_\lambda)$, since $|h(x)| \leq L |S^{T, \mathcal{A}} f(x)|$ for some constant L (here we use the presence of the characteristic function in the definition of h) and $S^{T, \mathcal{A}} f \in L^2(\mathbb{R}_+^d, d\mu_\lambda)$ (which follows from (11) and the inclusion $f \in L^2(\mathbb{R}_+^d, d\mu_\lambda)$). Plugging h into the above chain of inequalities, we get

$$\begin{aligned}
& \int_{\mathbb{R}_+^d} |S^{T, \mathcal{A}} f(x)| \left(\exp\left(\frac{|S^{T, \mathcal{A}} f(x)|}{K}\right) - 1 \right) \chi_{\{|S^{T, \mathcal{A}} f(x)| \leq M\}} d\mu_\lambda(x) \\
& \leq K \int_{\mathbb{R}_+^d} \left[\exp\left(\frac{|S^{T, \mathcal{A}} f(x)|}{K}\right) \left(\frac{|S^{T, \mathcal{A}} f(x)|}{K} - 1 \right) + 1 \right] \chi_{\{|S^{T, \mathcal{A}} f(x)| \leq M\}} d\mu_\lambda(x) \\
& \quad + \frac{1}{2(K-1)} \|f\|_{L^1(\mathbb{R}_+^d, d\mu_\lambda)}.
\end{aligned}$$

This is equivalent to

$$\int_{\mathbb{R}_+^d} \Phi\left(\frac{|S^{T, \mathcal{A}} f(x)|}{K}\right) \chi_{\{|S^{T, \mathcal{A}} f(x)| \leq M\}} d\mu_\lambda(x) \leq \frac{1}{2K(K-1)} \|f\|_{L^1(\mathbb{R}_+^d, d\mu_\lambda)},$$

so letting $M \rightarrow \infty$ and using Fatou's lemma gives

$$\int_{\mathbb{R}_+^d} \Phi\left(\frac{|S^{T, \mathcal{A}} f(x)|}{K}\right) d\mu_\lambda(x) \leq \frac{1}{2K(K-1)} \|f\|_{L^1(\mathbb{R}_+^d, d\mu_\lambda)}.$$

It remains to let $T \rightarrow \infty$ and repeat the argument used above in the proof of the L^p estimate. \square

Proof of (4). We will use duality. We have

$$\int_A |T^{a, \lambda} f(x)| d\mu_\lambda(x) = \int_{\mathbb{R}_+^d} T^{a, \lambda} f(x) \overline{\eta(x)} d\mu_\lambda(x),$$

where $\eta(x) = \chi_A(x)T^{a,\lambda}f(x)/|T^{a,\lambda}f(x)|$ (again with the convention $\eta(x) = 0$ when $T^{a,\lambda}f(x) = 0$). Applying Hankel transform, we see that the latter integral equals

$$\begin{aligned} \int_{\mathbb{R}_+^d} \mathcal{H}^\lambda(T^{a,\lambda}f)(\xi)\overline{\mathcal{H}^\lambda(\eta)(\xi)}d\mu_\lambda(\xi) &= \int_{\mathbb{R}_+^d} \frac{\langle \mathcal{A}\xi, \xi \rangle}{|\xi|^2} \mathcal{H}^\lambda f(\xi)\overline{\mathcal{H}^\lambda(\eta)(\xi)}d\mu_\lambda(\xi) \\ &= \int_{\mathbb{R}_+^d} \mathcal{H}^\lambda f(\xi)\overline{\mathcal{H}^\lambda(T^{\bar{a},\lambda}\eta)(\xi)}d\mu_\lambda(\xi) \\ &= \int_{\mathbb{R}_+^d} f(x)\overline{T^{\bar{a},\lambda}\eta(x)}d\mu_\lambda(x), \end{aligned}$$

where A is the diagonal matrix with the numbers a_1, a_2, \dots, a_n on the main diagonal, and \bar{a} stands for the conjugate sequence $\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n$. Now we exploit the inequalities (5) and (12), together with the fact that η is bounded by 1. As the result, we get

$$\begin{aligned} &\int_A |T^{a,\lambda}f(x)|d\mu_\lambda(x) \\ &\leq K \int_{\mathbb{R}_+^d} \Psi(|f(x)|)d\mu_\lambda(x) + K \int_{\mathbb{R}_+^d} \Phi\left(\frac{|T^{\bar{a},\lambda}\eta(x)|}{K}\right)d\mu_\lambda(x) \\ &\leq K \int_{\mathbb{R}_+^d} \Psi(|f(x)|)d\mu_\lambda(x) + \frac{1}{2(K-1)}\|\eta\|_{L^1(\mathbb{R}_+^d, d\mu_\lambda)} \\ &= K \int_{\mathbb{R}_+^d} \Psi(|f(x)|)d\mu_\lambda(x) + \frac{\mu_\lambda(A)}{2(K-1)}, \end{aligned}$$

which is the desired assertion. \square

Proof of (6). Arguing as in the proof of the L^p estimate (3), we deduce the inequality

$$(13) \quad \|T^{a,\lambda}f\|_{L^q(\mathbb{R}_+^d, d\mu_\lambda)}^q \leq K_p^q \|f\|_{L^1(\mathbb{R}_+^d, d\mu_\lambda)} \|f\|_{L^\infty(\mathbb{R}_+^d, d\mu_\lambda)}^{q-1}$$

from (8). Now, fix an arbitrary function $f \in L^p(\mathbb{R}_+^d, d\mu_\lambda)$ and a Borel set $A \subset \mathbb{R}_+^d$. As in the proof of (4), we write

$$\int_A |T^{a,\lambda}f(x)|d\mu_\lambda(x) = \int_{\mathbb{R}_+^d} T^{a,\lambda}f(x)\overline{\eta(x)}d\mu_\lambda(x) = \int_{\mathbb{R}_+^d} f(x)\overline{T^{\bar{a},\lambda}\eta(x)}d\mu_\lambda(x),$$

where $\eta(x) = \chi_A(x)T^{a,\lambda}f(x)/|T^{a,\lambda}f(x)|$. This implies, by Hölder's inequality and (13),

$$\begin{aligned} \int_A |T^{a,\lambda}f(x)|d\mu_\lambda(x) &\leq \|f\|_{L^p(\mathbb{R}_+^d, d\mu_\lambda)} \|T^{\bar{a},\lambda}\eta\|_{L^q(\mathbb{R}_+^d, d\mu_\lambda)} \\ &\leq K_p \|f\|_{L^p(\mathbb{R}_+^d, d\mu_\lambda)} \|\eta\|_{L^1(\mathbb{R}_+^d, d\mu_\lambda)}^{1/q} \|\eta\|_{L^\infty(\mathbb{R}_+^d, d\mu_\lambda)}^{1-1/q} \\ &\leq K_p \|f\|_{L^p(\mathbb{R}_+^d, d\mu_\lambda)} \mu_\lambda(A)^{1/q}. \end{aligned}$$

This yields (6), since A was arbitrary. \square

ACKNOWLEDGMENT

The author would like to express his gratitude to the anonymous Referee for the careful reading of the paper and several helpful suggestions. The research was supported by Narodowe Centrum Nauki (Poland), grant DEC-2014/14/E/ST1/00532.

REFERENCES

- [1] M. Abramowitz and I. A. Stegun, editors, *Handbook of Mathematical Functions with formulas, graphs and mathematical tables*, Reprint of the 1972 edition, Dover Publications, Inc., New York, 1992.
- [2] R. Bañuelos and G. Wang, *Sharp inequalities for martingales with applications to the Beurling-Ahlfors and Riesz transformations*, Duke Math. J. **80** (1995), 575–600.
- [3] J. J. Betancor, J. C. Fariña, D. Buraczewski, T. Martínez and J. L. Torrea, *Riesz transforms related to Bessel operators*, Proc. Roy. Soc. Edinburgh Sect. A **137** (2007), no. 4, 701–725.
- [4] J. J. Betancor, J. C. Fariña, T. Martínez and L. Rodríguez-Mesa, *Higher order Riesz transforms associated with Bessel operators*, Ark. Mat. **46** (2008), no. 2, 219–250.
- [5] J. J. Betancor and K. Stempak, *Relating multipliers and transplantation for Fourier-Bessel expansions and Hankel transform*, Tohoku Math. J. (2) **53** (2001), no. 1, 109–129.
- [6] D. L. Burkholder, *Boundary value problems and sharp inequalities for martingale transforms*, Ann. Probab. **12** (1984), 647–702.
- [7] D. L. Burkholder, *A sharp and strict L^p -inequality for stochastic integrals*, Ann. Probab. **15** (1987), pp. 268–273.
- [8] C. Dellacherie and P.-A. Meyer, *Probabilities and potential B: Theory of martingales*, North Holland, Amsterdam, 1982.
- [9] A. Osękowski, *Sharp martingale and semimartingale inequalities*, Monografie Matematyczne **72** (2012), Birkhäuser, 462 pp.
- [10] A. Osękowski, *Logarithmic inequalities for Fourier multipliers*, Math. Z. **274** (2013), pp. 515–530.
- [11] A. Osękowski, *Weak type inequalities for Fourier multipliers with applications to Beurling-Ahlfors transform*, J. Math. Soc. Japan **66** (2014), pp. 745–764.
- [12] D. Revuz and M. Yor, *Continuous martingales and Brownian motion*, third edition, Springer, 1999.
- [13] G. Wang, *Differential subordination and strong differential subordination for continuous time martingales and related sharp inequalities*, Ann. Probab. **23** (1995), 522–551.
- [14] G. N. Watson, *A treatise on the theory of Bessel functions*, Cambridge Univ. Press, Cambridge, 1966.

FACULTY OF MATHEMATICS, INFORMATICS AND MECHANICS, UNIVERSITY OF WARSAW, BANACHA 2, 02-097 WARSAW, POLAND
Email address: ados@mimuw.edu.pl