

SHARP WEAK TYPE INEQUALITIES FOR DIFFERENTIALLY SUBORDINATED MARTINGALES

ADAM OŚĘKOWSKI

ABSTRACT. Let M, N be real valued martingales such that N is differentially subordinate to M . The paper contains the proofs of the following weak type inequalities.

(i) If $M \geq 0$ and $0 < p \leq 1$, then

$$\|N\|_{p,\infty} \leq 2\|M\|_p$$

and the constant is the best possible.

(ii) If $M \geq 0$ and $p \geq 2$, then

$$\|N\|_{p,\infty} \leq \frac{p}{2}(p-1)^{-1/p}\|M\|_p$$

and the constant is the best possible.

(iii) If $1 \leq p \leq 2$ and M and N are orthogonal, then

$$\|N\|_{p,\infty} \leq K_p\|M\|_p,$$

where

$$K_p = \frac{1}{\Gamma(p+1)} \cdot \frac{1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots}{1 - \frac{1}{3^{p+1}} + \frac{1}{5^{p+1}} - \frac{1}{7^{p+1}} + \dots}.$$

The constant is the best possible.

We also provide related estimates for harmonic functions on Euclidean domains.

1. INTRODUCTION

The purpose of this paper is to study some sharp estimates for continuous-time martingales. However, to introduce the main concepts and to present the motivations, we will start from the discrete time setting. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, filtered by nondecreasing family (\mathcal{F}_n) of sub- σ -algebras of \mathcal{F} . Let $f = (f_n)$, $g = (g_n)$ be two sequences adapted to (\mathcal{F}_n) , taking values in a separable Hilbert space. Let $df = (df_n)$, $dg = (dg_n)$ be the difference sequences of f and g , defined by

$$f_n = \sum_{k=0}^n df_k, \quad g_n = \sum_{k=0}^n dg_k, \quad n = 0, 1, 2, \dots$$

Following Burkholder [5], we say that g is differentially subordinate to f , if

$$(1.1) \quad |dg_n| \leq |df_n|, \quad n = 0, 1, 2, \dots,$$

almost surely. For example, this takes place if g is a transform of f by a predictable real sequence $v = (v_n)$, bounded in absolute value by 1; that is, we have

2000 *Mathematics Subject Classification.* Primary: 60G42. Secondary: 31B05.

Key words and phrases. Martingale, harmonic function, differential subordination.

Partially supported by MEiN Grant 1 PO3A 012 29 and Foundation for Polish Science.

$dg_n = v_n df_n$, $\mathbb{P}(|v_n| \leq 1) = 1$ and v_n is measurable with respect to $\mathcal{F}_{(n-1) \vee 0}$, $n = 0, 1, 2, \dots$

Throughout the paper we assume that f and g are (\mathcal{F}_n) -martingales. The problem of comparing the sizes of f and g under the assumption of differential subordination has been studied in depth in the literature. For $p \in (0, \infty)$, let

$$\|f\|_p = \sup_n \|f_n\|_p = \sup_n (\mathbb{E}|f_n|^p)^{1/p},$$

$$\|f\|_{p,\infty} = \sup_n \|f_n\|_{p,\infty} = \sup_{\lambda>0} \lambda (\mathbb{P}(f^* \geq \lambda))^{1/p}$$

denote the strong and weak p -norms of a martingale. Here $f^* = \sup_n |f_n|$. For $1 < p < \infty$, let $p^* = \max\{p, p/(p-1)\}$. Let us start with the result by Burkholder [4].

Theorem 1.1. *Assume g is differentially subordinate to f .*

We have

$$(1.2) \quad \|g\|_{p,\infty} \leq \frac{2}{\Gamma(p+1)} \|f\|_p, \quad 1 \leq p \leq 2$$

and

$$(1.3) \quad \|g\|_p \leq (p^* - 1) \|f\|_p, \quad 1 < p < \infty.$$

Both constants $2/\Gamma(p+1)$ and $p^ - 1$ are optimal.*

One can check that neither of the estimates above holds for $p < 1$. The weak type inequality for the remaining set of parameters p was proved by Suh in [12]. Precisely,

Theorem 1.2. *Assume f and g are real valued and g is differentially subordinate to f . Then*

$$(1.4) \quad \|g\|_{p,\infty} \leq \left(\frac{p^{p-1}}{2}\right)^{1/p} \|f\|_p, \quad 2 \leq p < \infty.$$

The inequality is sharp.

If one imposes the additional assumption on the sign of the dominating martingale f , the optimal constants change for some values of p . Here is one of the main results of [8].

Theorem 1.3. *Assume g is differentially subordinate to f and f is nonnegative. Then*

$$\|g\|_p \leq C_p \|f\|_p, \quad 1 < p < \infty,$$

where the optimal constant C_p equals

$$C_p = \begin{cases} (p-1)^{-1} & \text{if } 1 < p \leq 2, \\ [p(\frac{p-1}{2})]^{1/p} & \text{if } 2 < p < \infty. \end{cases}$$

Hence the optimal constant in the moment inequalities (1.3) decreases if and only if $2 < p < \infty$. Furthermore, a closer inspection to the proof of (1.2) (see the example (4.24) on the page 657 in [4]) shows that the best constant in the inequality (1.2) for nonnegative martingales f is still $2/\Gamma(p+1)$. There is a natural question what can be said if $0 < p < 1$ or $p > 2$. The answer is contained in the following theorem.

Theorem 1.4. *Assume f is nonnegative, g is real valued and g is differentially subordinate to f . Then*

$$(1.5) \quad \|g\|_{p,\infty} \leq 2\|f\|_p, \quad 0 < p < 1$$

and

$$(1.6) \quad \|g\|_{p,\infty} \leq \frac{p}{2}(p-1)^{-1/p}\|f\|_p, \quad 2 \leq p < \infty.$$

The inequalities are sharp. They already sharp if g is assumed to be a transform of f .

Now let us turn to the continuous-time setting. Suppose $(\Omega, \mathcal{F}, \mathbb{P})$ is a complete probability space, equipped with a filtration $(\mathcal{F}_t)_{t \geq 0}$, such that \mathcal{F}_0 contains all the events of probability 0. Let $M = (M_t)$, $N = (N_t)$ be two real-valued semimartingales, which have right-continuous paths with limits from the left. The continuous-time extension of the differential subordination, which is due to Bañuelos and Wang (see [2] and [13]), can be formulated as follows. The semimartingale N is differentially subordinate to M , if the process $([M, M]_t - [N, N]_t)$ is nonnegative and nondecreasing. Here $([M, M]_t)$ denotes the quadratic variance process of M , see Delacherie and Meyer [10]. This notion is a generalization of (1.1). To see this, note that if one treats discrete-time sequences f, g as continuous time processes, then

$$[f, f]_t - [g, g]_t = \sum_{k=0}^{\lfloor t \rfloor} (|df_k|^2 - |dg_k|^2)$$

is nonnegative and nondecreasing if and only if (1.1) is valid.

As an example, assume X is a real-valued martingale and $K = (K_s)$, $H = (H_s)$ are predictable processes such that $|H| \leq |K|$ with probability 1. Let M, N be the Itô integrals of K, H , with respect to X ; that is,

$$M_t = K_0 X_0 + \int_0^t K_s dX_s, \quad N_t = H_0 X_0 + \int_0^t H_s dX_s.$$

Then, as

$$[M, M]_t - [N, N]_t = (|K_0|^2 - |H_0|^2)|X_0|^2 + \int_0^t (|K_s|^2 - |H_s|^2) d[X, X]_s,$$

we have that N is differentially subordinate to M .

All the results above have their counterparts in this new setting. For Theorem 1.1, see the paper [13] by Wang, where also a lot of information on transferring inequalities from discrete- to the continuous time setting can be found. Burkholder's method of proving martingale inequalities involves a construction of a special function, satisfying certain convexity-type properties. Once such a function is found, the continuous-time version follows from Itô's lemma and smoothing or stopping time argument. For other examples and discussion, see the papers by Bañuelos and Wang [2] and Suh [12].

Our approach follows the same pattern. To establish Theorem 1.4, we invent a special function and prove the following stronger result.

Theorem 1.5. *Assume M is a nonnegative martingale and N is differentially subordinate to M . Then*

$$(1.7) \quad \|N\|_{p,\infty} \leq 2\|M\|_p, \quad 0 < p < 1,$$

and

$$(1.8) \quad \|N\|_{p,\infty} \leq \frac{p}{2}(p-1)^{-1/p}\|M\|_p, \quad 2 < p < \infty,$$

and the inequalities are sharp.

We prove the case $p < 1$ in Section 2. Then we deal with the second part of the theorem. As the proof is quite complicated, we divide it into few steps. First, in Section 3, we show that the constant $p/(2(p-1)^{1/p})$ can not be replaced by a smaller one. Section 4 contains the study of a certain auxiliary differential equation, the solution of which will be needed in Section 5 in order to construct the special function. We complete the proof of Theorem 1.5 in Section 6.

In the second part of the paper we drop the condition $M \geq 0$ and deal with weak type estimates for differentially subordinated continuous-time martingales under the additional orthogonality assumption. We say that M and N are orthogonal, if their covariance process $[M, N]$ is constant with probability 1. Our result can be stated as follows.

Theorem 1.6. *Assume M, N are real-valued orthogonal martingales with N differentially subordinate to X . Then, for $1 \leq p \leq 2$,*

$$(1.9) \quad \|N\|_{p,\infty} \leq K_p \|M\|_p,$$

where

$$(1.10) \quad K_p^p = \frac{1}{\Gamma(p+1)} \cdot \frac{1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots}{1 - \frac{1}{3^{p+1}} + \frac{1}{5^{p+1}} - \frac{1}{7^{p+1}} + \dots}.$$

The inequality is sharp.

The theorem above for the particular case $p = 1$ was proved by Bañuelos and Wang in [3], using the ideas of Choi [9]. Their approach, again based on a construction of a special function, is analytic. We propose here a different proof, which is more probabilistic in nature. This is the contents of Section 7.

Finally, the last section of the paper is devoted to related results in harmonic analysis. As exhibited by Burkholder in [8] and [6] (consult also the paper [2] by Bañuelos and Wang for the orthogonal case), the inequalities for differentially subordinated martingales are accompanied by their analogues for harmonic functions on Euclidean domains. The Section 8 contains such extensions: the harmonic versions of the inequalities (1.7), (1.8) and (1.9).

2. THEOREMS 1.4 AND 1.5: THE CASE $0 < p < 1$

We start with an auxiliary lemma. Recall that for any semimartingale X there exists a unique continuous local martingale part X^c of X satisfying

$$[X, X]_t = |X_0|^2 + [X^c, X^c]_t + \sum_{0 < s \leq t} |\Delta X_s|^2$$

for all $t \geq 0$. Furthermore, $[X^c, X^c] = [X, X]^c$, the pathwise continuous part of $[X, X]$. Here is Lemma 1 from [13].

Lemma 2.1. *The process Y is differentially subordinate to X if and only if Y^c is differentially subordinate to X^c , the inequality $|\Delta Y_t| \leq |\Delta X_t|$ holds for all $t > 0$ and $|Y_0| \leq |X_0|$.*

Now let us introduce the special function $W : \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$, constructed by Burkholder in [7] to study the weak type inequality for nonnegative supermartingales. It is given by

$$W(x, y) = \begin{cases} 2x - x^2 + |y|^2 & \text{if } x + |y| \leq 1, \\ 1 & \text{if } x + |y| \geq 1. \end{cases}$$

The following functions $\phi : \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$, $\psi : \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$ will be needed later:

$$\begin{aligned} \phi(x, y) &= 2 - 2x, & \psi(x, y) &= 2y & \text{if } x + |y| \leq 1, \\ \phi(x, y) &= 0, & \psi(x, y) &= 0 & \text{if } x + |y| > 1. \end{aligned}$$

Note that ϕ and ψ coincide with the partial derivatives W_x, W_y except for the set $\{(x, y) : x + |y| = 1\}$. It can be shown (see page 1016 in [7]) that if $x \geq 0$, $x + h \geq 0$, $y, h \in \mathbb{R}$ and $|h| \leq |k|$, then

$$(2.1) \quad W(x + h, y + k) \leq W(x, y) + \phi(x, y)h + \psi(x, y)k.$$

Furthermore, we have

$$(2.2) \quad W(x, y) \geq 1_{\{x+|y|\geq 1\}}$$

and, if $|y| \leq x$, then

$$(2.3) \quad W(x, y) \leq (2x)^p.$$

Indeed, the inequality (2.2) is clear; to see (2.3), observe that it suffices to prove it for $|y| = x$ and then the inequality becomes $2x1_{\{2x < 1\}} + 1_{\{2x \geq 1\}} \leq (2x)^p$, which is immediate.

Proof of the inequality (1.7). We will prove a stronger statement: for any $\lambda > 0$,

$$\lambda^p \mathbb{P}((M + |N|)^* \geq \lambda) \leq 2^p \|M\|_p^p.$$

Here, as in the discrete-time case, $X^* = \sup_t |X_t|$. Obviously, we may assume $\lambda = 1$. Introduce the stopping time

$$\tau = \inf\{t : M_t + |N_t| \geq 1\}.$$

By Itô's formula,

$$(2.4) \quad W(M_{\tau \wedge t}, N_{\tau \wedge t}) = W(M_0, N_0) + I_1 + I_2 + I_3,$$

where

$$\begin{aligned} I_1 &= \frac{1}{2} \int_0^{\tau \wedge t} W_{xx}(M_{s-}, N_{s-}) d[M^c, M^c]_s + 2W_{xy}(M_{s-}, N_{s-}) d[M^c, N^c]_s \\ &\quad + W_{yy}(M_{s-}, N_{s-}) d[N^c, N^c]_s = -[M^c, M^c]_{\tau \wedge t} + [N^c, N^c]_{\tau \wedge t}, \\ I_2 &= \sum_{0 < s \leq \tau \wedge t} [W(M_s, N_s) - W(M_{s-}, N_{s-}) - W_x(M_{s-}, N_{s-}) \Delta M_s \\ &\quad - W_y(M_{s-}, N_{s-}) \Delta N_s], \\ I_3 &= \int_0^{\tau \wedge t} W_x(M_{s-}, N_{s-}) dM_s + \int_0^{\tau \wedge t} W_y(M_{s-}, N_{s-}) dN_s. \end{aligned}$$

Note that $I_1 \leq 0$, which is a consequence of Lemma 2.1. Moreover, as

$$W_x(M_{s-}, N_{s-}) = \phi(W_x(M_{s-}, N_{s-})) \text{ and } W_y(M_{s-}, N_{s-}) = \psi(W_x(M_{s-}, N_{s-})),$$

we have $I_2 \leq 0$: apply (2.1) to $x = M_{s-}$, $y = N_{s-}$, $h = \Delta M_s$, $k = \Delta N_s$ and observe that $|k| \leq |h|$ by Lemma 2.1. Finally, note that I_3 is a local martingale.

Therefore, there exists a sequence $(T_n)_{n=1}^\infty$ such that $T_n \uparrow \infty$ and with the further property that if we replace t with $T_n \wedge t$ ($n = 1, 2, \dots$) in the expression defining I_3 , then this expression has zero expectation. Combining this with the previous observations about I_1 and I_2 , we see that (2.4) gives

$$\mathbb{E}W(M_{\tau \wedge T_n \wedge t}, N_{\tau \wedge T_n \wedge t}) \leq \mathbb{E}W(M_0, N_0).$$

Now we let $n \rightarrow \infty$. As $0 \leq W \leq 1$, the Lebesgue's dominated convergence theorem gives

$$\mathbb{E}W(M_{\tau \wedge t}, N_{\tau \wedge t}) \leq \mathbb{E}W(M_0, N_0).$$

Apply (2.2) and (2.3) to obtain

$$\mathbb{P}(M_{\tau \wedge t} + |N_{\tau \wedge t}| \geq 1) \leq 2^p \mathbb{E}M_0^p = 2^p \|M\|_p^p.$$

To conclude the proof, fix $\varepsilon > 0$ and consider processes $M^\varepsilon = M(1+\varepsilon)$, $N^\varepsilon = N(1+\varepsilon)$. Clearly, N^ε is differentially subordinate to M^ε , so, if $\eta = \inf\{t : M_t^\varepsilon + |N_t^\varepsilon| \geq 1\}$, we get, by the above argumentation,

$$\begin{aligned} \mathbb{P}((M + |N|)^* \geq 1) &= \mathbb{P}((M^\varepsilon + |N^\varepsilon|)^* \geq (1 + \varepsilon)) \leq \lim_{t \rightarrow \infty} \mathbb{P}(M_{\eta \wedge t}^\varepsilon + |N_{\eta \wedge t}^\varepsilon| \geq 1) \\ &\leq 2^p \|M^\varepsilon\|_p^p = 2^p \|M\|_p^p (1 + \varepsilon)^p. \end{aligned}$$

As ε was arbitrary, the proof is complete. \square

Sharpness of (1.5): Consider the following example. Assume the probability space is the interval $[0, 1]$ with Lebesgue measure. Let $f_0 = g_0 \equiv 1/2$ and

$$df_1 = -dg_1 = -\frac{1}{2}[0, 3/4] + \frac{3}{2}(3/4, 1]$$

and $df_n = dg_n = 0$ for $n \geq 2$. Here and in the next section, we identify a set with its indicator function. Then $|g_1| = 1$ with probability 1 and hence

$$\|g\|_{p, \infty} \geq 1 = 2\|f_0\|_p = 2\|f\|_p,$$

as needed. \square

3. SHARPNESS OF (1.4) AND (1.5)

The optimality of the constant will be proved by constructing appropriate example. Let $p > 2$ be a fixed number. Let $\delta > 0$, $x \in (0, 1/p)$ be numbers satisfying

$$(3.1) \quad x \left(1 + \frac{2\delta}{p}\right)^N = \frac{1}{p}$$

for certain integer $N = N(\delta, x)$. It is clear that we may choose δ and x to be arbitrarily small.

Consider a two-dimensional Markov martingale $(X_n, Y_n) = (X_n^{x, \delta}, Y_n^{x, \delta})$, uniquely determined by the following properties.

- (i) $X_0 = x$, $Y_0 = (p-1)x$.
- (ii) We have $dX_n = (-1)^{n+1} dY_n$ for $n = 1, 2, \dots$
- (iii) If (X_n, Y_n) lies on the line $y = (p-1)x$ and $X_n < 1/p$, then in the next step it moves either to the line $x = 0$, or to the line $y = (p-1+\delta)x/(1+\delta)$, $n = 0, 1, 2, \dots$
- (iv) If (X_n, Y_n) lies on the line $y = (p-1+\delta)x/(1+\delta)$ then in the next step it moves either to the line $y = (p-1)x$, or to the line $y = (p-2)x/2$, $n = 0, 1, 2, \dots$
- (v) If $(X_n, Y_n) = (1/p, 1-1/p)$ (which happens only if $n = 2N$) then (X_{n+1}, Y_{n+1}) equals either $(0, 1)$, or $(2/p, 1-2/p)$.

(vi) The states on the line $x = 0$ and $y = (p - 2)x/2$ are absorbing. The exemplary trajectories are presented on Figure 1.

FIGURE 1. Four trajectories of the process (X, Y) corresponding to the parameters $p = 3$, $x = \frac{1}{24}$, $\delta = \frac{3}{2}$ and $N = 3$.

To be more precise, let the sequence (p_n) , $n = 0, 1, 2, \dots, 2N$ be given by

$$p_{2n} = \left(\frac{p - p\delta + 4\delta}{(p + 2\delta)(1 + \delta)} \right)^n, \quad p_{2n+1} = p_{2n} \cdot \frac{1}{1 + \delta}, \quad n = 0, 1, 2, \dots, N.$$

Let the probability space be the interval $[0, 1]$ with Lebesgue measure. Set

$$(3.2) \quad X_0 = x[0, 1], \quad dX_{2n+1} = \delta X_{2n}[0, p_{2n+1}] - X_{2n}(p_{2n+1}, p_{2n}], \\ dX_{2n+2} = -(1 - 2/p)\delta X_{2n}[0, p_{2n+2}] + X_{2n}(1 + \frac{4}{p}\delta - \delta)(p_{2n+2}, p_{2n+1}]$$

for $n = 0, 1, 2, \dots, N - 1$, and

$$(3.3) \quad dX_{2N+1} = X_{2N}[0, p_{2N}/2] - X_{2N}(p_{2N}/2, p_{2N}].$$

Furthermore,

$$Y_0 = (p - 1)x[0, 1], \quad dY_n = (-1)^{n+1}dX_n, \quad n = 0, 1, 2, \dots, 2N + 1.$$

Note that $Y_{2N+1} = 1$ on $[0, p_{2N}/2]$ and $0 < Y_{2N+1} < 1$ on $(p_{2N}/2, 1]$, so we have $\mathbb{P}(Y^* \geq 1) = p_{2N}/2$. Furthermore, by (3.2) and (3.3), X_{2N+1} equals 0 on the union of the sets $(p_{2n+1}, p_{2n}]$, $n = 0, 1, \dots, N$ and the interval $(p_{2N}/2, p_{2N}]$. Moreover, X_{2N+1} is equal to

$$2 \left(1 + \frac{2\delta}{p} \right)^{n+1} x \quad \text{on} \quad (p_{2n+2}, p_{2n+1}], \quad n = 0, 1, \dots, N - 1.$$

Finally, it equals $2/p$ on $[0, p_{2N}/2]$. Hence we may write

$$\begin{aligned} \mathbb{E}X_{2N+1}^p &= \frac{2^p(p-2)\delta}{(p+2\delta)(1+\delta)} \left(1 + \frac{2\delta}{p} \right)^p x^p \sum_{k=0}^{N-1} \left(\frac{p-p\delta+4\delta}{(p+2\delta)(1+\delta)} \left(1 + \frac{2\delta}{p} \right)^p \right)^k \\ &\quad + \left(\frac{2}{p} \right)^p \cdot \frac{1}{2} \left(\frac{p-p\delta+4\delta}{(p+2\delta)(1+\delta)} \right)^N \\ &= \frac{2^p(p-2)\delta}{(p+2\delta)(1+\delta)} \left(1 + \frac{2\delta}{p} \right)^p x^p \frac{\left(\frac{p-p\delta+4\delta}{(p+2\delta)(1+\delta)} \left(1 + \frac{2\delta}{p} \right)^p \right)^N - 1}{\frac{p-p\delta+4\delta}{(p+2\delta)(1+\delta)} \left(1 + \frac{2\delta}{p} \right)^p - 1} \\ &\quad + \left(\frac{2}{p} \right)^p \cdot \frac{1}{2} \left(\frac{p-p\delta+4\delta}{(p+2\delta)(1+\delta)} \right)^N. \end{aligned}$$

Now keep x fixed and let δ to 0 (so that (3.1) holds, with $N = N(\delta, x) \rightarrow \infty$). Then

$$\left(\frac{p-p\delta+4\delta}{(p+2\delta)(1+\delta)} \right)^N = \left(1 + \frac{2\delta}{p} \cdot \frac{p(1-p-\delta)}{(p+2\delta)(1+\delta)} \right)^N \rightarrow (px)^{p-1}$$

and

$$\frac{\delta}{\frac{p-p\delta+4\delta}{(p+2\delta)(1+\delta)} \left(1 + \frac{2\delta}{p} \right)^p - 1} \rightarrow \frac{p}{2},$$

so we have

$$\begin{aligned}\mathbb{E}X_{2N+1}^p &\rightarrow \frac{2^p(p-2)}{p} x^p \frac{((px)^{p-1}(px)^{-p} - 1)p}{2} + \left(\frac{2}{p}\right)^p \cdot \frac{1}{2} (px)^{p-1} \\ &= \frac{(2x)^{p-1}}{p} (p-1 - p(p-2)x)\end{aligned}$$

and

$$(3.4) \quad \frac{\mathbb{P}(Y_{2N+1} \geq 1)}{\mathbb{E}X_{2N+1}^p} \rightarrow \frac{\frac{1}{2}(px)^{p-1}}{\frac{(2x)^{p-1}}{p} (p-1 - p(p-2)x)} = \frac{p^p}{2^p(p-1) - 2^p p(p-2)x}.$$

Observe that Y is a transform of X and $|dX_n| = |dY_n|$ for $n \geq 1$. However, Y is not differentially subordinate to X as $Y_0 = (p-1)X_0 > X_0$. To overcome this difficulty, introduce the processes $Y'_n = (Y_n - (p-2)x)/(1 - (p-2)x)$, $X'_n = X_n/(1 - (p-2)x)$, $n = 0, 1, \dots, 2N+1$. Then Y' is a transform of X' by the deterministic sequence $(1, 1, -1, 1, -1, 1, -1, 1, \dots)$ and $\mathbb{P}(Y'_{2N+1} \geq 1) = \mathbb{P}(Y_{2N+1} \geq 1)$. In terms of these new processes, (3.4) reads

$$\frac{\mathbb{P}(Y'_{2N+1} \geq 1)}{\mathbb{E}(X'_{2N+1})^p} \rightarrow \frac{p^p(1 - x(p-2))^p}{2^p(p-1) - 2^p p(p-2)x}$$

and it is clear that the limit can be made arbitrarily close to $p^p/2^p(p-1)$ by choosing x sufficiently small. This proves the sharpness of (1.6) and hence the sharpness of (1.8) as well.

4. A DIFFERENTIAL EQUATION

Let $p > 2$ be fixed. The purpose of this section is to study a solution to a certain differential equation. A very similar equation appears in [12] and our arguments are parallel to those used there. We will show that there exists a function $h : [1, \infty) \rightarrow [2/p, \infty)$, which enjoys the following properties.

(4.1) the function h is increasing and continuous on $[1, \infty)$,

(4.2) $h(t) > t - 1$ for all $t \geq 1$,
 h is differentiable on $(1, \infty)$ and

(4.3) $h'(t) = \left(\frac{2}{p}\right)^{p+1} (h(t))^{2-p} (h(t) - t + 1)^{-2}$,

(4.4) $h(1) = h'(1+) = 2/p$.

The problem above is equivalent to the existence of the function $G : [2/p, \infty) \rightarrow [1, \infty)$ satisfying the following properties:

(4.5) the function G is increasing and continuous on $[2/p, \infty)$,

(4.6) $G(t) < t + 1$ for all $t \geq 2/p$,
 G is differentiable on $(2/p, \infty)$ and

(4.7) $G'(t) = \left(\frac{p}{2}\right)^{p+1} t^{p-2} (t + 1 - G(t))^2$.

(4.8) $G\left(\frac{2}{p}\right) = 1$, $G'\left(\frac{2}{p}+\right) = \frac{p}{2}$.

To see the equivalence, note that if h satisfies (4.1)–(4.4), then $G = h^{-1}$ satisfies (4.5)–(4.8), and if G satisfies (4.5)–(4.8), then $h = G^{-1}$ satisfies (4.1)–(4.4).

As (4.7) has Riccati form, we can use the transformation

$$k(t) = \exp \left[\int_{2/p}^t \left(\frac{p}{2}\right)^{p+1} y^{p-2} (y+1 - G(y)) dy \right]$$

to obtain the following differential equation for k :

$$(4.9) \quad yk''(y) + (2-p)k'(y) - \left(\frac{p}{2}\right)^{p+1} y^{p-1}k(y) = 0.$$

For a fixed $\alpha > -1$, let I_α be the modified Bessel function of the first kind (see Abramowitz and Stegun [1]). That is,

$$I_\alpha(z) = \sum_{k=0}^{\infty} \frac{(z/2)^{2k+\alpha}}{k!!\Gamma(\alpha+k+1)}$$

and we have

$$z^2 I_\alpha''(z) + z I_\alpha'(z) - (z^2 + \alpha^2) I_\alpha(z) = 0.$$

One can check that the functions

$$k_1(t) = t^{(p-1)/2} I_{-\frac{p-1}{p}}(z_0),$$

$$k_2(t) = t^{(p-1)/2} I_{\frac{p-1}{p}}(z_0),$$

where $z_0 = \sqrt{(p/2)^{p-1} t^p}$, are two linearly independent solutions on $(0, \infty)$ to the equation (4.9). As the functions I_α are infinitely differentiable on $(0, \infty)$, so are k_1 and k_2 . Let a_1, a_2 be two numbers such that $k = a_1 k_1 + a_2 k_2$ satisfies

$$(4.10) \quad k(2/p) = 1 \quad \text{and} \quad k'(2/p) = \frac{p^2}{4}.$$

If one rewrites (4.9) in the form

$$yk''(y) = (p-2)k'(y) + \left(\frac{p}{2}\right)^{p+1} y^{p-1}k(y),$$

then it follows from (4.10) that k, k' and k'' are strictly positive on $[2/p, \infty)$. Now it is straightforward to check that the function

$$(4.11) \quad G(t) = t + 1 - \left(\frac{2}{p}\right)^{p+1} \frac{k'(t)}{k(t)t^{p-2}}$$

has all the properties (4.5)–(4.8).

We conclude this section by

Lemma 4.1. *We have*

$$(4.12) \quad h'(t) \leq 1 \quad \text{for all } t > 1.$$

Proof. This is equivalent to $G'(t) \geq 1$ for all $t > 2/p$. We have

$$G''(t) = \left(\frac{p}{2}\right)^{p+1} [(p-2)t^{p-3}(t+1-G(t))^2 + 2t^{p-2}(t+1-G(t))(1-G'(t))],$$

implying that if $G'(t) \leq 1$, then $G''(t) > 0$. Now suppose $G'(t_0) < 1$ for some $t_0 > 2/p$. Then, as $G'(2/p+) = p/2 > 1$, there exists $t_1 \in (2/p, t_0)$ such that

$G'(t_1) = 1$ and $G'(t) < 1$ for $t \in (t_1, t_0)$. Now by mean value property, for some $t_2 \in (t_1, t_0)$,

$$G''(t_2) = \frac{G'(t_0) - G'(t_1)}{t_0 - t_1} < 0,$$

a contradiction. \square

5. THE SPECIAL FUNCTION AND ITS PROPERTIES

Now we are ready to define the special function U . Due to the lack of symmetry with respect to the y axis, this function is much more complicated than the one constructed in [12]. Consider the following subsets of $\mathbb{R}_+ \times \mathbb{R}$.

$$\begin{aligned} D_0 &= \{(x, y) : |y| \geq 1\}, \\ D_1 &= \left\{ (x, y) : (p-1)x \leq |y| < x + 1 - \frac{2}{p} \right\}, \\ D_2 &= \left\{ (x, y) : \frac{p-2}{2}x \leq |y| < \min\{1-x, (p-1)x\} \right\}, \\ D_3 &= \left\{ (x, y) : x + 1 - \frac{2}{p} \leq y < 1 - x \right\}, \\ D_4 &= \left\{ (x, y) : \max\left\{1-x, \left(x + 1 - \frac{2}{p}\right)\right\} \leq |y| < 1 \right\}, \\ D_5 &= \left\{ (x, y) : h(x + |y|) \geq x > \frac{-1 + h(x + |y|) + x + |y|}{2} \text{ and } x + |y| \geq 1 \right\}, \\ D_6 &= \left\{ (x, y) : \frac{1 - h(x + |y|) + x + |y|}{2} \leq |y| < \min\left\{\left(x + 1 - \frac{2}{p}\right), 1\right\} \right\}, \\ D_7 &= (\mathbb{R}_+ \times \mathbb{R}) \setminus (D_0 \cup D_1 \cup D_2 \cup D_3 \cup D_4 \cup D_5 \cup D_6). \end{aligned}$$

See Figure 2 for the case $p = 3$.

FIGURE 2. The regions $D_0 - D_7$ for $p = 3$, intersected with $\mathbb{R}_+ \times \mathbb{R}_+$.

Let

$$V(x, y) = \begin{cases} 1 - \frac{p^p}{2^p(p-1)}x^p & \text{on } D_0 \\ -\frac{p^p}{2^p(p-1)}x^p & \text{on } (\mathbb{R}_+ \times \mathbb{R}) \setminus D_0 \end{cases}$$

and let us define U by

$$\begin{cases} 1 - \frac{p^p}{2^p(p-1)}x^p & \text{on } D_0, \\ \frac{p^p}{2(p-1)(p-2)^{p-2}}x(|y| - x)^{p-1} & \text{on } D_1, \\ \frac{1}{p-1}(x + |y|)^{p-1} \left[(p-1)|y| - \frac{p^2-2p+2}{2}x \right] & \text{on } D_2, \\ \frac{x}{2(p-1)(1+x-|y|)} \left[-(p-2)^2 + p^2(|y| - x) \right] & \text{on } D_3, \\ 1 - \frac{p^2}{2(p-1)}(1 - |y|) - \frac{p^p}{2^p(p-1)}(x + |y| - 1)(x + 1 - |y|)^{p-1} & \text{on } D_4, \\ \frac{p^p}{2^p(p-1)}(h(x + |y|))^{p-1} [(p-1)h(x + |y|) - px] & \text{on } D_5, \\ 1 - \frac{2(1-|y|)}{2+x-|y|-G(x-|y|+1)} - \frac{p^p}{2^p(p-1)}(x - |y| + 1)^{p-1} [x - (p-1)(1 - |y|)] & \text{on } D_6, \\ -\frac{p^p}{2^p(p-1)}x^p & \text{on } D_7. \end{cases}$$

The properties of U are described in the sequence of the lemmas below.

Lemma 5.1. *The function U is continuous on $\mathbb{R}_+ \times \mathbb{R} \setminus \{(0, \pm 1)\}$ and of class C^1 except for the set $\partial D_0 \cup (\partial D_3 \cap \partial D_4 \setminus (\frac{2}{p}, \pm \frac{p-2}{p}))$.*

Proof. Clearly, U is of class C^1 in the interior of each D_j , $j = 0, 1, 2, \dots, 7$, so we need to check the properties on the boundaries. By symmetry, we may restrict ourselves to positive y 's. Using (4.4), (4.8) and the definitions of the boundaries, the continuity of U can be verified readily. For the second part of the lemma, we calculate the partial derivatives of U : we have

$$U_x(x, y) = \begin{cases} -\frac{p^{p+1}}{2^p(p-1)}x^{p-1} & \text{on } D_0^o, \\ \frac{p^p}{2(p-1)(p-2)^{p-2}}(y-x)^{p-2}(y-px) & \text{on } D_1^o, \\ \frac{p}{2(p-1)}(x+y)^{p-2}[(p-2)y - (p^2 - 2p + 2)x] & \text{on } D_2^o, \\ -\frac{p^2}{2(p-1)} + \frac{2(1-y)}{(1+x-y)^2} & \text{on } D_3^o, \\ -\frac{p^p}{2^p(p-1)}(x+1-y)^{p-2}[px - (p-2)(1-y)] & \text{on } D_4^o, \\ \frac{2(h(x+y)-x)}{(h(x+y)-(x+y)+1)^2} - \frac{p^{p+1}}{2^p(p-1)}(h(x+y))^{p-1} & \text{on } D_5^o, \\ \frac{2(1-|y|)}{(2+x-|y|-G(x-|y|+1))^2} - \frac{p^{p+1}}{2^p(p-1)}(x-|y|+1)^{p-1} & \text{on } D_6^o, \\ -\frac{p^{p+1}}{2^p(p-1)}x^{p-1} & \text{on } D_7^o, \end{cases}$$

while

$$U_y(x, y) = \begin{cases} 0 & \text{on } D_0^o, \\ \frac{p^p}{2(p-2)^{p-2}}x(y-x)^{p-2} & \text{on } D_1^o, \\ \frac{p}{2}(x+y)^{p-2}[2y - (p-2)x] & \text{on } D_2^o, \\ \frac{2x}{(1+x-y)^2} & \text{on } D_3^o, \\ \frac{p^2}{2(p-1)} + \frac{p^p}{2^p(p-1)}(x+1-y)^{p-2}[(p-2)x - p(1-y)] & \text{on } D_4^o, \\ \frac{2(h(x+y)-x)}{(h(x+y)-(x+y)+1)^2} & \text{on } D_5^o, \\ \frac{2}{(2+x-y-G(x-y+1))^2}[1+x-G(x-y+1)] & \text{on } D_6^o, \\ 0 & \text{on } D_7^o. \end{cases}$$

All that is left is to check that the partial derivatives agree on the boundaries. \square

Lemma 5.2. *For $y \geq 0$, U_y is nonnegative.*

Proof. This is clear on D_1^o , D_2^o , D_3^o , D_5^o and D_7^o . On D_4 , we have

$$U_{xy}(x, y) = \frac{p^p(p-2)}{2^p}(x+1-y)^{p-3}(x-y) \leq 0$$

and, consequently,

$$U_y(x, y) \geq U_y\left(y-1 + \frac{2}{p}, y\right) = \frac{p^2}{2} \left(y-1 + \frac{2}{p}\right) \geq 0.$$

On D_6 , we have $G(1+x-y) \leq x+y$ (as it is equivalent to $1+x-y \leq h(x+y)$, one of the inequalities defining D_6). This can be further bounded from above by $1+x$, which yields the claim. \square

The most technical lemma is the following.

Lemma 5.3. *Suppose (x, y) belongs to the interior of D_j for some $0 \leq j \leq 7$. Then for any h, k we have*

$$(5.1) \quad U_{xx}(x, y)h^2 + 2U_{xy}(x, y)hk + U_{yy}(x, y)k^2 \leq 0.$$

Proof. We start with the observation that the inequality holds if (x, y) belongs to D_0^o or D_7^o ; indeed, $U_{xy} = U_{yy} = 0$ and $U_{xx} \leq 0$ there. For (x, y) lying in the interior of one of the remaining sets, note that U has the following property: one of the functions $t \mapsto U(x+t, y+t)$, $t \mapsto U(x+t, y-t)$, is linear on some neighbourhood of 0. Hence

$$(5.2) \quad U_{xx}(x, y) + 2U_{xy}(x, y) + U_{yy}(x, y) = 0$$

or

$$(5.3) \quad U_{xx}(x, y) - 2U_{xy}(x, y) + U_{yy}(x, y) = 0.$$

Now if (5.2) holds, we may write

$$\begin{aligned} & U_{xx}(x, y)h^2 + 2U_{xy}(x, y)hk + U_{yy}(x, y)k^2 \\ &= \frac{U_{xx}(x, y) + U_{yy}(x, y)}{2}(h-k)^2 + \frac{U_{xx}(x, y) - U_{yy}(x, y)}{2}(h^2 - k^2), \end{aligned}$$

while if (5.3) is valid, we have

$$\begin{aligned} & U_{xx}(x, y)h^2 + 2U_{xy}(x, y)hk + U_{yy}(x, y)k^2 \\ &= \frac{U_{xx}(x, y) + U_{yy}(x, y)}{2}(h+k)^2 + \frac{U_{xx}(x, y) - U_{yy}(x, y)}{2}(h^2 - k^2). \end{aligned}$$

Therefore (5.1) will hold once we have established the inequality

$$(5.4) \quad U_{xx} \leq -|U_{yy}|.$$

As previously, with no loss of generality may assume $y > 0$.

Straightforward computations show that

$$U_{xx}(x, y) = \begin{cases} \frac{p^p}{2(p-2)^{p-2}}(y-x)^{p-3}(px-2y) & \text{on } D_1^o, \\ -p(x+y)^{p-3} \left(\frac{p^2-2p+2}{2}x+y \right) & \text{on } D_2^o, \\ -\frac{4}{(1+x-y)^3}(1-y) & \text{on } D_3^o, \\ -\frac{p^p}{2^p}(x+1-y)^{p-3}(px+(p-4)(y-1)) & \text{on } D_4^o, \\ \frac{2}{(h(x+y)-(x+y)+1)^2} \left[-2+h'(x+y) - \frac{2(h(x+y)-x)(h'(x+y)-1)}{h(x+y)-(x+y)+1} \right] & \text{on } D_5^o, \\ -\frac{4(1-y)(1-G'(x-y+1))}{(2+x-y-G(x-y+1))^3} - \frac{p^{p+1}}{2^p}(x-y+1)^{p-2} & \text{on } D_6^o \end{cases}$$

and

$$U_{yy}(x, y) = \begin{cases} \frac{p^p}{2(p-2)^{p-2}}(y-x)^{p-3} \cdot (p-2)x & \text{on } D_1^o, \\ -p(x+y)^{p-3} \left(\frac{p^2-4p+2}{2}x - (p-1)y \right) & \text{on } D_2^o, \\ \frac{4}{(1+x-y)^3}x & \text{on } D_3^o, \\ \frac{p^p}{2^p}(x+1-y)^{p-3}(-(p-4)x+p(1-y)) & \text{on } D_4^o, \\ \frac{2}{(h(x+y)-(x+y)+1)^2} \left[-h'(x+y) - \frac{2(h(x+y)-x)(h'(x+y)-1)}{h(x+y)-(x+y)+1} \right] & \text{on } D_5^o, \\ -\frac{4(1-y)(1-G'(x-y+1))}{(2+x-y-G(x-y+1))^3} + \frac{2(2-G'(x-y+1))}{(2+x-y-G(x-y+1))^2} & \text{on } D_6^o. \end{cases}$$

Now let us check (5.4). On D_1^o it is equivalent to

$$px - 2y \leq -(p-2)x \text{ or } y \geq (p-1)x,$$

which follows from the definition of D_1 . On D_2^o , the inequalities $U_{xx} + U_{yy} \leq 0$, $U_{xx} - U_{yy} \leq 0$ can be transformed to $-p(x+y) \leq 0$ and $(p-2)(y - (p-1)x) \leq 0$,

respectively, which are valid. On D_3^o the inequality is verified easily. On D_4^o , the estimates $U_{xx} + U_{yy} \leq 0$, $U_{xx} - U_{yy} \leq 0$ are equivalent to

$$2(p-2)(1-x-y) \leq 0 \quad \text{and} \quad -4(1-y+x) \leq 0,$$

respectively, which hold true. On D_5^o it is obvious that $U_{xx} \leq U_{yy}$, while the inequality $U_{xx} \leq -U_{yy}$ reduces to

$$\frac{(h'(x+y)-1)(-h(x+y)+x-y+1)}{h(x+y)-(x+y)+1} \leq 0,$$

which is guaranteed by (4.12) and the definition of D_5 . Finally, assume $(x, y) \in D_6^o$. Then, by (4.7),

$$\frac{p^{p+1}}{2^p}(x-y+1)^{p-2} = \frac{2G'(x-y+1)}{(2+x-y-G(x-y+1))^2}$$

and it is easy to see that $U_{xx}(x, y) \leq U_{yy}(x, y)$. The inequality $U_{xx}(x, y) \leq -U_{yy}(x, y)$ is equivalent to

$$\frac{4(1-G'(x-y+1))}{(2+x-y-G(x-y+1))^3}(x+y-G(x-y+1)) \leq 0.$$

To prove its validity, note that $G' \geq 1$, which is a consequence of (4.12), and $x+y \geq G(x-y+1)$, which is equivalent to $h(x+y) \geq x-y+1$, one of the inequalities in the definition of D_6 .

The proof is complete. \square

As in the proof of the case $p < 1$, we extend the partial derivatives of the special function to the whole $\mathbb{R}_+ \times \mathbb{R}$. Let $\phi, \psi : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$ be given by

$$\phi(x, y) = \begin{cases} U_x(x, y) & \text{if } (x, y) \notin \partial D_3 \cap \partial D_4, \\ U_x(x+, y) & \text{if } (x, y) \in \partial D_3 \cap \partial D_4, \end{cases}$$

$$\psi(x, y) = \begin{cases} U_y(x, y) & \text{if } (x, y) \notin \partial D_0 \cup (\partial D_3 \cap \partial D_4), \\ U_y(x+, y) & \text{if } (x, y) \in \partial D_3 \cap \partial D_4 \setminus \{(0, 0)\}, \\ U_y(x, y+) & \text{if } (x, y) \in \partial D_0 \end{cases}$$

and extend them to the whole $\mathbb{R} \times \mathbb{R}$ by $\phi(x, y) = \phi(x, -y)$, $\psi(x, y) = -\psi(x, -y)$.

The further properties of U are described in

Lemma 5.4. (i) Let $x \geq 0$, $x+h \geq 0$ and $y, k \in \mathbb{R}$. Then

$$(5.5) \quad U(x+h, y+k) \leq U(x, y) + \phi(x, y)h + \psi(x, y)k.$$

(ii) Let $x \in \mathbb{R}_+$ and $y \in (-1, 1)$. Then the function $H_{x,y}$ defined on $\{t : x+t \geq 0 \text{ and } -1 < y+t < 1\}$, given by

$$H_{x,y}(t) = \phi(x+t, y+t) - \psi(x+t, y+t)$$

is nonincreasing.

Proof. (i) Consider a continuous function $L = L_{x,y,h,k}$ defined on $\{t : x+th \geq 0\}$, given by

$$L(t) = u(x+th, y+tk).$$

The inequality (5.5) is equivalent to $L(1) \leq L(0) + L'(0)$ (with $L'(0)$ replaced by left- or right-sided derivative if (x, y) belongs to ∂D_0 or $\partial D_3 \cap \partial D_4$) and will follow if we show that L is concave. To this end, it suffices to prove that $L''(t) \leq 0$ for

those t , where the second derivative exists, and $L'(t-) \geq L'(t+)$ for remaining t , satisfying $x + th > 0$. The first inequality follows from

$$L''(t) = U_{xx}(x + th, y + tk)h^2 + 2U_{xy}(x + th, y + tk)hk + U_{yy}(x + th, y + tk)k^2 \leq 0,$$

due to (5.1). To deal with the second, recall that L is of class C^1 except for $(x + th, t + tk)$ belonging to ∂D_0 or $\partial D_3 \cap \partial D_4$. Hence, by the transity property $L_{x,y,h,k}(t+s) = L_{x+th,y+tk,h,k}(s)$, all we need is $L'(0-) \geq L'(0+)$ for (x, y) belonging to one of these sets. As $L_{x,y,h,k}$ is concave if and only if $L_{x,y,-h,-k}$ is concave, we may also assume $h > 0$. Now if $(x, y) \in \partial D_0$, then U_x is continuous in (x, y) and

$$L'(0-) - L'(0+) = (U_y(x, y-) - U_y(x, y+))|k| = U_y(x, y-)|k| \geq 0,$$

by Lemma 5.2. Suppose then, that $(x, y) \in \partial D_3 \cap \partial D_4$. We have

$$(5.6) \quad L'(0-) - L'(0+) = \left(-\frac{p^2}{2(p-1)} + \frac{1}{2x} + \frac{p^p}{2(p-1)}x^{p-1} \right) (h+k) \geq 0.$$

The latter inequality is a consequence of $h \geq k$ and

$$-\frac{p^2}{2(p-1)} + \frac{1}{2x} + \frac{p^p}{2(p-1)}x^{p-1} = \frac{1}{2x(p-1)} [p(1-px) - (1-(px)^p)] \geq 0,$$

which follows from the mean value property.

(ii) If $(x+t, y+t)$ lies in the interior of one of the sets D_k , $k = 1, 2, \dots, 7$, then

$$H'_{x,y}(t) = U_{xx}(x+t, y+t) - U_{yy}(x+t, y+t) \leq 0.$$

Therefore we will be done if we show that $H_{x,y}$ is continuous. By Lemma 5.1, we only need to check continuity for t determined by $(x+t, y+t) \in \partial D_3 \cap \partial D_4$, $(x+t, y+t) \neq (2/p, \pm(p-2)/p)$. If $y+t > 0$, then one can check, using the formulae for U_x, U_y , that

$$\lim_{t' \rightarrow t} H_{x,y}(t') = \lim_{t' \rightarrow t} [U_x(x+t', y+t') - U_y(x+t', y+t')] = -\frac{p^2}{2(p-1)} = H_{x,y}(t).$$

For $y+t < 0$ this follows from the fact that $H_{x,y}(s) = U_x(x+s, y+s) - U_y(x+s, y+s)$ for s lying in some neighborhood of t (both the partial derivatives are defined by the formulae for D_4 and hence are continuous). \square

Lemma 5.5. (i) For any $x \geq 0, y \in \mathbb{R}$ satisfying $|y| \leq x$ we have $U(x, y) \leq 0$.

(ii) We have $U \geq V$.

Proof. (i) Using (5.5), we may write

$$U(x, y) \leq U(0, 0) + \phi(0, 0)x + \psi(0, 0)y = 0,$$

as claimed.

(ii) The inequality is clear on D_0 . For $(x, y) \notin D_0$, use Lemma 5.2 to obtain

$$U(x, y) \geq U(x, 0) = V(x, 0) = V(x, y).$$

This finishes the proof. \square

6. THE PROOFS OF THE INEQUALITIES (1.6) AND (1.8)

For the sake of convenience, the proof is divided into a few steps.

Step 1. We start with a smoothing argument and correct the function U in such a way that the key properties are still valid (the inequalities (6.1), (6.2) and (6.3) below). Let $\varepsilon > 0$ be fixed and m be a positive integer satisfying $1/m < \varepsilon$. Let $g_m : \mathbb{R}^2 \rightarrow \mathbb{R}_+$ be a C^∞ function with support inside the ball B_m centered at 0 and radius $1/m$. Furthermore, assume g_m has integral 1 and define $U^m : [1/m, \infty) \times \mathbb{R} \rightarrow \mathbb{R}$ as the convolution of U with the function g_m . Note that U^m is infinitely differentiable. Moreover, by Lemma 5.4 (i), U^m is concave along the lines of slope not greater than 1 in absolute value, as convolving with a positive function does not affect this property. Therefore, we have

$$(6.1) \quad U_{xx}^m \pm 2U_{xy}^m + U_{yy}^m \leq 0.$$

and, for $x, x+h > 1/m, y, k \in \mathbb{R}$ such that $|h| \geq |k|$,

$$(6.2) \quad U^m(x+h, y+k) \leq U^m(x, y) + U_x^m(x, y)h + U_y^m(x, y)k.$$

Furthermore, by part (ii) of this lemma, for any $x > 1/m, |y| < 1 - 1/m$, the function $H_{x,y}^m$, defined on a (small) neighborhood of 0 by

$$H^m(t) = U_x^m(x+t, y+t) - U_y^m(x+t, y+t),$$

is nonincreasing. To see this, note that by integration by parts and continuity of U on $(0, \infty) \times \mathbb{R}$, we have

$$H_{x,y}^m(t) = \int_{B_m} H_{x-u, y-v}(t) g_m(u, v) du dv$$

(here H under the integral is defined as in Lemma 5.4 (ii)). Thus

$$(6.3) \quad (H_{x,y}^m)'(0) = U_{xx}^m(x, y) - U_{yy}^m(x, y) \leq 0 \quad \text{for } x > 1/m, |y| < 1 - 1/m.$$

Step 2. Here we will introduce the key stopping time. With no loss of generality we may assume $\|M\|_p < \infty$. For $\eta < 1$, let $R = R(\eta, \varepsilon)$ denote the greatest number r such that

$$(6.4) \quad U(x, y) \geq \eta 1_{\{|y| \geq 1-\varepsilon\}} - \frac{p^p}{2^p(p-1)} x^p \quad \text{for all } x \leq r, y \in \mathbb{R}.$$

Note that if η is fixed and $\varepsilon \downarrow 0$, then $R(\eta, \varepsilon) \rightarrow \infty$. This is a consequence of Lemma 5.5 (ii).

Let $\tilde{M}_t = M_t + \varepsilon$ and introduce the stopping time

$$\tau = \inf\{t : |N_t| \geq 1 - \varepsilon \text{ or } \tilde{M} \geq R\}.$$

Step 3. Applying Itô's formula to U^m , we obtain

$$U^m(\tilde{M}_{\tau \wedge t}, N_{\tau \wedge t}) = U^m(\tilde{M}_0, N_0) + I_1 + I_2 + I_3,$$

where

$$\begin{aligned} I_1 &= \sum_{0 < s < \tau \wedge t} [U^m(\tilde{M}_s, N_s) - U^m(\tilde{M}_{s-}, N_{s-}) \\ &\quad - U_x^m(\tilde{M}_{s-}, N_{s-}) \Delta M_s - U_y^m(\tilde{M}_{s-}, N_{s-}) \Delta N_s], \\ I_2 &= \int_0^{\tau \wedge t} U_{xx}^m(\tilde{M}_{s-}, N_{s-}) d[M^c, M^c]_s + 2U_{xy}^m(\tilde{M}_{s-}, N_{s-}) d[M^c, N^c]_s \\ &\quad + U_{yy}^m(\tilde{M}_{s-}, N_{s-}) d[N^c, N^c]_s. \end{aligned}$$

$$I_3 = \int_0^{\tau \wedge t} U_x^m(\tilde{M}_{s-}, N_{s-}) dM_s + \int_0^{\tau \wedge t} U_y^m(\tilde{M}_{s-}, N_{s-}) dN_s,$$

We have $I_1 \leq 0$ in view of (6.2). Moreover, $I_2 = J_1 + J_2 + J_3$, where

$$\begin{aligned} J_1 &= \int_0^{\tau \wedge t} U_{xx}^m(\tilde{M}_{s-}, N_{s-}) + 2U_{xy}^m(\tilde{M}_{s-}, N_{s-}) + U_{yy}^m(\tilde{M}_{s-}, N_{s-}) d[X, X]_s, \\ J_2 &= \int_0^{\tau \wedge t} U_{xx}^m(\tilde{M}_{s-}, N_{s-}) - 2U_{xy}^m(\tilde{M}_{s-}, N_{s-}) + U_{yy}^m(\tilde{M}_{s-}, N_{s-}) d[Y, Y]_s, \\ J_3 &= \int_0^{\tau \wedge t} U_{xx}^m(\tilde{M}_{s-}, N_{s-}) - U_{yy}^m(\tilde{M}_{s-}, N_{s-}) d[X, Y]_s \end{aligned}$$

and

$$X = \frac{M^c + N^c}{2}, \quad Y = \frac{M^c - N^c}{2}.$$

That J_1 and J_2 are nonpositive, follows from (6.1). The inequality $J_3 \leq 0$ is a consequence of (6.3) and the fact that the process $[X, Y] = \frac{1}{4}([M^c, M^c] - [N^c, N^c])$ is nonnegative and nondecreasing. Hence $I_2 \leq 0$. Finally, the expression in I_3 is a local martingale, which, by the stopping time argument (exactly as in the proof of the inequality (1.7)) can be assumed to satisfy $\mathbb{E}I_3 = 0$.

Summarizing, we have proved that

$$(6.5) \quad \mathbb{E}U^m(\tilde{M}_{\tau \wedge t}, N_{\tau \wedge t}) \leq \mathbb{E}U^m(\tilde{M}_0, N_0).$$

Step 4. This is the final part. Note that we have the estimate

$$(6.6) \quad 1 \geq U^m(x, y) \geq \inf_{(x', y') \in B_m} U(x - x', y - y') \geq -\frac{p^p}{2^p(p-1)}(x + \varepsilon)^p,$$

The martingales $(\tilde{M}_{\tau \wedge t}, (N_{\tau \wedge t}))$ converge almost surely, so by Lebesgue's dominated convergence theorem, if we let $t \rightarrow \infty$ in (6.5), we get

$$\mathbb{E}U^m(\tilde{M}_\tau, N_\tau) \leq \mathbb{E}U^m(\tilde{M}_0, N_0).$$

We have $U^m \rightarrow U$ pointwise as $m \rightarrow \infty$ and by (6.6) we may again use Lebesgue's theorem to obtain

$$\mathbb{E}U(\tilde{M}_0, N_0) \geq \mathbb{E}U(\tilde{M}_\tau, N_\tau) = \mathbb{E}U(\tilde{M}_\tau, N_\tau)1_{\{\tilde{M}_\tau < R\}} + \mathbb{E}U(\tilde{M}_\tau, N_\tau)1_{\{\tilde{M}_\tau \geq R\}}$$

Now $\mathbb{E}U(\tilde{M}_0, N_0) \leq 0$ by Lemma 5.5 (i). For the first expression appearing on the right, we use the inequality (6.4). For the second one, we use the bound

$$U(x, y) \geq -\frac{p^p}{2^p(p-1)}x^p,$$

which is a trivial consequence of Lemma 5.5 (ii). We arrive at

$$0 \geq \eta \mathbb{P}(|N_\tau| \geq 1 - \varepsilon, \tilde{M}_\tau < R) - \frac{p^p}{2^p(p-1)} \mathbb{E}\tilde{M}_\tau^p.$$

Therefore,

$$\begin{aligned}
\eta\mathbb{P}(N^* \geq 1) &\leq \eta\mathbb{P}(N^* \geq 1, \tilde{M}_\tau < R) + \eta\mathbb{P}(\tilde{M}_\tau \geq R) \\
&\leq \eta\mathbb{P}(N_\tau \geq 1 - \varepsilon, \tilde{M}_\tau < R) + \eta\mathbb{P}(\tilde{M}_\tau \geq R) \\
&\leq \frac{p^p}{2^p(p-1)}\mathbb{E}\tilde{M}_\tau^p + \frac{\eta}{R^p}\mathbb{E}\tilde{M}_\tau^p \leq \left(\frac{p^p}{2^p(p-1)} + \frac{\eta}{R^p}\right)\|\tilde{M}\|_p^p \\
&\leq \left(\frac{p^p}{2^p(p-1)} + \frac{\eta}{R^p}\right)(\varepsilon + \|M\|_p)^p.
\end{aligned}$$

Here, in the third passage, we used Chebyshev's inequality. Now let $\varepsilon \rightarrow 0$ to obtain

$$\eta\mathbb{P}(N^* \geq 1) \leq \frac{p^p}{2^p(p-1)}\|M\|_p^p.$$

As η was arbitrary, the proof is complete.

7. THE PROOF OF THEOREM 1.6

Let us begin with the following inequality.

Lemma 7.1. *Let $1 \leq p \leq 2$ and $x, h \in \mathbb{R}$. Then*

$$(7.1) \quad |x+h|^p + |x-h|^p \leq 2|x|^p + 2|h|^p.$$

Proof. Consider a map $T : \mathbb{R}^2 \rightarrow L_\infty(\Omega, \mathcal{F}, \mathbb{P})$ given by

$$T(x, h) = x + \varepsilon h,$$

where ε is a random variable such that $\mathbb{P}(\varepsilon = -1) = \mathbb{P}(\varepsilon = 1) = 1/2$. It follows from the triangle inequality that $\|T\|_{\ell_1^2 \rightarrow L_1} = 1$, and from the parallelogram identity, that $\|T\|_{\ell_2^2 \rightarrow L_2} = 1$. Thus, by interpolation, we have $\|T\|_{\ell_p^2 \rightarrow L_p} \leq 1$ for $1 \leq p \leq 2$. This yields (7.1). \square

Let $(B, \mathbb{P}) = ((B^1, B^2), (\mathbb{P}_{x,y})_{(x,y) \in \mathbb{R}^2})$ denote the family of 2-dimensional Brownian Motions such that for any (x, y) ,

$$\mathbb{P}_{x,y}(B_0 = (x, y)) = 1.$$

Recall the constant K_p given by (1.10) and define $V : \mathbb{R}^2 \rightarrow \mathbb{R}$ by $V(x, y) = 1_{\{|y| \geq 1\}} - K_p^p|x|^p$. Throughout this section, τ will denote the stopping time

$$\tau = \inf\{t \geq 0 : |B_t^2| \geq 1\}.$$

The special function is defined by

$$U(x, y) = \mathbb{E}_{x,y}|B_\tau^1|^p.$$

Note that in particular we have $U(x, y) = |x|^p$ for $|y| \geq 1$. It follows from the very definition that U is harmonic on S and continuous on \mathbb{R}^2 . Moreover, it is obvious that for any x, y we have

$$(7.2) \quad U(x, y) = U(-x, y) = U(x, -y) = U(-x, -y).$$

We study the further properties of the function U in the lemmas below. First we will provide an explicit formula for U . To do this, let $H = \{(\alpha, \beta) : \beta > 0\}$ denote the upper halfplane and define $\mathcal{W} : H \rightarrow \mathbb{R}$ as the Poisson integral

$$\mathcal{W}(\alpha, \beta) = \frac{2}{\pi^2} \int_{-\infty}^{\infty} \frac{\beta |\log |t||^p}{(\alpha - t)^2 + \beta^2} dt.$$

Clearly, \mathcal{W} is harmonic on H . Furthermore, we have

$$(7.3) \quad \lim_{(\alpha, \beta) \rightarrow (t, 0)} \mathcal{W}(\alpha, \beta) = \frac{2}{\pi} |\log |t||^p.$$

Consider the conformal map ϕ on $S = \{(x, y) : |y| < 1\}$, defined by

$$\phi(x, y) = \phi(z) = ie^{\pi z/2}.$$

It is easy to check that ϕ maps S onto H .

Lemma 7.2. (i) *We have the identity*

$$(7.4) \quad U(x, y) = \begin{cases} |x| & \text{if } |y| \geq 1, \\ \mathcal{W}(\phi(x, y)) & \text{if } |y| < 1. \end{cases}$$

(ii) *We have $U(0, 0) = K_p^{-p}$.*

Proof. (i) Note that both sides of (7.4) are harmonic on S , continuous on \mathbb{R}^2 (for the right-hand side, use (7.3)) and equal on the set $\{(x, y) : |y| \geq 1\}$. This proves the claim.

(ii) Using the first part of the lemma, we may write

$$\begin{aligned} U(0, 0) &= \mathcal{W}(0, 1) = \frac{4}{\pi^2} \int_0^\infty \frac{|\log t|^p}{t^2 + 1} dt \\ &= \frac{4}{\pi^2} \int_{-\infty}^\infty \frac{|s|^p e^s}{e^{2s} + 1} ds \\ &= \frac{8}{\pi^2} \int_0^\infty s^p e^{-s} \sum_{k=0}^\infty (-e^{-2s})^k ds \\ &= \frac{8}{\pi^2} \Gamma(p+1) \sum_{k=0}^\infty \frac{(-1)^k}{(2k+1)^{p+1}} = K_p^{-p}. \end{aligned}$$

Here we have used the identity

$$\sum_{k=0}^\infty \frac{1}{(2k+1)^2} = \frac{\pi^2}{8}. \quad \square$$

Lemma 7.3. *For fixed $x \in \mathbb{R}$, the function $U(x, \cdot)$ is concave on $(-1, 1)$.*

Proof. Since U is harmonic on S , an equivalent formulation is that for any fixed $y \in (-1, 1)$ the function $U(\cdot, y)$ is convex. For any $x_1, x_2 \in \mathbb{R}$ we have

$$\mathbb{E}_{x_1, y} |B_\tau^1|^p = \mathbb{E}_{x_2, y} |x_1 - x_2 + B_\tau^1|^p,$$

so if $\lambda \in (0, 1)$, then, by Jensen's inequality,

$$\begin{aligned} \lambda U(x_1, y) + (1 - \lambda)U(x_2, y) &= \mathbb{E}_{x_2, y} [\lambda |x_1 - x_2 + B_\tau^1|^p + (1 - \lambda)|B_\tau^1|^p] \\ &\geq \mathbb{E}_{x_2, y} |\lambda x_1 - \lambda x_2 + B_\tau^1|^p \\ &= \mathbb{E}_{\lambda x_1 + (1 - \lambda)x_2, y} |B_\tau^1|^p = U(\lambda x_1 + (1 - \lambda)x_2, y). \quad \square \end{aligned}$$

Lemma 7.4. (i) *For any $x > 0$, $y \in (0, 1)$, $U_{xyy} \geq 0$.*

(ii) *$U_{xy} \geq 0$ on S .*

Proof. (i) Since U_x is harmonic on S , the equivalent statement is $U_{xxx} \leq 0$. Fix x, y as in the statement and let $\varepsilon \in (0, x)$. We have

$$2U_x(x, y) - U_x(x - \varepsilon, y) - U_x(x + \varepsilon, y) = \mathbb{E}_{0, y} f(B_\tau^1),$$

where

$$f(h) = 2|x + h|^{p-2}(x + h) - |x - \varepsilon + h|^{p-2}(x - \varepsilon + h) - |x + \varepsilon + h|^{p-2}(x + \varepsilon + h).$$

Note that $f \geq 0$ for $h \geq -x$ and $f(-x + h) = -f(-x - h)$ for any h . As τ is independent of B^1 , we infer that the density of B_τ^1 under $\mathbb{P}_{0, y}$ is decreasing on $[0, \infty)$. This implies $\mathbb{E}_{0, y} f(B_\tau^1) \geq 0$ and, since $x > 0$ and $\varepsilon \in (0, x)$ were arbitrary, we conclude that U_x is concave.

(ii) We have $U_y(x, 0) = 0$ due to (7.2). Thus $U_{xy}(x, 0) = 0$ and the claim follows from the first part of the lemma. \square

Lemma 7.5. (i) We have $U(x, y) \geq U(0, 0)$ for any x, y such that $|y| \leq |x|$.

(ii) We have $U(x, y) \leq |x|^p + K_p^{-p} 1_{\{|y| < 1\}}$ for any $x, y \in \mathbb{R}$.

Proof. (i) For fixed x , the function $U(x, \cdot)$ is even, concave on $(-1, 1)$ and constant on $(-\infty, 1] \cup [1, \infty)$. This implies that it suffices to show the inequality for $x = y$. Let F be the function given by $F(x) = U(x, x)$, $x \in \mathbb{R}$. By (7.2), F is even. Moreover, by harmonicity of U and Lemma 7.4 (ii), we have

$$F''(x) = U_{xx}(x, x) + 2U_{xy}(x, x) + U_{yy}(x, x) \geq 0 \quad \text{for } x \in (0, 1),$$

from which we conclude that $F(x) \geq F(0)$ for $x \in (0, 1)$. Now note that F is continuous on \mathbb{R} and increasing on $(1, \infty)$. This finishes the proof.

(ii) It suffices to prove the inequality on S . The function

$$y \mapsto U(x, y) - |x|^p - K_p^{-p} 1_{\{|y| < 1\}} = U(x, y) - |x|^p - K_p^{-p}$$

is even (by (7.2)) and concave (due to Lemma 7.3), so attains its maximum at 0. Hence we must show that

$$U(x, 0) - |x|^p - U(0, 0) \leq 0,$$

or

$$\mathbb{E}_{0, 0} |x + B_\tau^1|^p \leq |x|^p + \mathbb{E}_{0, 0} |B_\tau^1|^p,$$

an inequality which follows from (7.1). \square

Proof of (1.6): We may assume $\|M\|_p < \infty$. If N is differentially subordinate and orthogonal to M , then N has continuous paths (see Lemma 2.1 in [3]). We proceed as in the proof of Theorem 1.5. Introduce the stopping time

$$\sigma = \inf\{t : |N_t| \geq 1\}.$$

Itô's formula gives

$$U(M_{\sigma \wedge t}, N_{\sigma \wedge t}) = U(M_0, N_0) + I_1 + I_2 + I_3,$$

where

$$\begin{aligned} I_1 &= \sum_{0 < s < \sigma \wedge t} [U(M_s, N_s) - U(M_{s-}, N_{s-}) - U_x(M_{s-}, N_{s-}) \Delta M_s], \\ I_2 &= \int_0^{\sigma \wedge t} U_{xx}(M_{s-}, N_{s-}) d[M^c, M^c]_s + 2U_{xy}(M_{s-}, N_{s-}) d[M^c, N^c]_s \\ &\quad + U_{yy}(M_{s-}, N_{s-}) d[N^c, N^c]_s. \end{aligned}$$

$$I_3 = \int_0^{\sigma \wedge t} U_x(M_{s-}, N_{s-}) dM_s + \int_0^{\sigma \wedge t} U_y(M_{s-}, N_{s-}) dN_s,$$

Note that $I_1 \geq 0$ by Lemma 7.3 and $I_2 = 0$ as U is harmonic in S and M, N are orthogonal. Furthermore, using the stopping time argument, we may assume $\mathbb{E}I_3 = 0$ and obtain

$$\mathbb{E}U(M_{\sigma \wedge t}, N_{\sigma \wedge t}) \geq \mathbb{E}U(M_0, N_0).$$

This, combined with Lemma 7.2 (ii) and Lemma 7.5, leads to

$$\mathbb{E}|M_{\sigma \wedge t}|^p + K_p^{-p} \mathbb{P}(|N_{\sigma \wedge t}| < 1) \geq K_p^{-p},$$

or

$$\mathbb{P}(|N_{\sigma \wedge t}| \geq 1) \leq K_p^p \|M\|_p^p.$$

Now the proof is completed using the scaled martingales $M^\varepsilon, N^\varepsilon$, exactly in the same manner as in the proof the inequality (1.7). \square

Sharpness. Let $M = (B_{\tau \wedge t}^1)$, $N = (B_{\tau \wedge t}^2)$. Then N is differentially subordinate and orthogonal to M . Moreover,

$$\mathbb{P}(N^* \geq 1) - K_p^p \|M\|_p = 1 - K_p^p U(0, 0) = 0,$$

which implies that the best constant in (1.6) is not smaller than K_p . \square

8. INEQUALITIES FOR HARMONIC FUNCTIONS

In this section we study related weak type inequalities for harmonic functions on Euclidean domains. Let n be a positive integer and let D be an open connected subset of \mathbb{R}^n . Fix a point ξ , which belongs to D . For two real-valued harmonic functions u, v on D , we say that v is *differentially subordinate* to u , if the following two conditions are satisfied:

$$(8.1) \quad |v(\xi)| \leq |u(\xi)|$$

and

$$(8.2) \quad |\nabla v(x)| \leq |\nabla u(x)| \quad \text{for any } x \in D.$$

This concept was introduced by Burkholder; see [5] for more information and references. We say that u, v are orthogonal, if

$$(8.3) \quad \nabla v(x) \cdot \nabla u(x) = 0 \quad \text{for any } x \in D,$$

where \cdot denotes the scalar product in \mathbb{R}^n .

Let D_0 be a bounded domain satisfying $\xi \in D_0 \subset D_0 \cup \partial D_0 \subset D$. Let $\mu_{D_0}^\xi$ denote the harmonic measure on ∂D_0 , corresponding to ξ . Consider weak and strong p -norms given by

$$\|u\|_{p, \infty} = \sup \sup_{\lambda} [\lambda^p \mu_{D_0}^\xi(\{x \in \partial D_0 : |u(x)| \geq \lambda\})]^{1/p},$$

$$\|u\|_p = \sup \left[\int_{\partial D_0} |u(x)|^p d\mu_{D_0}^\xi(x) \right]^{1/p},$$

the first supremums being taken over all subdomains D_0 as above.

Now we state the harmonic analogue of Theorems 1.4 and 1.5.

Theorem 8.1. *Suppose v is differentially subordinate to u and u is nonnegative.*

(i) *For $0 < p < 1$ we have*

$$\|v\|_{p,\infty} \leq 2\|u\|_p$$

and the inequality is sharp.

(ii) *For $p \geq 2$ we have*

$$\|v\|_{p,\infty} \leq \frac{p}{2}(p-1)^{-1/p}\|u\|_p.$$

We do not know if the constant in (ii) are the best possible (except for the case $p = 2$, when, clearly, it is).

Proof. It suffices to show that for any bounded domain D_0 as above we have

$$(8.4) \quad \mu_{D_0}^\xi(\{x \in \partial D_0 : |v(x)| \geq 1\}) \leq \frac{p^p}{2^p(p-1)} \int_{\partial D_0} u(x)^p d\mu_{D_0}^\xi(x).$$

Let $B = (B_t)_{t \geq 0}$ be a Brownian Motion in \mathbb{R}^n , starting from ξ and introduce a stopping time

$$\tau = \tau_{D_0} = \inf\{t : B_t \in \partial D_0\}.$$

Let M, N be martingales defined by

$$(8.5) \quad M_t = u(B_{\tau \wedge t}) \quad \text{and} \quad N_t = v(B_{\tau \wedge t}), \quad t \geq 0.$$

Note that M is nonnegative and N is real-valued. The property (8.2) gives $|N_0| \leq |M_0|$ and, combined with (8.1), implies that N is differentially subordinate to M : this follows from identities

$$\begin{aligned} [M, M]_t &= |M_0|^2 + \int_0^{\tau \wedge t} |\nabla u(B_s)|^2 ds, \\ [N, N]_t &= |N_0|^2 + \int_0^{\tau \wedge t} |\nabla v(B_s)|^2 ds. \end{aligned}$$

Hence

$$\mu_{D_0}^\xi(\{x \in \partial D_0 : |v(x)| \geq 1\}) \leq \mathbb{P}(N^* \geq 1) \leq \frac{p}{2(p-1)^{1/p}} \|M_\tau\|_p = \frac{p}{2(p-1)^{1/p}} \|u\|_p.$$

Now we show the constant 2 is optimal in (i). Here we use the example of Burkholder [7]. Let $n = 1$, $D = (-1, 3)$, $\xi = 0$, $u(x) = 1 + x$ and $v(x) = 1 - x$. Then u, v are harmonic, u is nonnegative and v is differentially subordinate to u . We have $\|u\|_p = u(\xi) = 1$. Furthermore, for $0 < \lambda < 2$ we have $|v(x)| < \lambda$ if and only if $x \in (1 - \lambda, 1 + \lambda)$, which implies

$$\limsup_{\lambda \uparrow 2} \lambda (\mu_{D_0}(|v| \geq \lambda))^{1/p} = \lim_{\lambda \uparrow 2} \lambda = 2\|u\|_p.$$

Therefore we cannot replace 2 in (i) by a smaller number. \square

The harmonic extension of Theorem 1.6 can be stated as follows.

Theorem 8.2. *Suppose v is differentially subordinate to u and u, v are orthogonal. Then for $1 \leq p \leq 2$ we have*

$$\|v\|_{p,\infty} \leq K_p \|u\|_p.$$

The inequality is sharp.

Proof. The inequality is proved by the same argumentation as above, using the martingales M and N given by (8.5). Their orthogonality is guaranteed by

$$[M, N]_t = M_0 N_0 + \int_0^{\tau \wedge t} \nabla u(B_s) \cdot \nabla v(B_s) ds.$$

We omit the details.

To see that the inequality is sharp, let $n = 2$, $\varepsilon > 0$, $D = \mathbb{R} \times (-1 - \varepsilon, 1 + \varepsilon)$, $\xi = (0, 0)$, $u(x, y) = x$, $v(x, y) = y$. Clearly, u, v are harmonic, orthogonal and v is differentially subordinate to u . For $D_0 = (-R, R) \times (-1, 1)$ and $\mu = \mu_{D_0}^\xi$, we have

$$\lim_{R \rightarrow \infty} \left[\int_{\partial D_0} |u(x)|^p d\mu(x) \right]^{1/p} = \|B_\tau^1\|_p = \frac{1}{K_p},$$

which gives

$$\lim_{\varepsilon \downarrow 0} \|u\|_p = \frac{1}{K_p}.$$

To complete the proof, it suffices to note that

$$\lim_{R \rightarrow \infty} \mu(|v| \geq 1) = 1 \quad \text{and hence} \quad \|v\|_{p, \infty} \geq 1,$$

which implies that K_p is the best possible. \square

REFERENCES

- [1] M. Abramowitz and I. A. Stegun, editors, *Handbook of Mathematical Functions with formulas, graphs and mathematical tables*, Reprint of the 1972 edition, Dover Publications, Inc., New York, 1992.
- [2] R. Bañuelos and G. Wang, *Sharp inequalities for martingales with applications to the Beurling-Ahlfors and Riesz transformations*, Duke Math. J. 80 (1995), 575–600.
- [3] R. Bañuelos and G. Wang, *Davis’s inequality for orthogonal martingales under differential subordination*, Michigan Math. J. 47 (2000), 109–124.
- [4] D. L. Burkholder, *Boundary value problems and sharp inequalities for martingale transforms*, Ann. Probab. 12 (1984), 647–702.
- [5] D. L. Burkholder, *Differential subordination of harmonic functions and martingales*, Harmonic Analysis and Partial Differential Equations (El Escorial, 1987), Lecture Notes in Mathematics 1384 (1989), 1–23.
- [6] D. L. Burkholder, *Explorations in martingale theory and its applications*, École d’Ete de Probabilités de Saint-Flour XIX—1989, pp. 1–66, Lecture Notes in Math., 1464, Springer, Berlin, 1991.
- [7] D. L. Burkholder, *Strong differential subordination and stochastic integration*, Ann. Probab. 22 (1994), pp. 995–1025.
- [8] D. L. Burkholder, *Some extremal problems in martingale theory and harmonic analysis*, Harmonic analysis and partial differential equations (Chicago, IL, 1996), 99–115, Chicago Lectures in Math., Univ. Chicago Press, Chicago, IL, 1999.
- [9] C. Choi, *A weak-type inequality for differentially subordinate harmonic functions*, Trans. Amer. Math. Soc. 350, pp. 2687–2696.
- [10] C. Dellacherie and P. A. Meyer, *Probabilities and potential B*, North-Holland, Amsterdam, 1982.
- [11] R Development Core Team, *R: A Language and Environment for Statistical Computing*, R Foundation for Statistical Computing, Vienna, Austria (2008).
- [12] Y. Suh, *A sharp weak type (p, p) inequality $(p > 2)$ for martingale transforms and other subordinate martingales*, Trans. Amer. Math. Soc. 357 (2005), no. 4, 1545–1564 (electronic).
- [13] G. Wang, *Differential subordination and strong differential subordination for continuous time martingales and related sharp inequalities*, Ann. Probab. 23 (1995), 522–551.

DEPARTMENT OF MATHEMATICS, INFORMATICS AND MECHANICS, UNIVERSITY OF WARSAW, BANACHA 2, 02-097 WARSAW, POLAND

E-mail address: `ados@mimuw.edu.pl`