

Equivariant Hirzebruch class for quadratic cones via degenerations

Małgorzata Mikosz

Warsaw University of Technology,
ul. Koszykowa 75, 00-662, Warszawa, Poland
emmikosz@mini.pw.edu.pl

Andrzej Weber*

Department of Mathematics of Warsaw University
Banacha 2, 02-097 Warszawa, Poland
aweber@mimuw.edu.pl

August 2014

Let X be a smooth algebraic variety and Y a subvariety. The cohomology class of Y in $H^*(X)$, i.e. the Poincaré dual of the fundamental class of Y , does not change when we deform Y in a flat manner. A more subtle cohomological invariant of Y is the Hirzebruch class

$$td_y(Y \rightarrow X) \in H_*(X) \otimes \mathbb{Q}[y]$$

defined in [BSY]. A flat family member Y_t can be thought of as a fiber of a function

$$X \times \mathbb{C} \supset W \xrightarrow{\pi} \mathbb{C}.$$

The difference between the Hirzebruch class of the generic fiber and the Hirzebruch class of the special fiber is measured by the appropriate version of Milnor class, studied in [CMSS] for hypersurfaces and in [MSS] the general case. The same phenomenon happens for the equivariant Hirzebruch class developed in [We3], compare also with [Oh, Sec.4] for the equivariant Hirzebruch class in the context of quotient stacks. We fix our attention on the varieties with torus action. If we are interested in local invariants of singularities, we study the localization of the equivariant Hirzebruch class $td_y^{\mathbb{T}}(Y \rightarrow X)$ at a fixed point. The bottom degree of the Hirzebruch class is the equivariant fundamental class, also called the multi-degree of the variety. It does not change in the deformation class. For example, let $\widehat{Q}_n \subset \mathbb{C}^n$ be the cone over a quadric in \mathbb{P}^{n-1} , in other words, \widehat{Q}_n in some coordinates is

*Supported by NCN grant 2013/08/A/ST1/00804

described by the Morse function $\sum_{i=1}^n x_i^2$. Let $\mathbb{T} = \mathbb{C}^*$ act on \mathbb{C}^n diagonally. Then $[\widehat{Q}_n]$ is equal to $2t$, with

$$t = c_1(\mathbb{C}) \in H_{\mathbb{T}}^*(pt) \simeq H_{\mathbb{T}}^*(\mathbb{C}^n) \simeq \mathbb{Q}[t],$$

the first Chern class of the standard weight one representation. Indeed \widehat{Q}_n can be equivariantly degenerated to the sum of two transverse hyperplane. The difference of the Hirzebruch classes is supported by the singular locus of the special member of the family. In the case of quadratic cones (\widehat{Q}_n and intersection of planes) both varieties have only rational singularities, therefore ([BSY, Example 3.2]) their Hirzebruch classes for $y = 0$ are equal to the Todd classes constructed by Baum-Fulton-MacPherson. The Todd class of a hypersurface H of an ambient manifold M are expressed by the class $[H]$ and the Todd class of M , precisely $i_*td(H) = td(M)(1 - e^{-[H]})$, where i is the inclusion $i : H \hookrightarrow M$, see eg. [Fu, Th. 18.3(4)]. One easily generalizes this formula in the equivariant setting. Hence the Todd classes of \widehat{Q}_n and \widehat{X}_n are equal. (Alternatively one can apply Verdier specialization argument, which implies that the Todd class of singular spaces is constant in flat families, [Ve].) It follows that full Milnor class is divisible by y .

We would like to present how the equivariant Hirzebruch class degenerates for the cone singularities. Our work started when we tried to analyze the equivariant Hirzebruch class of the cone. For the fixed dimension n it is easy to compute the corresponding polynomial. From initial sequence of coefficients it was hard to guess a closed formula and, for example, to prove a kind of positivity studied in [We3, §13]. Applying the degeneration method we find an answer. An interesting reciprocity happens. The difference between the Hirzebruch classes of the projective quadric Q_n and two intersecting projective hyperplanes X_n is the Hirzebruch class of the complement of another projective quadric multiplied by y :

$$td_y^{\mathbb{T}}(Q_n) - td_y^{\mathbb{T}}(X_n) = y \cdot td_y^{\mathbb{T}}(\mathbb{P}^{n-3} \setminus Q_{n-2}) \quad (1)$$

(Formula 3). In the non-equivariant context this result should follow for example from [CMSS, Thm.1.4, Rem.1.5] and the methods of [PaPr, Sec.5]. (as explained later in Remark 2). In this paper we even prove more directly a corresponding result for the equivariant Hirzebruch classes. Using induction we find the equivariant Hirzebruch classes of Q_n and \widehat{Q}_n .

Having in mind the expression for Chern-Schwartz-MacPherson class of smooth open varieties via logarithmic forms [Al], it is more natural to compute the Hirzebruch class of the complement $Q_n^* = \mathbb{C}^n \setminus \widehat{Q}_n$. For $n = 2m$ we obtain the expression

$$(1+y)^2 T^2 \sum_{i=1}^m (-y)^{m-i} \frac{(1+yT)^{2i-2}}{(1-T)^{2i}}$$

and for $n = 2m + 1$

$$(-y)^m \frac{(y+1)T}{1-T} + (1+y)^2 T^2 \sum_{i=1}^m (-y)^{m-i} \frac{(1+yT)^{2i-1}}{(1-T)^{2i+1}}.$$

Here $T = e^{-t}$ and the given expression is equal to the Hirzebruch class divided by the Euler class of $0 \in \mathbb{C}^n$, that is $eu(0) = t^n$. The formulas are understood as elements of the completed $H_{\mathbb{T}}^*(\mathbb{C}^n)[y]$ and localized in t . This ring is isomorphic to the ring of Laurent series in t and polynomials in y , i.e. $\mathbb{Q}[[t]][t^{-1}, y]$. (We will omit the completion in our notation for cohomology.) The formulas follow from Corollary 10 by the specialization T_i to one. Taking the limit $y \rightarrow -1$ with $T = e^{-(y+1)t}$ we obtain the expression for the Chern-Schwartz-MacPherson class of X_n^* in equivariant cohomology of \mathbb{C}^n : for $n = 2m$

$$\sum_{i=0}^{m-1} t^{2i} (1+t)^{2(m-i-1)}$$

and for $n = 2m + 1$

$$t^{2m} + \sum_{i=0}^{m-1} t^{2i} (1+t)^{2(m-i-1)}$$

which, as one can check, agrees with the invariant of a conical set introduced in [AlMa], compare [We1, §8]. We note that the quadratic cone appears as a singularity of Schubert varieties: the quadric Q_n can be considered as a homogenous space with respect to $SO(n)$ and the codimension one Schubert variety is isomorphic to the projective cone over Q_{n-2} . It would be interesting to examine singularities of Schubert varieties from the point of view of degenerations, having in mind the work on smoothability [Co1, Co2] and intersection theory [CoVa].

The presented computation in fact is a baby example of what can happen. The aim of the paper is to show a bunch of computation of the Hirzebruch class based on Localization Theorem 4. The **Formulas 3, 8** and **12** are the outcome. They show how Milnor class may be realized geometrically. We hope that these formulas will find generalizations for some class of degenerations of Schubert varieties.

We would like to thank the Referee for very careful reading of the manuscript. He has suggested many important improvements. The meaningful Remark 2 is due to him.

1 Hirzebruch classes of projective quadrics

To understand systematically the situation we consider a bigger torus preserving the quadric. One has to distinguish between the cases of even and

odd n . Let us index the coordinates in \mathbb{C}^{2m} by integer numbers from $-m$ to m omitting 0 and consider the quadratic form in \mathbb{C}^{2m} given by the formula

$$\sum_{i=1}^m x_{-i}x_i.$$

For \mathbb{C}^{2m+1} allow the index 0 and fix the quadratic form

$$x_0^2 + \sum_{i=1}^m x_{-i}x_i.$$

Let $Q_n \subset \mathbb{P}^{n-1}$ be the quadric defined by vanishing of the quadratic form. It is an invariant variety with respect to the torus $\mathbb{T}_m = (\mathbb{C}^*)^m$ action coming from the representation with weights (i.e. characters)

$$(-t_m, -t_{-m+1}, \dots, t_{m-1}, t_m)$$

if $n = 2m$ and

$$(-t_m, -t_{-m+1}, \dots, 0, \dots, t_{m-1}, t_m)$$

for $n = 2m + 1$. Consider the equivariant Hirzebruch class

$$td_y^{\mathbb{T}_m}(Q_n \rightarrow \mathbb{P}^{n-1}) \in H_{\mathbb{T}_m}^*(\mathbb{P}^{n-1})[y]$$

and compare it with the Hirzebruch class of degeneration X_n of Q_n given by the equation $x_{-m}x_m = 0$. The variety X_n is the sum of the two coordinate planes. We think of X_n as the special fiber for $\lambda = 0$ of the equivariant family given by the equation

$$\lambda \sum_{i=1}^{m-1} x_{-i}x_i + x_{-m}x_m \quad \text{or} \quad \lambda \left(x_0^2 + \sum_{i=1}^{m-1} x_{-i}x_i \right) + x_{-m}x_m.$$

We will show that the difference of the Hirzebruch classes is the Hirzebruch class of $\mathbb{C}^{n-2} \setminus Q_{n-2}$ multiplied by y , i.e. Formula 3.

Remark 2 Let us explain why the formula (1) holds in non-equivariant cohomology¹. In $H^*(\mathbb{P}^{n-1})$

$$td_y(Q_n) - td_y(X_n) = y \cdot td_y(\mathbb{P}^{n-3} \setminus Q_{n-2})$$

should follow from results and techniques a la [CMSS, Thm.1.4, Rem.1.5] and [PaPr, Sec.5]:

$$g = \sum_{i=1}^{m-1} x_{-i}x_i \quad \text{and} \quad f = x_{-m}x_m$$

¹This remark is due to the Referee

are both sections of the line bundle $\mathcal{O}(2)$ on \mathbb{P}^{n-1} , with $Z' := \{g = 0\}$ and $Z := X_n = \{f = 0\}$ transversal in a stratified sense. Let

$$p : \mathcal{Z} := \{\lambda g + f = 0\} \subset \mathbb{P}^{n-1} \times \mathbb{C} \rightarrow \mathbb{C}$$

be the projection onto the last variable λ . Then the vanishing cycles $\phi_p(\mathbb{Q}_{\mathcal{Z}})$ are supported by the critical locus $\mathbb{P}^{n-3} = \{x_{-m} = 0 = x_m\} \subset X_n = \{p = 0\}$ of p . Moreover, the restriction of these vanishing cycles to $Z \cap Z' = Q_{n-3} \subset \mathbb{P}^{n-3}$ should be zero by the argument of [PaPr, Sec.5] (or [MSS, part a) of the proof of Prop. 4.1]). Moreover, the corresponding nearby cycles can be calculated in terms of the generic fiber $Q_n = \{p = 1\}$, since p is quasi-homogeneous (i.e. equivariant for a suitable \mathbb{C}^* -action). Then the stated formula above follows from [CMSS, Thm.1.4, Rem.1.5], with the factor y equal to the (reduced) χ_y -genus of the transversal Milnor fiber of an A_1 -singularity $z^2 + w^2 = 0 = x_{-m}x_m$ in \mathbb{C}^2 . This remark would be an alternative proof of our formula provided that one developed the general theory of Milnor class in the equivariant case.

It is more convenient to work with complements of the closed varieties from the beginning. We will give formulas for complements of the quadrics, since then the components have better geometric interpretation. To make the notation easier we identify the equivariant cohomology with respect to \mathbb{T}_m with the subspace of $H_{\mathbb{T}_{m+1}}^*(\mathbb{P}^{n-1}) \simeq H_{\mathbb{T}_m}^*(\mathbb{P}^{n-1}) \otimes \mathbb{Q}[t_{m+1}]$ given by $t_{m+1} = 0$ and omit the index m in \mathbb{T}_m . Also we will omit the ambient space in the notation. This should not lead to a confusion; enlarging the ambient space results in introducing of the factor, which is the Euler class of the normal bundle. We will use this for example for the inclusions $\iota : \mathbb{P}^{n-3} \rightarrow \mathbb{P}^{n-1}$ into the first coordinates and the corresponding inclusions of the affine spaces. For an isolated fixed point $p \in Q_{n-2} \subset \mathbb{P}^{n-3} \subset \mathbb{P}^{n-1}$ the quotient $\frac{td_y^{\mathbb{T}}(Q_{n-2})|_p}{eu(p)}$ (where $eu(p) \in H_{\mathbb{T}}^*(pt)$ is the Euler class of the ambient tangent representation) does not depend on the ambient space. After these remarks about notation we state our first formula:

Formula 3 *Consider the complements of the quadrics $X'_n = \mathbb{P}^{n-1} \setminus X_n$ and $Q'_n = \mathbb{P}^{n-1} \setminus Q_n$. We have the equation*

$$td_y^{\mathbb{T}}(X'_n) - td_y^{\mathbb{T}}(Q'_n) = y td_y^{\mathbb{T}}(Q'_{n-2})$$

in the equivariant cohomology $H_{\mathbb{T}}^(\mathbb{P}^{n-1})[y]$ for $n > 2$. For the closed varieties we have*

$$td_y^{\mathbb{T}}(Q_n) - td_y^{\mathbb{T}}(X_n) = y td_y^{\mathbb{T}}(Q'_{n-2}).$$

2 Topological and analytic localization theorems

First let us note that equivariant cohomology is a homotopy invariant, for example for any \mathbb{T} -representation V the restriction map $H_{\mathbb{T}}^*(V) \rightarrow H_{\mathbb{T}}^*(\{0\})$

is an isomorphism. Therefore we get for free $H_{\mathbb{T}}^*(V) \xrightarrow{\cong} H_{\mathbb{T}}^*(V^{\mathbb{T}})$. We need much stronger property of equivariant cohomology. The main tool for computations is the Localization Theorem, see [Bo, Ch.XII §6] or [Qu]:

Theorem 4 (Topological Localization Theorem) [Qu, Theorem 4.4] *Assume either X is a compact topological space or that X is paracompact, $cd_{\mathbb{Q}}(X) < \infty$. Suppose a compact torus \mathbb{T} acts on X and the set of identity components of the isotropy groups of points of X is finite. Then the restriction map*

$$H_{\mathbb{T}}^*(X) \rightarrow H_{\mathbb{T}}^*(X^{\mathbb{T}})$$

is an isomorphism after localization in the multiplicative system generated by nontrivial characters.

We apply Topological Localization Theorem to algebraic varieties with algebraic torus action. The fixed points of the compact torus are the same as the fixed points of the full torus. The theorem may be applied to *any* algebraic variety, but it may very well happen (exactly when $X^{\mathbb{T}} = \emptyset$) that the localized equivariant cohomology is trivial.

For differential manifolds the isomorphism was made explicit by Atiyah-Bott and Berline-Vergne, see also [EdGr].

Theorem 5 (Topological Localization Theorem) [AtBo, page 9], [BeVe] *Let \mathbb{T} be a compact torus and let M be a compact \mathbb{T} -manifold. Let*

$$M^T = \bigsqcup_{\alpha \in I} F_{\alpha}$$

be the decomposition of the fixed point set into connected components. Denote by $\iota_{\alpha} : F_{\alpha} \rightarrow M$ the inclusion. Let

$$eu(F_{\alpha}) \in H_{\mathbb{T}}^*(F_{\alpha}) \simeq H^*(F_{\alpha}) \otimes H_{\mathbb{T}}^*(pt)$$

be the equivariant Euler class of the normal bundle to F_{α} . Let S be the multiplicative system generated by nontrivial characters. Then

1. *The class $eu(F_{\alpha})$ is invertible in $S^{-1}H_{\mathbb{T}}^*(F_{\alpha})$.*
2. *For any equivariant cohomology class $\omega \in H_{\mathbb{T}}^*(M)$ the equality holds*

$$\omega = \sum_{\alpha \in I} \iota_{\alpha*} \left(\frac{\iota_{\alpha}^*(\omega)}{eu(F_{\alpha})} \right) \tag{6}$$

in $S^{-1}H_{\mathbb{T}}^(M)$.*

The resulting integration formula follows, [AtBo, Formula 3.8].

The case of compact algebraic smooth varieties is special. The equivariant cohomology with respect to an algebraic torus action is always a free

module over $H_{\mathbb{T}}^*(pt)$ (see [GKM] and the references therein). Therefore the restriction map

$$H_{\mathbb{T}}^*(M) \rightarrow H_{\mathbb{T}}^*(M^T)$$

is a monomorphism. *The equality of the classes restricted to the fixed point set implies their equality.* We will use just this principle. Nevertheless, having in mind the formula (6), it is natural and convenient to consider the *localized Hirzebruch class*

$$\frac{\iota_{\alpha}^*(td_y^{\mathbb{T}}(-))}{eu(F_{\alpha})}$$

in the localized cohomology of fixed point set components. The spaces we consider here have only isolated fixed point sets, thus the localized Hirzebruch classes are polynomials in y with coefficients in the ring of Laurent polynomials in t_i 's. In fact the coefficients are rational functions in $T_i = e^{-t_i}$.

3 Properties of equivariant Hirzebruch class

Now we would like to recall basic properties of the equivariant Hirzebruch class, which in fact formally do not differ from the properties of the non-equivariant class. For an equivariant line bundle L the class $td_y^{\mathbb{T}}(L)$ is given in equivariant cohomology by the power series

$$t \frac{1 + y e^{-t}}{1 - e^{-t}},$$

with t the first equivariant Chern class of L . Then the corresponding class of a vector bundle is given in terms of Chern roots, and the class for a smooth manifold M is the corresponding class of the tangent bundle TM . In the localized classes of a smooth manifold appears then the (corrected) factor

$$\Phi(T) = \frac{1 + yT}{1 - T}$$

with $T = e^{-t}$ at the normal directions to the fixed point set.

The important properties of the equivariant Hirzebruch classes of singular varieties used in this paper are:

1. the normalization for smooth spaces (the Hirzebruch class is a series in equivariant Chern classes of tangent bundle),
2. covariant functoriality under proper maps,
3. additivity.

For example: Let $\pi : \widetilde{M} \rightarrow M$ be an equivariant proper morphism, with $\pi|_{\widetilde{M} \setminus E}$ an isomorphism on the image for some $E \subset M$, a closed invariant

subspace (for example the blowup of the origin in $M = \mathbb{C}^n$ with $E = \mathbb{P}^{n-1}$ the exceptional divisor, as used later on). Then

$$\pi_*(td_y^{\mathbb{T}}(\widetilde{M}) - td_y^{\mathbb{T}}(E)) = td_y^{\mathbb{T}}(M) - td_y^{\mathbb{T}}(\pi(E)).$$

As an example for additivity (or the inclusion-exclusion principle) one can calculate:

$$td_y^{\mathbb{T}}(X_n) = td_y^{\mathbb{T}}(\{x_m = 0\}) + td_y^{\mathbb{T}}(\{x_{-m} = 0\}) - td_y^{\mathbb{T}}(\{x_{-m} = x_m = 0\});$$

since $X_n = \{x_m = 0\} \cup \{x_{-m} = 0\}$ but the intersection is counted twice. In particular one can calculate in this simple way the class of the singular space X_n in terms of classes of smooth spaces.

The next property follows from 1.-3.:

4. multiplicativity and, more generally, contravariant functoriality with respect to fibrations.

For example if $p : \nu \rightarrow X$ is an equivariant vector bundle, then the Hirzebruch class of the total space of ν is equal to

$$td_y^{\mathbb{T}}(Tot(\nu)) = p^* \left(td_y^{\mathbb{T}}(\nu) \cdot td_y^{\mathbb{T}}(X) \right). \quad (7)$$

Here $td_y^{\mathbb{T}}(\nu)$ is understood as a characteristic class of a vector bundle.

4 Proof of Formula 3.

By Localization Theorem 4 it is enough to check equality at each fixed point of \mathbb{T} -action. The fixed points p_i corresponds to the coordinate lines in \mathbb{C}^n . Let us show the calculation for even $n = 2m$. At the point p_i the quadric is given by the equation

$$u_{-i} + \sum_{j \neq i} u_{-j} u_j = 0$$

in coordinates $u_j = x_j/x_i$. For a fixed point p_i the Hirzebruch class $td_y^{\mathbb{T}}(Q_n)$ divided by Euler class of at p_i (i.e. the localized Hirzebruch class) is equal to the product

$$\frac{1}{eu(p_i)} td_y^{\mathbb{T}}(Q_n) = \prod_{\text{weights of } T_{p_i} Q_n} \Phi(e^{-w}).$$

Here the product is taken with respect to the weights appearing in the tangent representation $T_{p_i} Q_n$ (see [We3, §1]). Let us set $t_{-i} = -t_i$ and $T_i = e^{-t_i}$. The weights of the tangent representation $T_{p_i} \mathbb{P}^{n-1}$ are equal to $t_j - t_i$ for $j \neq i$. The normal direction has weight $t_{-i} - t_i = -2t_i$. Since Q'_{2m} is the complement of Q_{2m} in \mathbb{P}^{n-1} , one gets by additivity that

$$\frac{1}{eu(p_i)} td_y^{\mathbb{T}}(Q'_{2m})_{p_i} = \frac{1}{eu(p_i)} td_y^{\mathbb{T}}(\mathbb{P}^{n-1})_{p_i} - \frac{1}{eu(p_i)} td_y^{\mathbb{T}}(Q_{2m})_{p_i}.$$

- At each point p_i , $|i| \leq m$ the localized Hirzebruch class is equal to

$$(\Phi(T_i^{-2}) - 1) \cdot \prod_{j=1, j \neq i}^m \Phi(T_j T_i^{-1}) \Phi(T_j^{-1} T_i^{-1}).$$

The class $td_y^{\mathbb{T}}(X'_{2m})$ is equal to

$$td_y^{\mathbb{T}}(\mathbb{P}^{n-1}) - td_y^{\mathbb{T}}(\{x_m = 0\}) - td_y^{\mathbb{T}}(\{x_{-m} = 0\}) + td_y^{\mathbb{T}}(\{x_m = x_{-m} = 0\}).$$

Therefore the localized class $\frac{1}{eu(p_i)} td_y^{\mathbb{T}}(X'_{2m})|_{p_i}$ is the following

- at the points p_i , $|i| < m$

$$\Phi(T_i^{-2}) \cdot (\Phi(T_m T_i^{-1}) - 1) (\Phi(T_m^{-1} T_i^{-1}) - 1) \cdot \prod_{j=1, j \neq i}^{m-1} \Phi(T_j T_i^{-1}) \Phi(T_j^{-1} T_i^{-1})$$

since

$$\begin{aligned} & \Phi(T_m T_i^{-1}) \Phi(T_m^{-1} T_i^{-1}) - \Phi(T_m T_i^{-1}) - \Phi(T_m^{-1} T_i^{-1}) + 1 = \\ & = (\Phi(T_m T_i^{-1}) - 1) (\Phi(T_m^{-1} T_i^{-1}) - 1) \end{aligned}$$

- at the point p_i , $|i| = m$

$$(\Phi(T_i^{-2}) - 1) \cdot \prod_{j=1}^{m-1} \Phi(T_j T_i^{-1}) \Phi(T_j^{-1} T_i^{-1}).$$

For the points p_{-m} and p_m which do not belong to $\iota(Q_{n-2})$ the considered classes are equal. At the point p_i for $|i| < m$ the classes $td_y^{\mathbb{T}}(Q'_{2m})$, $y td_y^{\mathbb{T}}(Q'_{2m-2})$ and $td_y^{\mathbb{T}}(X'_{2m})$ have the common factor

$$\prod_{j=1, j \neq i}^{m-1} \Phi(T_j T_i^{-1}) \Phi(T_j^{-1} T_i^{-1})$$

and it is enough to check the equality

$$\begin{aligned} & \Phi(T_i^{-2}) \cdot (\Phi(T_m T_i^{-1}) - 1) \cdot (\Phi(T_m^{-1} T_i^{-1}) - 1) - \\ & - (\Phi(T_i^{-2}) - 1) \cdot \Phi(T_m T_i^{-1}) \cdot \Phi(T_m^{-1} T_i^{-1}) = y(\Phi(T_i^{-2}) - 1). \end{aligned}$$

After multiplying by

$$(1 - T_i^{-2}) \cdot (1 - T_i^{-1} T_m) \cdot (1 - T_i^{-1} T_m^{-1})$$

the equality reduces to

$$\begin{aligned} (1 + yT_i^{-2}) \cdot (y + 1)(T_m T_i^{-1}) \cdot (y + 1)(T_m^{-1} T_i^{-1}) - \\ -(y + 1)(T_i^{-2}) \cdot (1 + yT_m T_i^{-1}) \cdot (1 + yT_m^{-1} T_i^{-1}) = \\ = y(y + 1)(T_i^{-2}) \cdot (1 - T_m T_i^{-1}) \cdot (1 - T_m^{-1} T_i^{-1}), \end{aligned}$$

which one verifies easily. The proof for n odd is identical except that all the expressions are multiplied by $\Phi(T_i^{\pm 1})$. \square

Also for $n = 2$ if we admit that $Q_0 = \mathbb{P}^{-1} = \emptyset$ and $td_y(\emptyset) = 0$ the Formula 3 holds.

5 Affine cones

Let us extend the torus action by adding one factor to \mathbb{T} . Now we consider $\mathbb{T} = (\mathbb{C}^*)^{m+1}$ the character of the additional coordinate of \mathbb{T} is denoted by t and $T = e^{-t}$. The weights of the action on \mathbb{C}^n are

$$(t + t_{-m}, t - t_{-m+1}, \dots, t + t_{m-1}, t - t_m)$$

in the even case and

$$(t + t_{-m}, t - t_{-m+1}, \dots, t, \dots, t + t_{m-1}, t - t_m)$$

in the odd case. It does not change the action on \mathbb{P}^{n-1} on which the additional coordinate of \mathbb{T} acts trivially.

Formula 8 Consider the complements of the affine cones $Q_n^* = \mathbb{C}^n \setminus \widehat{Q}_n$ and $X_n^* = \mathbb{C}^n \setminus \widehat{X}_n$. In the equivariant cohomology $H_{\mathbb{T}}^*(\mathbb{C}^n)[y]$ we have the equation

$$td_y^{\mathbb{T}}(X_n^*) - td_y^{\mathbb{T}}(Q_n^*) = y td_y^{\mathbb{T}}(Q_{n-2}^*)$$

for $n \geq 2$.

Proof. Let Y denote X_n , Q_n or Q_{n-2} . Let $\pi : \widetilde{\mathbb{C}}^n \rightarrow \mathbb{C}^n$ be the blowup at the origin with $i : \mathbb{P}^{n-1} \hookrightarrow \widetilde{\mathbb{C}}^n$ the inclusion of the exceptional divisor. The Hirzebruch class of $Y^* \subset \mathbb{C}^n$ can be computed by push-forward of the class $td_y^{\mathbb{T}}(\pi^{-1}(Y^*))$, since $\pi : \widetilde{\mathbb{C}}^n \setminus \mathbb{P}^{n-1} \rightarrow \mathbb{C}^n \setminus \{0\}$ is an isomorphism. Here we are using functoriality of the equivariant Hirzebruch classes. The projection $p : \widetilde{\mathbb{C}}^n \rightarrow \mathbb{P}^{n-1}$ has a structure of a vector bundle $\nu = \mathcal{O}(-1)$. We apply the formula (7) and additivity for $\pi^{-1}(Y^*) = p^{-1}(Y') \setminus i(Y')$:

$$td_y^{\mathbb{T}}(Y^*) = \pi_* td_y^{\mathbb{T}}(\pi^{-1}(Y^*)) = \pi_* p^* \left((td_y^{\mathbb{T}}(\nu) - c_1(\nu)) \cdot td_y^{\mathbb{T}}(Y') \right).$$

The expression is linear with respect to $td_y^{\mathbb{T}}(Y')$. It follows that the linear relation (Formula 3) among Hirzebruch classes $td_y^{\mathbb{T}}(X_n^*)$, $td_y^{\mathbb{T}}(Q_n^*)$ and $td_y^{\mathbb{T}}(Q_{n-2}^*)$ in $H_{\mathbb{T}}^*(\mathbb{P}^{n-1})[y]$ implies the corresponding relation in $H_{\mathbb{T}}^*(\mathbb{C}^n)[y]$. \square

Remark 9 More generally for the degeneration

$$\lambda \sum_{i=1}^k x_{-i} x_i + \sum_{i=k+1}^m x_{-i} x_i$$

(and similarly for n odd) we have

$$td_y^{\mathbb{T}}(Q_n^*) - td_y^{\mathbb{T}}(Y^*) = (-y)^{m-k} td_y^{\mathbb{T}}(Q_{2k}^*)$$

where Y is the hypersurface corresponding to $\lambda = 0$. The general case follows from the case $k = m - 1$, which was studied here.

We obtain the explicit formula

Corollary 10 *The equivariant Hirzebruch class of the complement of the quadratic cone $Q_n^* = \mathbb{C}^n \setminus \widehat{Q}_n$ is equal to for $n = 2m$*

$$td_y^{\mathbb{T}}(Q_n^*) = \sum_{k=0}^{m-1} (-y)^k td_y^{\mathbb{T}}(X_{n-2k}^*)$$

for $n = 2m + 1$

$$td_y^{\mathbb{T}}(Q_n^*) = \sum_{k=0}^{m-1} (-y)^k td_y^{\mathbb{T}}(X_{n-2k}^*) + (-y)^m td_y^{\mathbb{T}}(\mathbb{C} \setminus 0),$$

where

$$\frac{td_y^{\mathbb{T}}(X_{2m}^*)}{eu(0)} = (\Phi(TT_m) - 1) \cdot (\Phi(TT_m^{-1}) - 1) \cdot \prod_{j=1}^{m-1} \Phi(TT_j) \Phi(TT_j^{-1})$$

and

$$\frac{td_y^{\mathbb{T}}(X_{2m+1}^*)}{eu(0)} = \Phi(T) \cdot (\Phi(TT_m) - 1) \cdot (\Phi(TT_m^{-1}) - 1) \cdot \prod_{j=1}^{m-1} \Phi(TT_j) \Phi(TT_j^{-1}),$$

$$\frac{td_y^{\mathbb{T}}(\mathbb{C} \setminus 0)}{eu(0)} = \Phi(T) - 1.$$

6 Positivity

Now we will show that the Hirzebruch classes of \widehat{Q}_n and Q_n^* satisfy certain positivity condition. For a weight $w \in \text{Hom}(\mathbb{T}, \mathbb{C}^*)$ let us set a new variable $S_w = e^{-w} - 1$. Also let us set $\delta = -1 - y$.

Corollary 11 *The Hirzebruch class of the complement of the affine cone $Q_n^* = \mathbb{C}^n \setminus \widehat{Q}_n$ is equal to a polynomial in δ and S_w with nonnegative coefficients divided by the product of the variables S_w , where w are the weights of the representation \mathbb{C}^n .*

Proof. It suffices to note that for the standard action of one dimensional torus on \mathbb{C} we have (with $T = e^{-t}$ as before)

$$\frac{td_y^{\mathbb{T}}(\mathbb{C})}{eu(0)} = \Phi(T) = \frac{1 - T + (1 + y)(T - 1 + 1)}{1 - T} = \frac{S_t + \delta(S_t + 1)}{S_t}$$

and

$$\frac{td_y^{\mathbb{T}}(\mathbb{C} \setminus \{0\})}{eu(0)} = \frac{td_y^{\mathbb{T}}(\mathbb{C})}{eu(0)} - \frac{td_y^{\mathbb{T}}(\{0\})}{eu(0)} = \Phi(T) - 1 = \frac{\delta(S_t + 1)}{S_t}.$$

Moreover, since $\widehat{X}'_{n-2k} = \mathbb{C}^{n-2-2k} \times (\mathbb{C}^*)^2$ for $k = 0, \dots, m-1$, by multiplicativity, the Hirzebruch class $td_y^{\mathbb{T}}(\widehat{X}'_n)$ is a nonnegative expression. The claim for Q_n^* follows from Corollary 10. \square

For the original closed varieties we have:

Formula 12

$$td_y^{\mathbb{T}}(\widehat{Q}_n) - td_y^{\mathbb{T}}(\widehat{X}_n) = y \left(td_y^{\mathbb{T}}(\mathbb{C}^{n-2}) - td_y^{\mathbb{T}}(\widehat{Q}_{n-2}) \right).$$

Proof. We rewrite the Formula 8 passing to the complement

$$\left(td_y^{\mathbb{T}}(\mathbb{C}^n) - td_y^{\mathbb{T}}(\widehat{X}_n) \right) - \left(td_y^{\mathbb{T}}(\mathbb{C}^n) - td_y^{\mathbb{T}}(\widehat{Q}_n) \right) = y \left(td_y^{\mathbb{T}}(\mathbb{C}^{n-2}) - td_y^{\mathbb{T}}(\widehat{Q}_{n-2}) \right).$$

Hence we obtain what is claimed. \square

Corollary 13 *The Hirzebruch class of the affine cone of \widehat{Q}_n is equal to a polynomial in δ and S_w with nonnegative coefficients divided by the product of the variables S_w , where w are the weights of the representation \mathbb{C}^n .*

Proof. Transforming the Formula 12 we obtain that

$$\begin{aligned} td_y^{\mathbb{T}}(\widehat{Q}_n) &= -y td_y^{\mathbb{T}}(\widehat{Q}_{n-2}) + \left(td_y^{\mathbb{T}}(\widehat{X}_n) + y td_y^{\mathbb{T}}(\mathbb{C}^{n-2}) \right) \\ &= -y td_y^{\mathbb{T}}(\widehat{Q}_{n-2}) + td_y^{\mathbb{T}}(\mathbb{C}^{n-2}) \cdot \frac{-(1+y)(T^2 - 1)}{(1 - TT_m^{-1})(1 - TT_m)}. \end{aligned} \quad (14)$$

Here we use additivity and multiplicativity of the Hirzebruch class applied to the decomposition $\widehat{X}^n = \mathbb{C}^{n-2} \times (\mathbb{C}_+ \cup \mathbb{C}_- \setminus \{0\})$ with

$$\frac{td_y^{\mathbb{T}}(\mathbb{C}_{\pm})}{eu(0)} = \frac{1 + yTT_m^{\pm 1}}{1 - TT_m^{\pm 1}}.$$

The formula (14) follows from the identity

$$\frac{1 + yTT_m}{1 - TT_m} + \frac{1 + yTT_m^{-1}}{1 - TT_m^{-1}} - 1 + y = \frac{-(1 + y)(T^2 - 1)}{(1 - TT_m^{-1})(1 - TT_m)}.$$

We note that

$$\frac{-(1 + y)(T^2 - 1)}{(1 - TT_m)(1 - TT_m^{-1})} = \frac{\delta(S_t^2 + 2S_t)}{S_{t+t_m}S_{t-t_m}}$$

is a positive expression. We proceed inductively having in mind that the coefficient before $td_y^T(\tilde{Q}_{n-2})$ is $-y = 1 + \delta$. \square

The Corollaries 11 and 13 confirm the general rule (not proved so far) that the local Hirzebruch classes of Schubert cells are positive expressions in the variables associated with tangent weights.

References

- [Al] P. Aluffi, *Differential forms with logarithmic poles and Chern-Schwartz-MacPherson classes of singular varieties*. C. R. Acad. Sci. Paris Sér. I Math., 329(7)
- [AlMa] P. Aluffi, M. Marcolli, *Algebraic-geometric Feynman rules*. Int. J. Geom. Methods Mod. Phys., 8, 203-237, (2011)
- [AtBo] M. Atiyah, R. Bott, *The moment map and equivariant cohomology* *Topology*. 23 (1984) 1-28.
- [Bo] A. Borel, *Seminar on transformation groups*. Annals of Mathematics Studies, No. 46, Princeton University Press (1960)
- [BeVe] N. Berline, M. Vergne, *Classes caractéristiques équivariantes. Formule de localisation en cohomologie équivariante*. C.R. Acad. Sc. Paris 295 (1982), 539-541.
- [BSY] J-P. Brasselet, J. Schürmann, S. Yokura, *Hirzebruch classes and motivic Chern classes for singular spaces*. J. Topol. Anal. 2, No. 1, 1-55 (2010)
- [Co1] I. Coskun, *Rigid and non-smoothable Schubert classes*. J. Diff. Geom. 87 (2011), 493-514.
- [Co2] I. Coskun, *Rigidity of Schubert classes in orthogonal grassmannians*. Israel J. Math. 200 (2014), 85-126.
- [CoVa] I. Coskun, R. Vakil, *Geometric positivity in the cohomology of homogeneous spaces and generalized schubert calculus*. Algebraic geometry, Seattle 2005. Part 1, 77-124, Proc. Sympos. Pure Math., 80, Part 1, Amer. Math. Soc., Providence, RI, 2009.

- [CMSS] S. Cappell, L. Maxim, J. Schürmann, J. Shaneson, *Characteristic classes of complex hypersurfaces*. Adv. Math. 225, No. 5, 2616-2647 (2010)
- [EdGr] D. Edidin, W. Graham, *Localization in equivariant intersection theory and the Bott residue formula*. Am. J. Math. 120, No.3, 619-636 (1998)
- [Fu] W. Fulton, *Intersection Theory*. Springer 1998
- [GKM] M. Goresky, R. Kottwitz, R. MacPherson, *Equivariant Cohomology, Koszul Duality, and the Localization Theorem*. Invent. Math. 131, No.1, (1998), 25-83
- [MSS] L. Maxim, M. Saito, J. Schürman, *Hirzebruch-Milnor classes of complete intersections*. Adv. Math. 241 (2013), 220245.
- [Oh] T. Ohmoto, *A note on the Chern-Schwartz-MacPherson class*. Singularities in geometry and topology, 117-131, IRMA Lect. Math. Theor. Phys., 20, Eur. Math. Soc., Zürich, 2012
- [PaPr] A. Parusiński, P. Pragacz, *Characteristic classes of hypersurfaces and characteristic cycles*. J. Algebraic Geom. 10 (2001), no. 1, 63-79.
- [Qu] D. Quillen, *The Spectrum of an Equivariant Cohomology Ring: I*. Ann. Math., Vol. 94, No. 3, 549-572
- [Ve] J.-L. Verdier, *Specialisation des classes de Chern*. 149-159, Astérisque, 82-83, Soc. Math. France, Paris, 1981
- [We1] A. Weber, *Equivariant Chern classes and localization theorem*. Journal of Singularities, Vol. 5, 153-176, (2012)
- [We2] A. Weber, *Computing equivariant characteristic classes of singular varieties*. RIMS Kkyuroku 1868, 109-129, (2013)
- [We3] A. Weber, *Equivariant Hirzebruch class for singular varieties*. arXiv:1308.0788