

Tutorial 6

13.11

1. Reachability for VASS with negative semantic belongs to NP.

Brief theoretical introduction (hint):

Linear programming (LP)

a minimization / maximization problem for a linear function of n arguments: x_1, x_2, \dots, x_n , where the arguments have to satisfy some conditions (linear equations or inequalities using x_i s)

Example:

$$\max \quad x_1 + 2x_2$$

$$\text{s.t.} \quad x_2 \leq x_1 + 2$$

$$2x_1 + x_2 \leq 4$$



solvable in
polynomial
time

Integer programming (IP)

↑ for each $i \in \{1, 2, \dots, n\}$ we add a condition $x_i \in \mathbb{Z}$
this problem is NP-complete

Given: VASS (Q, T)

Question: $(q, v) \xrightarrow{*} (q', v')$?

does there exist a run that can drop below zero

Attempt 1

* denote transition vectors by v_1, v_2, \dots, v_n

* solve a set of linear equations

$$\begin{bmatrix} \vdots & \vdots & \vdots & \vdots \\ v_1 & v_2 & \dots & v_n \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = v' - v$$

where x_i is the number of times transition t_i is to be fired (used)

Problem: we can obtain $x_i \notin \mathbb{Z}$ and $x_i < 0$

Attempt 2

and the goal of IP: $\min \sum_{i=1}^n x_i$

* add conditions: $x_i \in \mathbb{Z}$, $x_i \geq 0$ for each i

Problem: firing each transition t_i x_i times


gives $v' - v$ but the transitions may

not form a run in VASS

Attempt 3

- * add Kirchoff's law conditions - for each state except q, q' the number of incoming transitions (sum of x_i) equals the number of outgoing ones
- * for q there is one more outgoing transition and for q' - one more incoming

Problem: x_i can describe a path from q to q' and a disjoint cycle...




Attempt 4

- * after obtaining x_i 's, define $S = \{t_i \in T \mid x_i > 0\}$ and check whether each $t_i \in S$ can be reached from q using transitions in S
- * the condition above ensures that we can create the run given x_i 's (go from q using any available transition until reaching q' with the last transition incoming to q' ; then, if we still have some spare transitions, we can create some cycles and glue them to the run $q \xrightarrow{*} q'$)
- * it seems like it should be enough for S to be connected in order to create such a run

Problem: solution to our LP program may not satisfy the condition from attempt 4, but there can be some other run that does

received solution,
unable to create
a run based on it



another feasible
solution that corresponds
to a run but doesn't
minimize the goal function

Attempt 5 - final approach

- * to ensure that every possible run is taken into account we can guess the transition set S' that we allow the use of and then solve IP with conditions

$$\forall i \in S' \quad x_i > 0$$

$$\forall i \in T \setminus S' \quad x_i = 0$$

2. Using the insights from the previous problem present a way to check condition Θ_1

Θ_1 : for every $m \geq 1$ $(q, v) \dashrightarrow^* (q', v')$
using every transition $\geq m$ times

- * if there exists a run r' for some selected $m' > 0$ using every transition t_i $x_i \geq m'$ times, then there also exists a run r'' that uses each transition at least $m'' = \max(x_i) + 1$ times (we use notation $\# t_i \rightarrow y_i$ for r'')

- * both r' and r'' satisfy the Kirchhoff's law, hence so does $r'' - r'$ which uses each transition t_i at least $y_i - x_i > 1$ times
- * thus, $r'' - r'$ is a cycle that has no side effect on a given configuration
- * the solution idea:
 - define two sets of variables: x_1, x_2, \dots, x_n and $\Delta_1, \Delta_2, \dots, \Delta_n$ such that both sets satisfy Kirchhoff's law (first with a twist for q and q')
 - $x_i > 0$ (for a subset of transitions - as discussed before), firing all t_i s x_i times results in $v' - v$ change on the configuration
 - $\Delta_i > 0$ for all transitions and firing t_i s don't change the configuration

3. Prove that the following implication holds:

$\exists m$. every configuration reachable from (q, v) has some coordinate $< m$] (1)

\Downarrow

$\exists m$. every run from (q, v) , on some coordinate, is always $< m$] (2)


* if (1) holds, we can set m to be the greatest integer appearing in the coverability tree (it might take a lot of time to find m but we can afford it; also, for VASSes we have to remember the state we are in)

* now, we show that such an m satisfies (2)

* first, assume that the VASS is 2-dimensional and assume that there exists a run r such that r goes through points (v_1, v_2) with $v_1 \geq m$ and (u_1, u_2) with $u_2 \geq m$

* it means that we had to reach $(p, (w, -))$ in the coverability tree \Rightarrow there exists a sequence σ of transitions that starts and ends in p , increases the first coordinate and doesn't change the second one

* we can fire σ as many times as it is required to increase the first coordinate by $\geq m$, i.e., we extend r in such a way



* it gives us a new run r' that instead of (u_1, u_2) reaches (u_1+k, u_2) with $k \geq m$

* this configuration doesn't satisfy (1) - contradiction

* for higher dimensions the reasoning is similar (the only problem: going from (w, x_1, \dots, x_d) to (w, w, y_1, \dots, y_d) might decrease first coordinate)

4. Show reductions between the following problems:


1) reachability problem: $M \in \mathbb{N}^r$, is M reachable? (RP)

2) submarking reachability problem: \tilde{P} - subset of places, $M_{\tilde{P}}$ - configuration over \tilde{P} , does there exist a reachable configuration M such that over the places of \tilde{P} it agrees with $M_{\tilde{P}}$? (SRP)


3) zero reachability problem: is $\mathbf{0}$ reachable? (ZRP)

4) single-place zero reachability problem: given a place p , does there exist a reachable configuration M , s.t. $M(p) = 0$? (SPZRP)

The idea is to show the following reductions:



"Y is reducible to X"




\rightarrow easy reduction

\Rightarrow reduction to present

SP2RP \Rightarrow ZRP:

- * now, we can use the blackbox for answering single-place zero reachability problem and have to answer the zero reachability problem in N
- * the idea is to have a new place Π s.t. at all times Π contains as many tokens as there are in all places of N , i.e., at every configuration M in the new net N' :
$$M(\Pi) = \sum_{i=1}^r M(p_i)$$
- * to obtain it, we connect Π with all transitions of N with incoming and outgoing arcs - the weight of incoming arc is the sum of incoming arcs' weights in N and analogously for outgoing
- * initially, Π contains the total number of tokens in configuration M_0 for N
- * obviously, $t_j \in T$ is fireable in N iff t_j is fireable in N'
- * answer for ZRP in $N =$ answer for SP2RP in N'

* obtaining N' from N - example



Source of the net modification figures:

Michel Hack. Decidability Questions for Petri Nets

1.^D ZRP \Rightarrow SRP: homework (not obligatory)

2.^D Structural unboundedness for general Petri nets belongs to NP.

homework (not obligatory)