

# ON TWO CONSEQUENCES OF CH ESTABLISHED BY SIERPIŃSKI. II

R. POL AND P. ZAKRZEWSKI

ABSTRACT. We continue a study of the relations between two consequences of the Continuum Hypothesis discovered by Waław Sierpiński, concerning uniform continuity of continuous functions and uniform convergence of sequences of real-valued functions, defined on subsets of the real line of cardinality continuum.

## 1. INTRODUCTION

In [11] we studied the following two consequences of the Continuum Hypothesis (CH) distinguished by Waław Sierpiński in his classical treaty *Hypothèse du continu* [14] (the notation is taken from [14]):

- $C_8$  There exists a continuous function  $f : E \rightarrow \mathbb{R}$ ,  $E \subseteq \mathbb{R}$ ,  $|E| = \mathfrak{c}$ , not uniformly continuous on any uncountable subset of  $E$ .
- $C_9$  There is a sequence of functions  $f_n : E \rightarrow \mathbb{R}$ ,  $E \subseteq \mathbb{R}$ ,  $|E| = \mathfrak{c}$ , converging pointwise but not converging uniformly on any uncountable subset of  $E$ .

Sierpiński [13] checked that  $C_8$  implies  $C_9$ . The status of the converse implication remains unclear. Let us notice, however, that in *Topology I* by Kuratowski [7], footnote (3) on page 533 suggests that the two statements are in fact equivalent.

In [11] we considered the following stratifications of statements  $C_8$  and  $C_9$  for uncountable cardinals  $\kappa \leq \lambda \leq \mathfrak{c}$ :

- $C_8(\lambda, \kappa)$  There exist a set  $E \subseteq \mathbb{R}$  of cardinality  $\lambda$  and a continuous function  $f : E \rightarrow \mathbb{R}$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ .
- $C_9(\lambda, \kappa)$  There exist a set  $E \subseteq \mathbb{R}$  of cardinality  $\lambda$  (equivalently: for any set  $E \subseteq \mathbb{R}$  of cardinality  $\lambda$ ) and there is a sequence of functions  $f_n : E \rightarrow \mathbb{R}$ , converging on  $E$  pointwise but not converging uniformly on any subset of  $E$  of cardinality  $\kappa$ .

In particular, we proved in [11] that:

- $C_8(\mathfrak{c}, \mathfrak{c}) \Leftrightarrow C_9(\mathfrak{c}, \mathfrak{c})$ , and each of these statements is equivalent to the assertion  $\mathfrak{d} = \mathfrak{c}$ , provided that the cardinal  $\mathfrak{c}$  is regular.

---

*Date:* May 25, 2024.

*2020 Mathematics Subject Classification.* 03E20, 03E15, 26A15, 54C05 54F45 .

*Key words and phrases.* uniform continuity, uniform convergence, K-Lusin set, Henderson compactum.

- $C_8(\aleph_1, \aleph_1) \Leftrightarrow C_9(\aleph_1, \aleph_1)$ , and each of these statements is equivalent to the assertion  $\mathfrak{b} = \aleph_1$ .

Here  $\mathfrak{d}$  and  $\mathfrak{b}$  denote, as usual, the smallest cardinality of a dominating and, respectively, an unbounded family in  $\mathbb{N}^{\mathbb{N}}$  corresponding to the ordering of eventual domination  $\leq^*$  (cf. [4]).

An important role in our considerations was played by the notion of a  $\kappa$ -Lusin set (cf. [1]) which we extended declaring that an uncountable subset  $E$  of a Polish space  $X$  is a  $\kappa$ - $K$ -Lusin set in  $X$ ,  $\aleph_1 \leq \kappa \leq \mathfrak{c}$ , if  $|E \cap K| < \kappa$  for every compact set  $K \subseteq X$ . We proved in [11] that  $C_9(\lambda, \kappa)$  is equivalent to the statement that there is a Polish space  $X$  and a  $\kappa$ - $K$ -Lusin set of cardinality  $\lambda$  in  $X$ .

In this note we present additional results related to the subject of [11]. Most of them were earlier announced in [11, Section 4].

In Sections 2, 3 and 4 we investigate  $C_8$ -like phenomena in a more general setting. We are interested in two closely related problems ( $X$  and  $Y$  are fixed separable metric spaces).

*Problem 1.* Is the existence of a set  $E \subseteq X$  of cardinality  $\lambda$  and a continuous function on  $E$  with values in  $Y$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ , related to either  $C_8(\lambda, \kappa)$  or  $C_9(\lambda, \kappa)$ ?

*Problem 2.* Does the existence of a set  $E \subseteq X$  of cardinality  $\lambda$  and a continuous function on  $E$  with values in  $Y$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ , imply that there also exists such a function on  $E$  with values in  $\mathbb{R}$ ?

Concerning Problem 1, we observe that the existence of a separable metric space  $X$  of cardinality  $\lambda$ , a metric space  $Y$ , and a continuous function on  $X$  with range in  $Y$ , which is not uniformly continuous on any subset of  $X$  of cardinality  $\kappa$ , already implies that there exists a  $\kappa$ - $K$ -Lusin set of cardinality  $\lambda$  in some Polish space and, consequently, that  $C_9(\lambda, \kappa)$  holds true (cf. Proposition 2.1).

Conversely,  $C_9(\lambda, \kappa)$  implies that there exists a  $\kappa$ - $K$ -Lusin set  $E$  of cardinality  $\lambda$  in  $\mathbb{P}$ , the set of irrationals of the unit interval  $I = [0, 1]$ , such that for every non- $\sigma$ -compact Polish space  $Y$  there is a continuous function on  $E$  which is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$  (cf. Theorem 2.2).

On the other hand, if  $Y$  is compact, then the existence of a set  $E \subseteq I$  of cardinality  $\lambda$ , and a continuous function  $f : E \rightarrow Y$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ , implies  $C_8(\lambda, \kappa)$  (cf. Theorem 3.2).

Concerning Problem 2, we show that if a set  $E$  in the Hilbert cube  $I^{\mathbb{N}}$  is zero-dimensional and there exists a continuous function on  $E$  with range in any uncountable compact metric space  $Y$ , not uniformly continuous on any subset of  $X$  of cardinality  $\kappa$ , then there is also such

a function with range in any uncountable compact metric space  $Z$ , and in particular, in  $I$  (cf. Theorem 3.1).

On the other hand, assuming CH, we prove the existence of a set  $E \subseteq I^{\mathbb{N}}$  of cardinality  $\mathfrak{c}$  such that there is a continuous function  $f : E \rightarrow I^{\mathbb{N}}$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\mathfrak{c}$  but each continuous function  $g : E \rightarrow \mathbb{R}$  is constant on a subset of  $E$  of cardinality  $\mathfrak{c}$ . The construction of a witnessing pair  $E$  and  $f$  falls under a general scheme, described in Section 4, of constructions of  $C_8$ -like examples based on a generalization of the notion of a  $\kappa$ -K-Lusin set.

Section 5 is a slight departure from the topic but it is closely related to a reasoning of Sierpiński concerning  $C_9$ . We shall show that a Hausdorff space  $X$  is Čech-complete and Lindelöf if and only if there is a sequence  $f_0 \geq f_1 \geq \dots$  of continuous functions  $f_n : X \rightarrow I$  converging pointwise to zero but not converging uniformly on any closed non-compact set in  $X$  (cf. Theorem 5.2). The existence of such a sequence for any Polish space  $X$  was a key step in proving that statement  $C_9(\lambda, \kappa)$  is equivalent to the existence of a  $\kappa$ -K-Lusin set  $E$  of cardinality  $\lambda$  in  $\mathbb{P}$  (cf. [11, Theorem 2.3]).

In this note  $\mathbb{P}$  always denotes the set of irrationals of the unit interval  $I = [0, 1]$ . It is homeomorphic to the Baire space  $\mathbb{N}^{\mathbb{N}}$ , the countable product of the set of natural numbers  $\mathbb{N} = \{0, 1, 2, \dots\}$  with the discrete topology (cf. [6]).

## 2. MAPPINGS INTO NON-COMPACT SPACES AND $C_9$

We start with a general observation.

**Proposition 2.1.** *If there exist a separable metric space  $(X, d_X)$  of cardinality  $\lambda$ , a metric space  $(Y, d_Y)$ , and a continuous function on  $X$  with range in  $Y$ , which is not uniformly continuous on any subset of  $X$  of cardinality  $\kappa$ , then there exists a  $\kappa$ -K-Lusin set of cardinality  $\lambda$  in some Polish space and, consequently,  $C_9(\lambda, \kappa)$  holds true.*

*Proof.* Let  $f : X \rightarrow Y$  be a continuous function which is not uniformly continuous on any subset of  $X$  of cardinality  $\kappa$ . The function  $f$  extends (cf. [6, Theorem 3.8]) to a continuous function  $\tilde{f} : G \rightarrow \hat{Y}$  over a  $G_\delta$ -set  $G$  (with  $X \subseteq G$ ) in the Polish completion  $(\hat{X}, \hat{d}_X)$  of  $(X, d_X)$  into the completion  $(\hat{Y}, \hat{d}_Y)$  of  $(Y, d_Y)$ . Now, if  $K \subseteq G$  is compact, then  $\tilde{f}$  is uniformly continuous on  $K$ , hence  $f$  is uniformly continuous on  $K \cap X$ . By the choice of  $X$ , we have that  $|K \cap X| < \kappa$ , which shows that  $X$  is a  $\kappa$ -K-Lusin set of cardinality  $\lambda$  in  $G$ . By [11, Theorem 2.3], this proves  $C_9(\lambda, \kappa)$ . □

On the other hand, statement  $C_9(\lambda, \kappa)$  already implies (and in view of Proposition 2.1, is equivalent to) the existence of  $C_9$ -like example for functions with values in arbitrary non- $\sigma$ -compact Polish spaces.

**Theorem 2.2.** *For any uncountable cardinals  $\kappa \leq \lambda \leq \mathfrak{c}$  the following are equivalent:*

- (1)  $C_9(\lambda, \kappa)$ ,
- (2) *there exist a set  $E \subseteq \mathbb{R}$  of cardinality  $\lambda$ , a non- $\sigma$ -compact Polish space  $Y$ , and a continuous function  $f : E \rightarrow Y$  which is not uniformly continuous (with respect to arbitrary complete metric on  $Y$ ) on any subset of  $E$  of cardinality  $\kappa$ .*

*Moreover, any  $\kappa$ -K-Lusin set  $E$  of cardinality  $\lambda$  in  $\mathbb{P}$  has the property expressed in (2) with respect to any non- $\sigma$ -compact Polish space  $Y$ .*

*Proof.* (1)  $\Rightarrow$  (2). By Theorem [11, Theorem 2.3], statement  $C_9(\lambda, \kappa)$  implies that there is a  $\kappa$ -K-Lusin set  $E$  of cardinality  $\lambda$  in  $\mathbb{P}$ .

Let  $Y$  be an arbitrary non- $\sigma$ -compact Polish space. Let  $h : \mathbb{P} \rightarrow Y$  be a homeomorphic embedding of  $\mathbb{P}$  onto a closed subspace  $h(\mathbb{P})$  of  $Y$  (cf. [6, Theorem 7.10]). We will show that  $E$  together with  $f = h|_E$  have the required properties.

To that end, let us fix a set  $A \subseteq E$  with  $|A| = \kappa$ . Then  $\bar{A}$ , the closure of  $A$  in  $I$ , is not contained in  $\mathbb{P}$  since otherwise  $\bar{A}$  would be a compact set in  $\mathbb{P}$ , intersecting  $E$  on a set of cardinality  $\kappa$ . So let us pick  $a_k \in A$ ,  $k \in \mathbb{N}$ , such that  $\lim_{k \rightarrow \infty} a_k = a$  and  $a \in I \setminus \mathbb{P}$ .

Now, if  $f$  were uniformly continuous on  $A$  with respect to a complete metric  $d$  on  $Y$ ,  $f$  would take Cauchy sequences in  $A$  to Cauchy sequences in  $Y$ . In particular, the sequence  $(f(a_n))_{n \in \mathbb{N}}$  would be Cauchy in  $Y$ , hence  $\lim_{k \rightarrow \infty} f(a_k) = z$  for some  $z \in Y$ . This, however, is not the case: since the set  $\{a_0, a_1, \dots\}$  has no accumulation point in  $\mathbb{P}$ , the set  $\{f(a_0), f(a_1), \dots\}$  has no accumulation point in  $h(\mathbb{P})$  and hence also in  $Y$ , as  $h(\mathbb{P})$  is closed in  $Y$ .

The implication (2)  $\Rightarrow$  (1) follows immediately from Proposition 2.1.  $\square$

### 3. MAPPINGS INTO COMPACT SPACES AND $C_8$

The results of the previous section show that  $C_8$ -like statements for functions with values in non- $\sigma$ -compact Polish spaces are actually equivalent to statement  $C_9$ . Apparently, the situation changes when we consider functions with values in compact spaces. As the following result shows, if a set  $E \subseteq I^{\mathbb{N}}$  is zero-dimensional and there exists a continuous function on  $E$  with range in any uncountable compact metric space  $Y$ , not uniformly continuous on any subset of  $X$  of cardinality  $\kappa$ , then there is also such a function with values in  $\mathbb{R}$ , witnessing that  $C_8(\lambda, \kappa)$  holds true.

**Theorem 3.1.** *Let  $E$  be a zero-dimensional subset of a compact metric space  $X$ . If there exists a continuous function on  $E$  with range in a compact metric space  $Y$ , not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ , then there is also such a function with range in the Cantor ternary set  $C$  in  $I$ . Consequently, for any uncountable compact metric space  $Z$ , there is also such a function with values in  $Z$ .*

*Proof.* Let  $h : E \rightarrow Y$  be a continuous function not uniformly continuous on any set of cardinality  $\kappa$ .

Using the compactness of  $Y$ , let us fix a sequence  $(K_n, L_n)_{n \in \mathbb{N}}$  of pairs of disjoint compact sets in  $Y$  such that for any pair  $(K, L)$  of disjoint compact sets in  $Y$ , there is  $n$  with  $K \subseteq K_n$  and  $L \subseteq L_n$ .

For each  $n \in \mathbb{N}$ , we let  $C_n = h^{-1}(K_n)$ ,  $D_n = h^{-1}(L_n)$ , and using the fact that  $E$  is zero-dimensional and separable, we choose a continuous function  $u_n : E \rightarrow \{0, 2\}$  taking on  $C_n$  value 0 and on  $D_n$  value 2, cf. [3, Theorem 1.2.6].

We shall show that the function  $f : E \rightarrow I$  defined by the formula

$$f(x) = \sum_{i=0}^{\infty} \frac{1}{3^{i+1}} u_i(x),$$

which takes values in the Cantor ternary set  $C$ , is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ .

To that end, let us fix a set  $A \subseteq E$  with  $|A| = \kappa$ .

We shall first make the following observation. Let  $a \in \bar{A}$ , the closure of  $A$  in  $X$ . Then for any  $n$ , since  $u_n$  takes on  $A$  values 0 or 2 only, the oscillation

$$\inf\{\text{diam}(u_n(A \cap V)) : V \text{ is a neighbourhood of } a \text{ in } X\}$$

of  $u_n|_A$  at  $a$  is either 0 or 2.

Let us note that  $h|_A : A \rightarrow Y$  cannot be extended to a continuous (hence, by the compactness of  $\bar{A}$ , uniformly continuous) function  $\bar{h} : \bar{A} \rightarrow Y$ , since otherwise the function  $h|_A = \bar{h}|_A$  would itself be uniformly continuous, contrary to the fact that  $|A| = \kappa$ . Consequently, there must be closed disjoint sets  $K, L$  in  $Y$  such that the closures of  $(h|_A)^{-1}(K)$  and  $(h|_A)^{-1}(L)$  in  $X$  meet, cf. [2, Theorem 3.2.1]. Then for an  $n$  with  $K \subseteq K_n$  and  $L \subseteq L_n$ , we infer that  $\overline{C_n \cap A} \cap \overline{D_n \cap A} \neq \emptyset$ , so let us fix  $a \in \overline{C_n \cap A} \cap \overline{D_n \cap A}$ .

It follows that  $u_n|_A$  has the oscillation at  $a$  equal to 2 and let us assume that  $n$  is the smallest index with this property. The oscillation of each of the functions  $u_0|_A, \dots, u_{n-1}|_A$  is then equal to 0, hence we can find a neighbourhood  $V$  of  $a$  in  $X$  such that all these functions have constant values on  $A \cap V$ .

Let us pick  $x_k, y_k \in A \cap V$ ,  $k \in \mathbb{N}$ , such that  $\lim_{k \rightarrow \infty} x_k = \lim_{k \rightarrow \infty} y_k = a$  and for every  $k$ ,  $u_n(x_k) = 0$ ,  $u_n(y_k) = 2$ , hence

$$\begin{aligned} f(y_k) - f(x_k) &= \frac{2}{3^{n+1}} + \sum_{i=n+1}^{\infty} \frac{1}{3^{i+1}} (u_i(y_k) - u_i(x_k)) \geq \\ &\frac{2}{3^{n+1}} - \sum_{i=n+1}^{\infty} \frac{2}{3^{i+1}} = \frac{1}{3^{n+1}}. \end{aligned}$$

It follows that the oscillation of  $f|A$  at  $a$  is at least  $3^{-n-1}$ . In effect,  $f|A$  has no continuous extension over  $\bar{A}$ , which means that  $f$  is not uniformly continuous on  $A$ .

Finally, if  $(Z, d)$  is an arbitrary compact metric space and  $e : C \rightarrow Z$  is a homeomorphic embedding of  $C$  into  $Z$ , then since  $e^{-1}$  is uniformly continuous, the function  $e \circ f : E \rightarrow Z$  has desired properties.  $\square$

As an immediate corollary we obtain the following equivalent form of statement  $C_8(\lambda, \kappa)$ .

**Theorem 3.2.** *For any uncountable cardinals  $\kappa \leq \lambda \leq \mathfrak{c}$  if the cofinality of  $\lambda$  is uncountable, then the following are equivalent:*

- (1)  $C_8(\lambda, \kappa)$ ,
- (2) *there exists a set  $E \subseteq \mathbb{R}$  of cardinality  $\lambda$ , a compact metric space  $Y$  and a continuous function  $f : E \rightarrow Y$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ .*

Moreover, any set  $E \subseteq I$  that witnesses  $C_8(\lambda, \kappa)$  for some continuous function from  $E$  to  $I$  has the property expressed in (2) with respect to any uncountable compact metric space.

*Proof.* (1)  $\Rightarrow$  (2). If  $f : E \rightarrow \mathbb{R}$  is a continuous function on a set  $E \subseteq \mathbb{R}$  of cardinality  $\lambda$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ , then since the cofinality of  $\lambda$  is uncountable, by shrinking  $E$ , if necessary, we may assume that the range of  $f$  is contained in a closed interval  $Y$  of length 1.

If additionally  $E \subseteq I$  and  $f : E \rightarrow I$ , then since  $E$  contains no non-trivial interval, it is zero-dimensional, and Theorem 3.1 applies.

(2)  $\Rightarrow$  (1). Now let  $f : E \rightarrow Y$  be a continuous function with values in a compact metric space  $Y$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ . We may again assume that  $E$  is contained in a closed interval  $X$ . Then,  $E$  being zero-dimensional, it is enough to apply the final part of the assertion of Theorem 3.1 to  $Z = I$ .  $\square$

By the results of [11] (cf. Section 1), Theorems 2.2 and 3.2 lead to the following corollary.

**Corollary 3.3.** *If either  $\kappa = \lambda = \mathfrak{c}$  and the cardinal  $\mathfrak{c}$  is regular or  $\kappa = \lambda = \aleph_1$ , then the following are equivalent:*

- (1) *There exist a set  $E \subseteq \mathbb{R}$  of cardinality  $\lambda$ , a non- $\sigma$ -compact Polish space  $Y$ , and a continuous function  $f : E \rightarrow Y$  which is not uniformly continuous (with respect to any complete metric on  $Y$ ) on any subset of  $E$  of cardinality  $\kappa$ .*
- (2) *There exist a set  $E \subseteq \mathbb{R}$  of cardinality  $\lambda$ , a compact metric space  $Y$  and a continuous function  $f : E \rightarrow Y$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ .*

#### 4. CONSTRUCTING $C_8$ -LIKE EXAMPLES FROM $\mathcal{K}$ -LUSIN SETS

Throughout this section we assume that  $G$  is an uncountable  $G_\delta$ -set in a compact metric space  $X$ .

Given a collection  $\mathcal{K}$  of compact sets in  $X$  containing all singletons, we say that an uncountable subset  $E$  of  $G$  is a  $\kappa$ - $\mathcal{K}$ -Lusin set in  $G$ , where  $\aleph_1 \leq \kappa \leq \mathfrak{c}$ , if  $|E \cap K| < \kappa$  for every set  $K \in \mathcal{K}$ . In particular, if  $\mathcal{K} = K(G)$ , the collection of all compact sets in  $G$ , then a  $\kappa$ - $\mathcal{K}$ -Lusin set in  $G$  is just a  $\kappa$ - $K$ -Lusin set in  $G$ .

The  $C_8$ -like examples, presented later in this section by means of  $\mathcal{K}$ -Lusin sets, are based on the following observation, applied also previously in the proofs of [11, Theorems 3.8 and 3.9].

**Proposition 4.1.** *Let  $\varphi : X \rightarrow Y$  be a continuous map onto a compact metric space  $Y$  such that  $\varphi|_G$  is a homeomorphism onto  $\varphi(G)$ . Let  $\mathcal{K}$  be a collection of compact sets in  $X$  such that whenever  $A \subseteq G$  and  $\varphi|_A$  extends to a homeomorphism over  $\bar{A}$ , the closure of  $A$  in  $X$  (equivalently, whenever  $\varphi$  is injective on  $\bar{A}$ ), then  $\bar{A} \in \mathcal{K}$ .*

*If  $T \subseteq G$  is a  $\kappa$ - $\mathcal{K}$ -Lusin set in  $G$  of cardinality  $\lambda$ , where  $\aleph_1 \leq \kappa \leq \lambda \leq \mathfrak{c}$ , then letting  $E = \varphi(T)$  and  $f = (\varphi|_G)^{-1}|_E : E \rightarrow X$ , we obtain a continuous function on a set of cardinality  $\lambda$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ .*

*Proof.* Aiming at a contradiction, assume that  $f|_B$  is uniformly continuous (with respect to any metric compatible with the topology of  $Y$ ) on a set  $B \subseteq E$  of cardinality  $\kappa$  and let  $A = f(B) = (\varphi|_G)^{-1}(B)$ . Then, since  $\varphi|_A : A \rightarrow B$  is also uniformly continuous, the function  $\varphi|_A$  extends to a homeomorphism over  $\bar{A}$  (cf. [2, Theorem 4.3.17]). Consequently,  $\bar{A} \in \mathcal{K}$ . This, however, is impossible, since on one hand we have  $|T \cap \bar{A}| < \kappa$ ,  $T$  being a  $\kappa$ - $\mathcal{K}$ -Lusin set in  $G$ , but on the other hand  $A \subseteq T \cap \bar{A}$  has cardinality  $\kappa$ .  $\square$

**4.1. Zero-dimensional spaces.** Throughout this subsection let us additionally assume that the (compact metric) space  $X$  is zero-dimensional.

A proof of the following fact is given in [10, Lemma 4.2] (it is based on an idea similar to that in [3, proof of Lemma 5.3]).

**Lemma 4.2.** *For any  $G_\delta$ -set  $G$  in  $X$  there is a continuous map  $\varphi : X \rightarrow Y$  onto a compact metric space  $Y$ , such that  $\varphi|_G$  is a homeomorphism onto  $\varphi(G)$ ,  $\varphi(X \setminus G) \cap \varphi(G) = \emptyset$  and the set  $Y \setminus \varphi(G)$  is countable.*

Various  $C_8$ -like examples could be constructed with the help of Lemma 4.2 and the following observation. By a  $\sigma$ -ideal on  $X$  we mean a collection  $\mathcal{I}$  of Borel sets in  $X$ , closed under taking Borel subsets and countable unions of elements of  $\mathcal{I}$ , and containing all singletons.

**Proposition 4.3.** *Let  $\mathcal{I}$  be a  $\sigma$ -ideal on  $X$  and let  $\mathcal{K}$  be the collection of all compact sets from  $\mathcal{I}$ .*

*If  $G$  is a  $G_\delta$  set in  $X$  such that  $G \in \mathcal{I}$  and  $\varphi : X \rightarrow Y$  is a continuous map described in Lemma 4.2, then  $\bar{A} \in \mathcal{K}$  for any  $A \subseteq G$  such that  $\varphi|_A$  extends to a homeomorphism onto  $\bar{A}$ .*

*Consequently, if  $T \subseteq G$  is a  $\kappa$ - $\mathcal{K}$ -Lusin set in  $G$  of cardinality  $\lambda$ , where  $\aleph_1 \leq \kappa \leq \lambda \leq \mathfrak{c}$ , then letting  $E = \varphi(T)$  and  $f = \varphi^{-1}|_E : E \rightarrow X$ , we obtain a continuous function on a set of cardinality  $\lambda$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\kappa$ .*

*Proof.* Let us fix  $A \subseteq G$  such that  $\varphi|_A$  extends to a homeomorphism  $\tilde{\varphi}$  between  $\bar{A}$  and  $\overline{\varphi(A)}$ . Then the set  $\bar{A} \setminus G = \tilde{\varphi}^{-1}(\overline{\varphi(A)} \setminus \varphi(G))$  is countable, so  $\bar{A} \subseteq G \cup (\bar{A} \setminus G) \in \mathcal{I}$ , hence  $\bar{A} \in \mathcal{K}$ .

The final assertion follows directly from Proposition 4.1. □

As the following proposition shows, the above observation could be applied to various natural  $\sigma$ -ideals. Let us recall that an uncountable set  $T$  in a Polish space  $Y$  is a *Lusin set* in  $Y$ , if  $|T \cap D| < \aleph_1$  for every closed nowhere-dense subset of  $Y$ .

**Proposition 4.4.** *Let  $\mathcal{I}$  be a  $\sigma$ -ideal on  $X$ , let  $\mathcal{K}$  be the collection of all compact sets from  $\mathcal{I}$  and let us assume that  $\mathcal{I}$  is not generated by  $\mathcal{K}$  (i.e., there is a set from  $\mathcal{I}$  which is not covered by any  $F_\sigma$ -set from  $\mathcal{I}$ ). Then, there exists a  $G_\delta$ -set in  $X$  such that  $G \in \mathcal{I}$  but no non-empty, relatively open set in  $G$  is covered by an  $F_\sigma$ -set from  $\mathcal{I}$ . Consequently, every Lusin set  $T$  in  $G$  is  $\aleph_1$ - $\mathcal{K}$ -Lusin in  $G$  hence, it gives rise to a  $C_8$ -example (as described in Proposition 4.3).*

*Proof.* Let  $B \in \mathcal{I}$  be (a Borel) set not covered by any  $F_\sigma$ -set from  $\mathcal{I}$ . Then the existence of a  $G_\delta$ -set in  $X$  such that  $G \subseteq B$  but  $G$  is not covered by any  $F_\sigma$ -set from  $\mathcal{I}$  follows from a theorem of Solecki (see [15]). By shrinking  $G$ , if necessary, we may assume that  $G$  has the desired properties.

It follows that if  $K \in \mathcal{K}$ , then  $G \cap K$  is meager in  $G$ , and, consequently,  $T \cap K$  is countable for any Lusin set  $T$  in  $G$ . □

**Remark 4.5.** *A typical example of the situation described in Propositions 4.3 and 4.4 is when  $X$  is a copy of the Cantor in  $\mathbb{R}$  of positive*



Lebesgue measure,  $\mathcal{I}$  is the  $\sigma$ -ideal of Lebesgue measure zero Borel sets in  $X$  (then  $\mathcal{K}$  is the family of closed Lebesgue measure zero sets in  $X$ ) and  $G$  is a dense copy of irrationals in  $X$  of Lebesgue measure zero.

Then the function  $f = \varphi^{-1}|_H : H \rightarrow \mathbb{R}$ , where  $H = \varphi(G)$ , is a homeomorphic embedding of  $H$ , a copy of irrationals, to  $\mathbb{R}$ , with the property that for every Lusin set  $L$  of cardinality  $\mathfrak{c}$  in  $H$ , the function  $f|_L$  is not uniformly continuous on any uncountable subset of  $L$ . This provides an alternative proof of the theorem of Sierpiński that the existence of a Lusin set of cardinality  $\mathfrak{c}$  in  $\mathbb{P}$  implies  $C_8$  (cf. [14, proof of Théorème 6 on page 45]).

**4.2. Infinite-dimensional spaces.** The assumption that the space  $X$  is zero-dimensional in Theorem 3.1 is essential, as demonstrated by the following result (CH in this theorem can be weakened to the assumption that no family of less than  $\mathfrak{c}$  meager sets covers  $\mathbb{R}$ , cf. Remark 4.8).

**Theorem 4.6.** *Assuming CH, there exists a set  $E \subseteq I^{\mathbb{N}}$  of cardinality  $\mathfrak{c}$  such that*

- (1) *there is a continuous function  $f : E \rightarrow I^{\mathbb{N}}$ , which is not uniformly continuous on any subset of  $E$  of cardinality  $\mathfrak{c}$ ,*
- (2) *each continuous function  $g : E \rightarrow \mathbb{R}$  is constant on a subset of  $E$  of cardinality  $\mathfrak{c}$ .*

A key element of the proof of this theorem is a *Henderson compactum* – a compact metrizable infinite-dimensional space whose all compact subsets of finite dimension are zero-dimensional, cf. [9].

More specifically, we shall need the following fact, where *punctiform* sets are the sets without non-trivial subcontinua, cf. [3, 1.4.3].

**Lemma 4.7.** *There exists a non-empty punctiform  $G_\delta$ -set  $M$  in a Henderson compactum  $H \subseteq I^{\mathbb{N}}$  such that for each  $M' \subseteq M$  with  $\dim(M \setminus M') \leq 0$ , every continuous function  $u : M' \rightarrow \mathbb{R}$  is constant on a set of positive dimension.*

A justification of this lemma is rather standard, but since we did not find convenient references, we shall give a proof to this effect at the end of this subsection. For now, taking this fact for granted, we shall proceed with a proof of our main result.

*Proof of Theorem 4.6.* Let us adopt the notation of Lemma 4.7, and let  $\mathcal{K}$  be the collection of all compact zero-dimensional sets in the Henderson compactum  $H$ .

Using CH, we inductively construct a  $\mathfrak{c}$ - $\mathcal{K}$ -Lusin set  $T$  in  $M$  such that  $|T \cap L| = \mathfrak{c}$  for every Borel set  $L$  in  $M$  with  $\dim L > 0$ . To that end, we list all elements of  $\mathcal{K}$  as  $\langle K_\alpha : \alpha < \mathfrak{c} \rangle$  and all Borel sets  $L$  in  $M$  with  $\dim L > 0$  as  $\langle L_\alpha : \alpha < \mathfrak{c} \rangle$ , repeating each such set  $L$  continuum

many times. Then, we subsequently pick for  $\xi < \mathfrak{c}$

$$p_\xi \in L_\xi \setminus \left( \bigcup_{\alpha < \xi} K_\alpha \cup \{p_\alpha : \alpha < \xi\} \right),$$

using the fact that, under CH, the set in brackets is zero-dimensional, by the sum theorem for dimension zero (cf. [3, 1.3.1]). Finally, we let

$$T = \{p_\xi : \xi < \mathfrak{c}\}.$$

In particular,  $T$  meets each Borel set in  $M$  of positive dimension in continuum many points.

**Claim 1.** Each continuous function  $w : T \rightarrow \mathbb{R}$  is constant on a set of cardinality  $\mathfrak{c}$ .

Indeed,  $w$  can be extended to a continuous function  $\tilde{w} : \tilde{T} \rightarrow \mathbb{R}$  over a  $G_\delta$ -set  $\tilde{T}$  in  $M$  containing  $T$ . The Borel set  $M \setminus \tilde{T}$  is disjoint from  $T$ , hence  $\dim(M \setminus \tilde{T}) \leq 0$ . Setting in Lemma 4.7  $u = \tilde{w}$  and  $M' = \tilde{T}$ , we get  $r \in \mathbb{R}$  such that  $\dim u^{-1}(r) > 0$ . Being a Borel set in  $M$ ,  $u^{-1}(r)$  intersects  $T$  in a set of cardinality  $\mathfrak{c}$ , i.e.,  $|w^{-1}(r)| = \mathfrak{c}$ .

Let us now apply a counterpart of Lemma 4.2 for infinite-dimensional spaces to the following effect (cf. the proof of [3, Lemma 5.3.1]): there exists a continuous surjection  $\varphi : H \rightarrow Y$  onto a compact metrizable space  $Y$  such that  $\varphi|_M$  is a homeomorphism onto  $\varphi(M)$ ,  $\varphi(H \setminus M) \cap \varphi(M) = \emptyset$  and  $Y \setminus \varphi(M)$  is a countable union of finite-dimensional compact sets.

**Claim 2.** The set  $E = \varphi(T)$  satisfies the assertion of Theorem 4.6.

Since  $E$  is homeomorphic to  $T$ , Claim 1 shows that (2) in Theorem 4.6 is satisfied.

To check also assertion (1) in this theorem, we shall make sure that the function

$$f = \varphi^{-1}|_E : E \rightarrow H$$

is not uniformly continuous on any set of cardinality  $\mathfrak{c}$ .

Since  $T$  is a  $\mathfrak{c}$ - $\mathcal{K}$ -Lusin set in  $M$ , by Proposition 4.1 it is enough to verify that for any non-empty  $A \subseteq M$ , whenever  $\varphi|_{\bar{A}}$  is an embedding (the closure of  $A$  is in  $H$ ), then  $\bar{A} \in \mathcal{K}$ , i.e.,  $\dim \bar{A} = 0$ .

To check this, let us note that

$$\varphi(\bar{A} \setminus M) = \varphi(\bar{A}) \setminus \varphi(M) \subseteq Y \setminus \varphi(M)$$

is a countable union of compact finite-dimensional sets, and so is  $\bar{A} \setminus M$ . Since  $H$  is a Henderson compactum,  $\bar{A} \setminus M$  is a countable union of zero-dimensional compact sets, hence  $\dim(\bar{A} \setminus M) = 0$  by the sum theorem (cf. [3, 1.4.5]). By the enlargement theorem, cf. [3, 1.5.11],  $\bar{A} \setminus M$  can be enlarged to a zero-dimensional  $G_\delta$ -set  $L$  in  $H$ . The set  $\bar{A} \setminus L$  is a  $\sigma$ -compact subset in the punctiform space  $M$ , hence again by the sum theorem,  $\dim(\bar{A} \setminus L) \leq 0$  (cf. [3, 1.4.5]). By the addition theorem,

cf. [3, 1.5.10],  $\dim \bar{A} \leq 1$ , and  $H$  being a Henderson compactum,  $\dim \bar{A} = 0$ .

□

In the assumptions of Theorem 4.6, CH can be weakened to the assertion (usually denoted by  $\text{cov}(\mathcal{M}) = \mathfrak{c}$ , cf. [5]) that no family of less than  $\mathfrak{c}$  meager sets covers  $\mathbb{R}$ . The only change in the proof requires checking that the inductive definition of the sequence  $\langle p_\xi : \xi < \mathfrak{c} \rangle$  is correct. More precisely, the following is true.

**Remark 4.8.** *Assuming  $\text{cov}(\mathcal{M}) = \mathfrak{c}$ , the union of any family  $\mathcal{K}$  of compact zero-dimensional sets in  $I^\mathbb{N}$ , with  $|\mathcal{K}| < \mathfrak{c}$ , is zero-dimensional.*

*Proof.* Let  $\mathcal{K}$  be a family of compact zero-dimensional sets in  $I^\mathbb{N}$ , with  $|\mathcal{K}| < \mathfrak{c}$ . For any  $K \in \mathcal{K}$ , let us consider

$$(1) \mathcal{U}_K = \{f \in C(I^\mathbb{N}, \mathbb{R}) : f^{-1}(0) \cap K = \emptyset\}.$$

Then  $\mathcal{U}_K$  is open and dense in the function space  $C(I^\mathbb{N}, \mathbb{R})$  and this space being perfect Polish, the assumption  $\text{cov}(\mathcal{M}) = \mathfrak{c}$  guarantees that the set

$$(2) \mathcal{G} = \bigcap \{\mathcal{U}_K : K \in \mathcal{K}\}$$

is dense in  $C(I^\mathbb{N}, \mathbb{R})$  (cf. [6, 8.32]).

Let us recall (cf. [3]) that a closed set  $L$  in a topological space  $Z$  separates the space  $Z$  between sets  $A_1, A_2 \subseteq Z$ , if  $Z \setminus L = U_0 \cup U_1$ , where  $U_0, U_1$  are open, disjoint and  $A_i \subseteq U_i$ , for  $i = 0, 1$ . If  $A_1$  and  $A_2$  are singletons, then we say that  $L$  separates  $Z$  between the respective points.

Let  $F = \bigcup \mathcal{K}$ . To prove that  $\dim F = 0$ , it is enough to show that for any pair of disjoint compact sets  $A, B$  in  $I^\mathbb{N}$ ,  $I^\mathbb{N}$  can be separated between  $A$  and  $B$  by a closed set disjoint from  $F$ . Indeed, let  $x \in F$  and let  $U$  be a relatively open subset of  $F$  with  $x \in U$  and  $F \setminus U \neq \emptyset$ . Then  $U = V \cap F$  for an open  $V$  in  $I^\mathbb{N}$  with  $B = I^\mathbb{N} \setminus V \neq \emptyset$ , and separating  $I^\mathbb{N}$  between  $A = \{x\}$  and  $B$  by a closed set in  $I^\mathbb{N}$ , disjoint from  $F$ , leads to disjoint open sets  $V_1, V_2$  in  $I^\mathbb{N}$  with  $x \in V_1 \cap F \subseteq U$ ,  $V_2 \cap F \neq \emptyset$ , and  $F = (V_1 \cap F) \cup (V_2 \cap F)$ . Thus,  $V_1 \cap F$  is a relatively clopen in  $F$  neighbourhood of  $x$  contained in  $U$ .

So let  $A, B$  be disjoint compact sets in  $I^\mathbb{N}$ . We can pick  $f \in \mathcal{G}$  such that  $f(A) \subseteq (-\infty, 0)$ ,  $f(B) \subseteq (0, +\infty)$ , as such functions form an open, non-empty set in  $C(I^\mathbb{N}, \mathbb{R})$ . Then  $L = f^{-1}(0)$  is a closed set in  $I^\mathbb{N}$  separating  $I^\mathbb{N}$  between  $A$  and  $B$  with the property that  $L \cap K = \emptyset$  for all  $K \in \mathcal{K}$ , cf. (1) and (2).

□

*Proof of Lemma 4.7.* Let us fix a Henderson continuum  $K$  in  $I^\mathbb{N}$ . We shall consider on  $I^\mathbb{N}$  the metric assigning to points  $(s_0, s_1, \dots), (t_0, t_1, \dots)$  the distance  $\sum_i 2^{-i} |s_i - t_i|$ .

**Claim 1.** There exists a punctiform  $G_\delta$ -set  $S$  in  $K$  and distinct points  $a, b \in S$  such that each relatively closed set in  $S$  separating  $S$  between  $a$  and  $b$  has dimension at least 2.

Indeed, since  $\dim K \geq 4$ ,  $K$  contains a punctiform  $G_\delta$ -set  $W$  with  $\dim W \geq 3$ .

This theorem goes back to Mazurkiewicz, and can be justified by the reasoning in the proof of [9, Theorem 3.9.3] applied to a pair of disjoint compact sets  $A$  and  $B$  in  $K$  such that for any set  $N$  in  $K$  with  $\dim N \leq 2$  there is a continuum in  $K$  joining  $A$  and  $B$  and missing  $N$  (cf. [12, Theorem 4.2]), and to a continuous map  $\pi : K \rightarrow [-1, 1]$  sending  $A$  to  $-1$  and  $B$  to  $1$ .

Now, since  $\dim W \geq 3$ , there is  $a \in W$  such that all sufficiently small neighbourhoods of  $a$  in  $W$  have boundaries of dimension  $\geq 2$ . An argument in [8, proof of Theorem 8, p. 172] gives a point  $b \in K \setminus \{a\}$  such that for  $S = W \cup \{b\}$ , every relatively closed set in  $S$  separating  $S$  between  $a$  and  $b$  has dimension  $\geq 2$ . Since  $S$  is a punctiform  $G_\delta$ -set in  $K$ , it satisfies Claim 1.

**Claim 2.** There are embeddings  $h_n : I^\mathbb{N} \rightarrow I^\mathbb{N}$ ,  $n = 1, 2, \dots$ , such that, letting

$$\begin{aligned} M_0 &= S \times \{0\}, \quad M_n = h_n(S) \times \{\tfrac{1}{n}\} \text{ for } n = 1, 2, \dots, \\ M &= \bigcup_{n=0}^{\infty} M_n \subseteq I^\mathbb{N} \times I \text{ and } H = \overline{M} - \text{the closure of } M \text{ in } I^\mathbb{N} \times I, \end{aligned}$$

one obtains sets  $M \subseteq H$  satisfying the assertion of Lemma 4.7.

We adopt the notation from Claim 1. Since  $K$  is a compact, connected set in  $I^\mathbb{N}$ , we can find open connected neighbourhoods  $U_1 \supseteq U_2 \supseteq \dots$  of  $K$  such that each open neighbourhood of  $K$  in  $I^\mathbb{N}$  contains some  $U_n$ .

Let  $D$  be a countable set dense in  $S$ , and let  $\langle (c_n, d_n) : n \in \mathbb{N} \rangle$  be an enumeration of all ordered pairs of distinct points from  $D$ , such that each such pair appears in the sequence infinitely many times.

Let us fix  $n > 0$ . We shall define an embedding  $h_n : I^\mathbb{N} \rightarrow U_n$  such that the distance from  $h_n(a)$  to  $c_n$  and the distance from  $h_n(b)$  to  $d_n$  is less than  $\frac{1}{n}$ .

To that end, let us notice that  $U_n$  is arcwise connected, being an open, connected set in  $I^\mathbb{N}$  (cf. [16, Proposition 12.25]), and hence there is a continuous  $f : I^\mathbb{N} \rightarrow U_n$  with  $f(a) = c_n$ ,  $f(b) = d_n$ . Let  $m \geq n$  be large enough to ensure that for any  $x, y \in I^\mathbb{N}$ , whenever the first  $m$  coordinates of  $x$  and  $f(y)$  coincide, then  $x \in U_n$ . Now, denoting by  $p_j : I^\mathbb{N} \rightarrow I$  the projection onto  $j$ 'th coordinate, we define the embedding  $h_n$  by  $p_i(h_n(x)) = p_i(f(x))$  for  $i \leq m$  and  $p_{m+i}(h_n(x)) = p_i(x)$  for  $i = 1, 2, \dots$ .

Having defined the embeddings  $h_n$ , we shall check that the set  $M$  in Claim 2 satisfies the assertion of Lemma 4.7.

Clearly,  $M$  is a punctiform  $G_\delta$ -set in  $I^\mathbb{N} \times I$ .

Since each neighbourhood of  $K$  in  $I^\mathbb{N}$  contains all but finitely many  $h_n(K) \subseteq \overline{U_n}$ , the set

$$K \times \{0\} \cup \bigcup_{n=1}^{\infty} \left( h_n(K) \times \left\{ \frac{1}{n} \right\} \right)$$

contains  $H = \overline{M}$ , and since it is clearly a Henderson compactum containing  $M$ , so is  $H$ .

Let  $M' \subseteq M$  satisfy  $\dim(M \setminus M') \leq 0$  and let  $u : M' \rightarrow \mathbb{R}$  be continuous.

Since  $\dim(M_0) = \dim(S) \geq 2$ , the addition theorem yields that  $\dim(M' \cap M_0) \geq 1$ . Therefore, if  $u$  is constant on  $M' \cap M_0$ , we are done.

Let us assume that this is not the case. Then for some  $x, y \in M' \cap M_0$ ,  $u(x) \neq u(y)$ , hence for  $r = \frac{u(x)+u(y)}{2}$ ,  $u^{-1}(r)$  is a relatively closed set in  $M'$  which separates  $M'$  between  $x$  and  $y$ . One can extend  $u^{-1}(r)$  to a closed set  $L$  in  $H$  such that  $L$  separates  $H$  between  $x$  and  $y$ , cf. [7, §21, XI, Theorem 2, p. 226]. Let us pick  $c, d \in D$ , close enough to  $x$  and  $y$ , respectively, so that  $L$  separates  $H$  between  $(c, 0)$  and  $(d, 0)$ , and using the fact that  $(c, d) = (c_n, d_n)$  for infinitely many  $n$ 's and the distances from  $h_n(a)$  to  $c_n$ , and from  $h_n(b)$  to  $d_n$  tend to zero, let us pick  $n$  such that  $L$  also separates  $H$  between  $(h_n(a), \frac{1}{n})$  and  $(h_n(b), \frac{1}{n})$ . In effect, since  $x \mapsto (h_n(x), \frac{1}{n})$  is a homeomorphism of  $S$  onto  $M_n$ , and each relatively closed set separating  $S$  between  $a$  and  $b$  has dimension  $\geq 2$ , we infer that  $\dim(M_n \cap L) \geq 2$ . Since  $\dim(M_n \setminus M') \leq 0$ , applying again the addition theorem, we conclude that  $\dim(M' \cap L) \geq 1$ .

Finally, since  $M' \cap L = u^{-1}(r)$ ,  $u$  is constant on a set of positive dimension.

□

## 5. A CHARACTERIZATION OF COMPLETE METRIZABILITY

A key step in proving that statement  $C_9(\lambda, \kappa)$  is equivalent to the existence of a  $\kappa$ -K-Lusin set  $E$  of cardinality  $\lambda$  in  $\mathbb{P}$  (cf. [11, Theorem 2.3]) is Theorem 2.1 in [11] (closely related to a theorem of Sierpiński, cf. [11, Remark 2.2]) stating that for any Polish space  $X$  there is a sequence  $f_0 \geq f_1 \geq \dots$  of continuous functions  $f_n : X \rightarrow I$  which converges to zero pointwise but does not converge uniformly on any set with non-compact closure in  $X$ .

In fact, the existence of such a function sequence characterizes complete metrizability of a separable metrizable space  $X$ .

**Theorem 5.1.** *A separable metrizable space  $X$  is completely metrizable if and only if there is a sequence  $f_0 \geq f_1 \geq \dots$  of continuous functions  $f_n : X \rightarrow I$  which converges to zero pointwise but does not converge uniformly on any set with non-compact closure in  $X$ .*

*Proof.* In view of [11, Theorem 2.1] we only have to prove the "if" part of the above equivalence. So let  $(X, d)$  be a separable metric space. Let  $f_n : X \rightarrow I$ ,  $n \in \mathbb{N}$ , be a sequence of continuous functions with  $f_0 \geq f_1 \geq \dots$ , which converges to zero pointwise but does not converge uniformly on any set with non-compact closure in  $X$ .

Let  $(\hat{X}, \hat{d})$  be the completion of  $(X, d)$ ; clearly, the space  $\hat{X}$  is Polish. There exists a  $G_\delta$ -set  $\tilde{X}$  in  $\hat{X}$ , containing  $X$  such that each  $f_n$  extends to a continuous function  $\tilde{f}_n : \tilde{X} \rightarrow I$ . Since  $X$  is dense in  $\tilde{X}$ , we have  $\tilde{f}_0 \geq \tilde{f}_1 \geq \dots$  and in particular, the set

$$G = \{x \in \tilde{X} : \lim_{n \rightarrow \infty} \tilde{f}_n(x) = 0\}$$

is  $G_\delta$  in  $\tilde{X}$ .

Since  $X \subseteq G$ , it remains to make sure that  $G \subseteq X$ , to conclude that  $X = G$  is completely metrizable as a  $G_\delta$ -subset of the (Polish) space  $\hat{X}$ .

So let  $c \in G$  and let us pick  $x_k \in X$ ,  $k \in \mathbb{N}$ , such that  $\lim_{k \rightarrow \infty} x_k = c$ .

Let  $K = \{c\} \cup \{x_k : k \in \mathbb{N}\}$ . Then  $K$  is compact, so the sequence  $(\tilde{f}_n)_{n \in \mathbb{N}}$  converges uniformly on  $K$  (this is a special instance of the Dini theorem [2, Lemma 3.2.18]). It follows that the sequence  $(f_n)_{n \in \mathbb{N}}$  converges uniformly on  $K \cap X$ , a closed set in  $X$  which therefore, by the assumed property of  $(f_n)_{n \in \mathbb{N}}$ , is compact. Since the sequence  $(x_n)_{n \in \mathbb{N}}$  converges to  $c$  and all  $x_n$  are elements of  $K \cap X$ , so is  $c$ . In particular,  $c \in X$ . □

With the help of Theorem 5.1 we shall establish the following more general result (for terminology see [2]).

**Theorem 5.2.** *Let  $X$  be a Hausdorff space. Then  $X$  is a Čech-complete Lindelöf space if and only if there is a sequence  $f_0 \geq f_1 \geq \dots$  of continuous functions  $f_n : X \rightarrow I$  converging pointwise to zero but not converging uniformly on any closed non-compact set in  $X$ .*

*Proof.* First, let  $X$  be a Čech-complete Lindelöf space. By a theorem of Frolík, it follows that there is a perfect map  $p : X \rightarrow Y$  onto a Polish space  $Y$ , cf. [2, 5.5.9]. Let us recall that for a Hausdorff space  $X$  and a metrizable space  $Y$  this means, cf. [2, Theorems 3.7.2 and 3.7.18], that  $p$  is a continuous, closed mapping and the inverse image of every compact subset of  $Y$  is compact. It is straightforward to check that if functions  $f_n : Y \rightarrow I$ ,  $n \in \mathbb{N}$ , with  $f_0 \geq f_1 \geq \dots$ , satisfy the assertion of Theorem 5.1, the functions  $f_n \circ p : X \rightarrow I$ ,  $n \in \mathbb{N}$ , have the required properties.

Next, given a sequence  $f_0 \geq f_1 \geq \dots$  of functions  $f_n : X \rightarrow I$ , described in the theorem, let us consider the diagonal map

$$(1) \ F = (f_0, f_1, \dots) : X \rightarrow I^{\mathbb{N}} \text{ and let } Y = F(X).$$

Let  $p_n : I^{\mathbb{N}} \rightarrow I$  denote the projection onto the  $n$ 'th coordinate and let, cf. (1), for  $n \in \mathbb{N}$ ,

$$(2) \quad g_n = p_n|_Y : Y \rightarrow I.$$

Note that, since  $f_n = g_n \circ F$ , the properties of the sequence  $(f_n)_{n \in \mathbb{N}}$  are passed to the sequence  $(g_n)_{n \in \mathbb{N}}$ , namely:

$$(3) \quad g_0 \geq g_1 \geq \dots \text{ is a sequence of continuous functions converging pointwise to zero,}$$

and for any closed set  $A$  in  $Y$ ,

$$(4) \quad \text{if } (g_n)_{n \in \mathbb{N}} \text{ converges on } A \text{ uniformly, then } A \text{ is compact.}$$

To see (4), let us consider  $B = F^{-1}(A)$ . Then, as  $f_n = g_n \circ F$ ,  $(f_n)_{n \in \mathbb{N}}$  converges on  $B$  uniformly, hence  $B$  is compact, and so is  $A = F(B)$ .

By Theorem 5.1,

$$(5) \quad Y \text{ is a } G_\delta\text{-set in } I^{\mathbb{N}}.$$

To complete the proof, it is enough to show that

$$(6) \quad \text{the mapping } F \text{ is perfect.}$$

Indeed, if (6) holds, then since by (5),  $Y$  is a Čech-complete Lindelöf space, so is its perfect preimage  $X$ , cf. [2, Theorems 3.8.9 and 3.9.10].

To prove (6), it suffices to check that for every compact  $K \subseteq Y$  the inverse image  $F^{-1}(K)$  is compact, cf. [2, Theorem 3.7.18].

So let  $K \subseteq Y$  be compact. Then, again by the Dini theorem, the sequence  $(g_n)_{n \in \mathbb{N}}$  converges uniformly on  $K$ , and since  $f_n = g_n \circ F$ , cf. (2), the sequence  $(f_n)_{n \in \mathbb{N}}$  converges uniformly on  $F^{-1}(K)$ . By the assumed property of this sequence,  $F^{-1}(K)$  is compact, completing the proof. □

## REFERENCES

1. T. Bartoszyński, L. J. Halbeisen, *On a theorem of Banach and Kuratowski and  $K$ -Lusin sets* Rocky Mt. J. Math. **33** (2003), 1223—1231
2. R. Engelking, *General Topology*, Sigma Series in Pure Mathematics, Heldermann Verlag, 1989.
3. R. Engelking, *Theory of Dimensions Finite and Infinite*, Heldermann Verlag, 1995.
4. L. J. Halbeisen, *Combinatorial Set Theory, With a Gentle Introduction to Forcing*, Springer Monographs in Mathematics, Springer-Verlag, 2012.
5. W. Just, M. Weese, *Discovering modern set theory. II*, (Graduate studies in mathematics, AMS, 1997.
6. A. S. Kechris, *Classical descriptive set theory*, Graduate Texts in Math. 156, Springer-Verlag, 1995.
7. K. Kuratowski, *Topology*, Vol. I, Academic Press 1966.
8. K. Kuratowski, *Topology*, Vol. II, Academic Press 1966.
9. J. van Mill, *The Infinite-Dimensional Topology of Function Spaces*, Elsevier 2001.
10. R. Pol, P. Zakrzewski, *On Mazurkiewicz's sets, thin sigma-ideals of compact sets and the space of probability measures on the rationals*, RACSAM **115**(42) (2021).

11. R. Pol, P. Zakrzewski, *On two consequences of CH established by Sierpiński*, Archive for Mathematical Logic (2024), <https://doi.org/10.1007/s00153-024-00925-6>.
12. L.R. Rubin, R.M. Schori, J.J. Walsh, *New dimension-theory techniques for constructing infinite-dimensional examples*, General Topology Appl. **10** (1979), 93–102.
13. W. Sierpiński, *Sur une relation entre deux conséquences de l'hypothèse du continu*, Fund. Math. **31** (1938), 227–230.
14. W. Sierpiński *Hypothèse du continu*, Monografie Matematyczne vol IV, 1934.
15. S. Solecki, *Covering analytic sets by families of closed sets*, Journal of Symbolic Logic **59(3)** (1994) 1022–1031.
16. W.A. Sutherland, *Introduction to Metric and Topological Spaces*, Second Edition, Oxford University Press, 2009.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, UL. BANACHA 2,  
02-097 WARSAW, POLAND

*E-mail address:* pol@mimuw.edu.pl, piotrzak@mimuw.edu.pl